

Practice paper (calculator 2): full worked solutions

1 $85\% = \text{£}258$

$$1\% = \frac{\text{£}258}{85}$$

$$100\% = \frac{\text{£}258}{85} \times 100$$

$$= \text{£}303.53$$

2 $(0.45 \times 0.78)^2 \approx (0.5 \times 0.8)^2 = 0.4^2 = 0.16$

3 a $\frac{2x-5}{11} = 3$

$$2x - 5 = 33$$

$$2x = 38$$

$$x = 19$$

b $x^2 - x - 42 = 0$

$$(x + 6)(x - 7) = 0$$

$$x = -6 \text{ or } 7$$

4 a gradient $= \frac{k-2}{-2-3} = \frac{k-2}{-5}$

$$\frac{k-2}{-5} = \frac{4}{5}$$

$$5(k-2) = -20$$

$$5k - 10 = -20$$

$$5k = -10$$

$$k = -2$$

b $y - 2 = \frac{4}{5}(x - 3)$

$$5y - 10 = 4x - 12$$

$$5y = 4x - 2 \text{ or } y = \frac{4}{5}x - \frac{2}{5}$$

5 $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$ gives:

$$= \frac{3 \pm \sqrt{(-3)^2 - 4 \times 1 \times -6}}{2 \times 1}$$

$$= \frac{3 \pm \sqrt{9 + 24}}{2}$$

$$= \frac{3 \pm \sqrt{33}}{2}$$

$$x = \frac{3 + \sqrt{33}}{2} \text{ or } \frac{3 - \sqrt{33}}{2}$$

$$x = 4.37 \text{ or } -1.37 \text{ (to 2 d.p.)}$$

6 a Completing the square:

$$y = x^2 - 2x - 3 = (x-1)^2 - 1 - 3 = (x-1)^2 - 4$$

Coordinates of turning point are (1, -4).

b To find where the curve intersects the x -axis:


$$x^2 - 2x - 3 = 0$$

$$(x-3)(x+1) = 0$$

$$x = 3 \text{ or } -1$$

To find where the curve intersects the y -axis:

$$\text{when } x = 0, y = (0)^2 - 2 \times 0 - 3 = -3$$



7 a Equating expressions for y :

$$10x^2 - 5x - 2 = 2x - 3$$

$$10x^2 - 7x + 1 = 0$$

$$(5x - 1)(2x - 1) = 0$$

$$x = \frac{1}{5} \text{ or } x = \frac{1}{2}$$

$$\text{Substituting } x = \frac{1}{5} \text{ into } y = 2x - 3 \text{ gives } y = -2\frac{3}{5}$$

$$\text{Substituting } x = \frac{1}{2} \text{ into } y = 2x - 3 \text{ gives } y = -2$$

$$x = \frac{1}{5} \text{ and } y = -2\frac{3}{5} \text{ or } x = \frac{1}{2} \text{ and } y = -2$$

b the two points where the curve and line intersect

8 $\frac{3\sqrt{3} - \sqrt{2}}{\sqrt{3} - \sqrt{2}} = \frac{(3\sqrt{3} - \sqrt{2})(\sqrt{3} + \sqrt{2})}{(\sqrt{3} - \sqrt{2})(\sqrt{3} + \sqrt{2})}$


$$= \frac{9 + 3\sqrt{6} - \sqrt{6} - 2}{3 + \sqrt{6} - \sqrt{6} - 2} = \frac{7 + 2\sqrt{6}}{1} = 7 + 2\sqrt{6}$$

9 $y = f(x)$ to $y = f(x - 1) + 1$ is a translation of 1 unit to the right and 1 unit up (i.e. a translation of $\begin{pmatrix} 1 \\ 1 \end{pmatrix}$):

(1, 0) will become (2, 1)

(5, 0) will become (6, 1)

(3, 4) will become (4, 5)



10 a $f(-2) = 5 \times (-2)^2 + 4 = 24$

b Let $y = 5x^2 + 4$

$$\frac{y-4}{5} = x^2$$

$$x = \pm \sqrt{\frac{y-4}{5}}$$

$$f^{-1}(x) = \pm \sqrt{\frac{x-4}{5}}$$

c $fg(x) = f(x + 1)$

$$= 5(x + 1)^2 + 4$$

11 a Angles VRP and VRQ are right angles because V is directly above R .


Using Pythagoras' theorem:

$$20^2 = VR^2 + 16^2$$

$$400 = VR^2 + 256$$

$$VR^2 = 144$$

$$VR = 12 \text{ cm}$$




Using Pythagoras' theorem:

$$20^2 = VS^2 + (24 \div 2)^2$$

$$400 = VS^2 + 144$$

$$VS^2 = 256$$

$$VS = 16 \text{ cm}$$



Using Pythagoras' theorem:

$$16^2 = RS^2 + 12^2$$

$$256 = RS^2 + 144$$

$$RS^2 = 112$$

$$RS = 10.6 \text{ cm (to 3 s.f.)}$$

c $\cos \theta = \frac{10.6}{16}$

$$\theta = \cos^{-1}\left(\frac{10.6}{16}\right)$$

$$= 48.5^\circ \text{ (to 3 s.f.)}$$

12 Let Amy's age = x years.

Amy's mother's age = $3x$ years.

In 12 years time, Amy will be $x + 12$ and her mother will be $3x + 12$

$$x + 12 = \frac{3x + 12}{2}$$

$$2x + 24 = 3x + 12$$

$$x = 12$$

Amy's mother is 36 years old.

13 a $l = \frac{\theta}{360} \times 2\pi r = \frac{35}{360} \times 2\pi \times 8 = 4.89 \text{ cm (to 2 d.p.)}$

b $A = \frac{\theta}{360} \pi r^2 = \frac{35}{360} \pi \times 8^2 = 19.55 \text{ cm}^2 \text{ (to 2 d.p.)}$

14 $\frac{1}{3x^2 + 5x - 2} \div \frac{1}{9x^2 - 1} = \frac{1}{3x^2 + 5x - 2} \times \frac{9x^2 - 1}{1}$

$$= \frac{1}{(3x - 1)(x + 2)} \times (3x - 1)(3x + 1)$$

$$= \frac{3x + 1}{x + 2}$$

$$a = 3, b = 1, c = 1, d = 2$$

15 multiples of 12: 12, 24, 36, 48, 60, 72, 84, 96, 108, 120, 132, 144 ...

multiples of 16: 16, 32, 48, 64, 80, 96, 112, 128, 144 ...

multiples of 18: 18, 36, 54, 72, 90, 108, 126, 144 ...

a The first number to appear in 2 lists is 36.

36 days

b The first number to appear in all 3 lists is 144.

144 days

16 a i $\vec{AM} = \vec{AO} + \frac{1}{2}\vec{OB}$

$$= -\mathbf{a} + \frac{\mathbf{b}}{2}$$

$$= \frac{\mathbf{b}}{2} - \mathbf{a}$$

ii $\vec{AR} = \frac{2}{3}\vec{AM}$

$$= \frac{2}{3}\left(\frac{\mathbf{b}}{2} - \mathbf{a}\right)$$

$$= \frac{\mathbf{b}}{3} - \frac{2\mathbf{a}}{3}$$

b $\vec{BP} = \vec{BO} + \vec{OP} = -\mathbf{b} + \frac{\mathbf{a}}{2} = \frac{\mathbf{a}}{2} - \mathbf{b} = \frac{1}{2}(\mathbf{a} - 2\mathbf{b})$

$$\vec{BR} = \vec{BA} + \vec{AR} = \mathbf{a} - \mathbf{b} + \frac{\mathbf{b}}{3} - \frac{2\mathbf{a}}{3} = \frac{\mathbf{a}}{3} - \frac{2\mathbf{b}}{3} = \frac{1}{3}(\mathbf{a} - 2\mathbf{b})$$

\vec{BP} and \vec{BR} have the same vector part $(\mathbf{a} - 2\mathbf{b})$ and are therefore parallel. As they both pass through point B , points P and R lie on the same straight line.

17 $2.535 \leq T < 2.545$

$$9.75 \leq g < 9.85$$

Upper bound for h will be when T has its upper bound and g has its upper bound.

$$h = \frac{gT^2}{2} = \frac{9.8 \times 2.545^2}{2} = 31.73742 = 31.74 \text{ m (to 4 s.f.)}$$

Lower bound for h will be when T has its lower bound and g has its lower bound.

$$h = \frac{gT^2}{2} = \frac{9.85 \times 2.545^2}{2} = 31.8993 = 31.89 \text{ m (to 4 s.f.)}$$

The upper bound (31.74) and the lower bound (31.89) are the same when the numbers are given to 2 significant figures.

$$h = 32 \text{ m (to 2 s.f.)}$$

18 a Let $f(x) = x^3 - 4x + 2$

$$f(1) = 1^3 - 4(1) + 2 = -1$$

$$f(0) = 0^3 - 4(0) + 2 = 2$$

As there is a sign change, the root lies between 0 and 1.

b $x^3 - 4x + 2 = 0$

$$4x = x^3 + 2$$

$$x = \frac{x^3 + 2}{4}$$

$$x = \frac{x^3}{4} + \frac{1}{2}$$

c $x_1 = 0.53125$

$$x_2 = 0.53748$$

$$x_3 = 0.53881$$

$$x_4 = 0.53911 \approx 1.539 \text{ (to 3 d.p.)}$$

19 a 9 parts = 36 so 1 part = 4

$$\text{number of red counters} = 4 \times 4 = 16$$

b i $P(\text{1st counter is red}) = \frac{16}{36} = \frac{4}{9}$

$$P(\text{2nd counter is red}) = \frac{15}{35} = \frac{3}{7}$$

$$P(\text{both counters are red}) = \frac{4}{9} \times \frac{3}{7} = \frac{12}{63} = \frac{4}{21}$$

ii $P(\text{different colours}) = P(\text{red then blue}) + P(\text{blue then red})$

$$= \left(\frac{4}{9} \times \frac{3}{7}\right) + \left(\frac{5}{9} \times \frac{19}{35}\right)$$

$$= \frac{31}{63}$$