

# Edexcel Foundation Mathematics Revision Guide

## Full worked solutions


### Number

#### Factors, multiples and primes

1 a 5      b 1, 12      c 1, 5, 45

2  $70 = 2 \times 5 \times 7$   
 $150 = 2 \times 3 \times 5 \times 5$   
 HCF = 10, LCM = 1050

3  $2 \times 3^2 \times 5$



a  $2 \times 5 = 10$       b  $2 \times 2 \times 2 \times 3 \times 5 \times 7 = 840$

5 12 and 18

#### Ordering integers and decimals

1 a false      c true      e true  
 b true      d true

2  $-0.3, -1.5, -2.5, -4.2, -7.2$

3 0.049, 0.124, 0.412, 0.442, 1.002

4 a <      b <      c >

#### Calculating with negative numbers

**Stretch it!** Multiplying three negative numbers together always gives a negative answer.

1 a  $-8 - 3 = -11$       d  $14 + 4 = 18$   
 b 99      e 0  
 c  $-6$       f  $12 + 15 - 2 = 25$

2  $-8$  and  $9$

3  $32^\circ\text{C}$

#### Multiplication and division

**Stretch it!**      148419

$$\begin{array}{r} 621 \\ \times 239 \\ \hline 5589 \\ 18630 \\ 124200 \\ \hline 148419 \end{array}$$

1 a

$$\begin{array}{r} 235 \\ \times 9 \\ \hline 2115 \\ \hline \end{array}$$

2115

b

$$\begin{array}{r} 924 \\ \times 61 \\ \hline 924 \\ 55440 \\ \hline 56364 \end{array}$$

56364

2 a

$$\begin{array}{r} 047 \\ 5 \overline{)235} \\ \underline{5} \phantom{0} \\ 23 \\ \underline{23} \\ 0 \end{array}$$

47

b

$$\begin{array}{r} 0516 \\ 8 \overline{)4128} \\ \underline{32} \phantom{0} \\ 928 \\ \underline{736} \\ 192 \\ \underline{192} \\ 0 \end{array}$$

516

c

$$\begin{array}{r} 0126 \\ 17 \overline{)2146} \\ \underline{17} \phantom{0} \\ 44 \\ \underline{34} \\ 106 \\ \underline{102} \\ 4 \end{array}$$

126 remainder 4, or  $126 \frac{4}{17}$

3

$$\begin{array}{r} 033 \\ 8 \overline{)265} \\ \underline{16} \phantom{0} \\ 105 \\ \underline{96} \\ 9 \end{array}$$

remainder 1

a 33 boxes  
 b 1 pencil

4

$$\begin{array}{r} 091.25 \\ 4 \overline{)365.00} \\ \underline{32} \phantom{00} \\ 450 \\ \underline{40} \phantom{0} \\ 500 \\ \underline{48} \phantom{0} \\ 200 \\ \underline{200} \\ 0 \end{array}$$

£91.25

5

$$\begin{array}{r} 32 \\ \times 9 \\ \hline 288 \\ \hline \end{array}$$

£288

6

$$\begin{array}{r} 307.66 \\ 3 \overline{)923.00} \\ \underline{9} \phantom{00} \\ 2230 \\ \underline{21} \phantom{00} \\ 1300 \\ \underline{12} \phantom{00} \\ 1000 \\ \underline{9} \phantom{00} \\ 2000 \\ \underline{2000} \\ 0 \end{array}$$

$307.6 = 307\frac{2}{3}$

7

$$\begin{array}{r} 823 \\ \times 35 \\ \hline 4115 \\ 24690 \\ \hline 28805 \end{array}$$

28 805

$$\begin{array}{r} 037 \\ 12 \overline{)450} \\ \underline{36} \\ 90 \\ \underline{84} \\ 6 \end{array}$$

37 remainder 6, so  
37 boxes

- 9 He has not placed a zero in the ones column before multiplying through by 5. The  $\times 50$  line should have 5 digits: 36300, so his final three rows of working should look like this:

$$\begin{array}{r} 1452 \times 2 \\ 36300 \times 50 \\ \hline 37752 \end{array}$$

### Calculating with decimals

**Stretch it!**  $3.2 + 7.5 \times 2 = 3.2 + 15 = 18.2$

1 a 
$$\begin{array}{r} 3.40 \\ -1.07 \\ \hline 2.33 \end{array}$$
 2.33

c  $5 \times 7 = 35$   
 $0.05 \times 0.7 = 0.035$

b 
$$\begin{array}{r} 19.300 \\ + 5.091 \\ \hline 24.391 \end{array}$$

24.391

d 
$$\begin{array}{r} 321 \\ \times 19 \\ \hline 2889 \\ 3210 \\ \hline 6099 \end{array}$$

$3.21 \times 1.9 = 6.099$

e 
$$\begin{array}{r} 1.563 \\ 5 \overline{)7.815} \\ \underline{5} \phantom{.} \\ 28 \phantom{1} \\ \underline{25} \phantom{1} \\ 31 \\ \underline{30} \\ 15 \end{array}$$
 1.563

2 
$$\begin{array}{r} 64 \\ \times 24 \\ \hline 256 \\ 1280 \\ \hline 1536 \end{array}$$

$24 \times 64 = 1536$   
 $24 \times \text{£}0.64 = \text{£}15.36$   
 $\text{£}20 - \text{£}15.36 = \text{£}4.64$

3 
$$\begin{array}{r} 27.46 \\ 3 \overline{)82.38} \\ \underline{6} \phantom{.} \\ 22 \phantom{.} \\ \underline{21} \phantom{.} \\ 13 \\ \underline{12} \\ 18 \end{array}$$

Erica  $2 \times 27.46 = \text{£}54.92$   
Freya  $82.38 - 54.92 = \text{£}27.46$

### Rounding and estimation

**Stretch it!** a 1.0 b 1.00 c 1.000  
All the answers are 1

**Stretch it!**  $6.5 \times 8.5 = 55.25 \text{ m}^2$  – an overestimate.

- 1 a 0.35 c 32.6  
b 10 d 33100
- 2 a  $150 \leq x < 250$  c  $3.15 \leq x < 3.25$   
b  $5.5 \leq x < 6.5$  d  $5.055 \leq x < 5.065$
- 3  $\frac{30}{0.5 \times 6} = 10$

- 4 b is false since  $18 \times 1 = 18$  so  $18 \times 0.9$  cannot be 1.62  
c is false because if you divide by a number smaller than 1, the answer will be larger.

5 Night-time low tariff:  $2.32 \text{ units} \times 1.622\text{p} = 3.76\text{p}$   
 $7.151 \text{ units} \times 2.315\text{p} = 16.55\text{p}$   
20.31p

One tariff:  $2.320 + 7.151 = 9.471 \text{ units}$   
 $9.471 \text{ units} \times 1.923\text{p} = 18.21\text{p}$   
Tarik should choose One tariff, since it is cheaper.

### Converting between fractions, decimals and percentages

**Stretch it!**  $0.\dot{1}$ ,  $0.\dot{2}$ ,  $0.\dot{3}$ , ...  $0.\dot{4}$ ,  $0.\dot{5}$ . The number of ninths is the same as the digit that recurs. The exception is  $\frac{9}{9}$  which is the same as 1.

1 a  $\frac{32}{100} = \frac{8}{25}$  c  $\frac{33}{100}$   
b  $1\frac{24}{100} = 1\frac{6}{25}$  d  $\frac{95}{100} = \frac{19}{20}$

2 a 
$$\begin{array}{r} 0.41666\dots \\ 12 \overline{)5.00000} \\ \underline{48} \phantom{000} \\ 20 \phantom{00} \\ \underline{24} \phantom{00} \\ 60 \phantom{0} \\ \underline{60} \\ 0 \end{array}$$
 0.416

b 
$$\begin{array}{r} 0.375 \\ 8 \overline{)3.0000} \\ \underline{24} \phantom{000} \\ 60 \phantom{0} \\ \underline{60} \\ 0 \end{array}$$
 0.375

- c 0.49  
d 0.185

e 
$$\begin{array}{r} 0.42857142\dots \\ 7 \overline{)3.00000000} \\ \underline{28} \phantom{00000} \\ 20 \phantom{0000} \\ \underline{14} \phantom{0000} \\ 60 \phantom{000} \\ \underline{56} \phantom{000} \\ 40 \phantom{00} \\ \underline{35} \phantom{00} \\ 50 \phantom{0} \\ \underline{49} \\ 10 \end{array}$$
 0.428571

- 3 a  $\frac{91}{100} = 91\%$   
b  $\frac{3}{10} = \frac{30}{100} = 30\%$   
c  $\frac{4}{5} = \frac{80}{100} = 80\%$   
d  $\frac{9}{15} = \frac{3}{5} = \frac{6}{10} = \frac{60}{100} = 60\%$

4  $\frac{3}{8} \frac{0.375}{8 \overline{)3.0000}} = 0.375 = 37.5\%$

5  $0.35 \frac{2}{5} = \frac{4}{10} = 0.4$   $30\% = 0.3$   
 $30\%, 0.35, \frac{2}{5}$

6  $\frac{15}{20} = \frac{75}{100} = 75\%$  – Amy  
Rudi's mark was higher.

### Ordering fractions, decimals and percentages

1  $\frac{1}{3} = \frac{8}{24}$   $\frac{3}{8} = \frac{9}{24}$   $\frac{7}{12} = \frac{14}{24}$   
 $\frac{7}{12}, \frac{3}{8}, \frac{1}{3}$

2  $-2.2, 7, \frac{1}{5} = 0.2, -\frac{1}{10} = -0.1, 15\% = 0.15,$   
 $1\% = 0.01, 0.1$   
In order, this is:  $-2.2, -\frac{1}{10}, 1\%, 0.1, 15\%, \frac{1}{5}, 7.$   
The middle value is 0.1

- 3 Yes. If the numerator of a fraction is half the denominator then the fraction is equivalent to  $\frac{1}{2}$ .  
If the numerator is smaller than this the fraction must be less than  $\frac{1}{2}$ .

## Calculating with fractions

**Stretch it!** No – you could add the whole number parts, and then add the fraction parts.

- 1 **a**  $2\frac{3}{8} - \frac{3}{4} = \frac{19}{8} - \frac{3}{4} = \frac{19}{8} - \frac{6}{8} = \frac{13}{8} = 1\frac{5}{8}$   
**b**  $\frac{18}{17} \times \frac{2}{5} = \frac{6}{17}$   
**c**  $\frac{1}{7} \times 3\frac{1}{3} = \frac{1}{7} \times \frac{10}{3} = \frac{10}{21}$   
**d**  $2\frac{2}{5} + 5\frac{3}{4} = 7 + \frac{2}{5} + \frac{3}{4} = 7 + \frac{8}{20} + \frac{15}{20}$   
 $= 7\frac{23}{20}$   
 $= 8\frac{3}{20}$   
**e**  $\frac{1}{5} \div 2\frac{1}{2} = \frac{1}{5} \div \frac{5}{2} = \frac{1}{5} \times \frac{2}{5} = \frac{2}{25}$
- 2 **a**  $30 \div 5 = 6$   
 $6 \times 2 = 12$   
**b**  $40 \div 8 = 5$   
 $5 \times 7 = \text{£}35$   
**c**  $1818 \div 9 = 202$   
 $202 \times 4 = 808\text{mm}$
- 3  $35 \div 7 = 5$   
 $3 \times 7 = 15$   
 $35 - 15 = 20$
- 4 The number must be a multiple of 5, and  $\frac{2}{5}$  of it must be a multiple of 2.  
 $\frac{2}{5}$  of 45 = 18  
 $\frac{2}{5}$  of 40 = 16  
 $\frac{2}{5}$  of 35 = 14  
 $\frac{2}{5}$  of 30 = 12  
 $\frac{2}{5}$  of the number must be greater than 12, so the number is 35

## Percentages

- 1 **a**  $18 \div 100 = 0.18\text{cm}$   
 $0.18 \times 10 = 1.8\text{cm}$   
**b**  $1.20 \div 100 = \text{£}0.012$   
 $0.012 \times 25 = \text{£}0.30$   
**c**  $200\text{ml} \div 100 = 2\text{ml}$   
 $2\text{ml} \times 2 = 4\text{ml}$
- 2 **a**  $1.1 \times 30 = 33$   
**b**  $1.08 \times 500 = 540$   
**c**  $1.12 \times 91 = 101.92$ , so  $\text{£}101.92$
- 3 **a**  $0.8 \times 600 = 480$   
**b**  $0.95 \times 140 = 133$   
**c**  $0.81 \times 18 = 14.58$ , so  $\text{£}14.58$
- 4  $1.09 \times 2800 = 3052$
- 5  $0.65 \times 22\,000 = \text{£}14\,300$

## Order of operations

- 1 **a** 7  
**b**  $0.9 + 3.2 - \sqrt{36}$   
 $= 0.9 + 3.2 - 6$   
 $= -1.9$

$$\begin{aligned} \text{c } (-1)^2 - 14 \\ &= 1 - 14 \\ &= -13 \end{aligned}$$

- 2 30  
 3  $(8 - 3 + 5) \times 4$

## Exact solutions

- 1 **a**  $\pi$  **b**  $36\pi$  **c**  $2\frac{1}{2}\pi$  or  $\frac{5}{2}\pi$   
 2 **a**  $7\pi$  **b**  $\frac{5}{8}\pi$   
 3 Area  $= \frac{2}{7} \times \frac{3}{4} = \frac{6}{28} = \frac{3}{14}\text{cm}^2$   
 Perimeter  $= (2 \times \frac{3}{4}) + (2 \times \frac{2}{7}) = \frac{3}{2} + \frac{4}{7} = \frac{21+8}{14} = \frac{29}{14}$   
 $= 2\frac{1}{14}\text{cm}$   
 4 **a**  $2 \times 9 \times \pi = 18\pi\text{cm}$   
**b**  $12^2 \times \pi = 144\pi\text{cm}^2$   
 5 Circumference  $= 2 \times \pi \times 1 = 2\pi\text{cm}$   
 Length of one side of square  $= 2\pi \div 4 = \frac{1}{2}\pi\text{cm}$

## Indices and roots

- 1 **a**  $\frac{1}{3}$  **b**  $\frac{1}{0.4} = \frac{10}{4} = 2\frac{1}{2}$   
**c**  $\frac{1}{0.9} = \frac{10}{9} = 1\frac{1}{9}$
- 2  $3^2 = 9$ ,  $1^3 = 1$ ,  $\sqrt[3]{27} = 3$ ,  $\sqrt[3]{8} = 2$   
 In order, this gives  $1^3, \sqrt[3]{8}, \sqrt[3]{27}, 3^2$
- 3 **a** -8 **b** 1 **c** 81 **d** 1  
 4 **a**  $\frac{1}{4}$  **b\***  $\frac{1}{7^2} = \frac{1}{49}$   
**c\***  $\frac{1}{1^4} = 1$  **d**  $\frac{1}{3}$
- 5  $\frac{5^9}{5^5} = 5^4$

## Standard form

- 1 **a** 45 000 000 **b** 0.091  
 2 **a**  $6.45 \times 10^8$  **b**  $7.9 \times 10^{-8}$   
 3  $350000 - 4200 = 345800$   
 4  $3.2 \times 10^2 = 320$   $3.1 \times 10^{-2} = 0.031$   
 $3.09 \times 10 = 30.9$   $3 + (2.1 \times 10^2) = 213$   
 In order, this gives:  $3.1 \times 10^{-2}$   $3.09 \times 10$   
 $3 + (2.1 \times 10^2)$   $3.2 \times 10^2$
- 5  $3 \times 10^8\text{m/s}$   
 6  $200 \times 1.1 \times 10^{-4} = 2.2 \times 10^{-2} = 0.022\text{m} = 2.2\text{cm}$

## Listing strategies

### Stretch it!

red + small, red + medium, red + large,  
 green + small, green + medium, green + large,  
 blue + small, blue + medium, blue + large.

- 1 111,  
 112, 121, 211, 113, 131, 311,  
 222  
 221, 212, 122, 223, 232, 322  
 333  
 331 313 133 332 323 233  
 123 132 213 231 312 321

- 2 444 446 449  
464 466 469  
494 496 499
- 3 Small A, Small B, Small C, Small D  
Medium A, Medium B, Medium C, Medium D  
Large A, Large B, Large C, Large D.

### Review it!

- 1 7 and 6 (or 11 and 2, where both are prime and 2 is also a factor of 12)
- 2  $630 = 2 \times 3 \times 3 \times 5 \times 7 = 2 \times 3^2 \times 5 \times 7^*$
- 3  $18 = 2 \times 3 \times 3$   
 $36 = 2 \times 2 \times 3 \times 3$   
 $40 = 2 \times 2 \times 2 \times 5$   
HCF = 2
- 4 -11.5, -8.3, -3.5, -3.2, 1.4

5 a 
$$\begin{array}{r} 32.99 \\ +18.74 \\ \hline 51.73 \\ \small{1 \quad 1 \quad 1} \end{array}$$
 £51.73

b 
$$\begin{array}{r} 18.33 \\ 3 \overline{)54.99} \end{array}$$
 £18.33

6 a 
$$\begin{array}{r} 23 \\ \times 14 \\ \hline 92 \times 4 \\ 230 \times 10 \\ \hline 322 \\ \small{1} \end{array}$$
  $23 \times 0.14 = 3.22$

b 
$$\begin{array}{r} 149 \\ \times 27 \\ \hline 1043 \times 7 \\ 2980 \times 20 \\ \hline 4023 \\ \small{1 \quad 1} \end{array}$$
 4023

7  $81 \div 3 = 27$

8 
$$\begin{array}{r} 031 \\ 11 \overline{)3345} \text{ remainder } 4 \end{array}$$
  $31\frac{4}{11}$

9 a 
$$\begin{array}{r} 0.375 \\ 8 \overline{)3.0000} \end{array}$$
 0.375

b  $0.7 \times 100 = 70\%$

10 a  $70\% = \frac{70}{100} = \frac{7}{10}$

b  $0.8 = \frac{8}{10} = \frac{4}{5}$

11  $\frac{1}{2} = \frac{2}{4}$   $\frac{1}{2}$  is larger

$\frac{2}{7} = \frac{8}{28}$   $\frac{1}{4} = \frac{7}{28}$   $\frac{2}{7}$  is larger

$\frac{3}{11} = \frac{12}{44}$   $\frac{1}{4} = \frac{11}{44}$   $\frac{3}{11}$  is larger

$\frac{2}{5} = \frac{8}{20}$   $\frac{1}{4} = \frac{5}{20}$   $\frac{2}{5}$  is larger

They all are

12 a  $\frac{3}{5} + \frac{1}{7} = \frac{21}{35} + \frac{5}{35} = \frac{26}{35}$

b  $2\frac{1}{5} - \frac{7}{10} = \frac{11}{5} - \frac{7}{10} = \frac{22}{10} - \frac{7}{10} = \frac{15}{10} = 1\frac{1}{2}$

c  $\frac{2}{3} \div \frac{4}{9} = \frac{2}{3} \times \frac{9}{4} = \frac{3}{2} = 1\frac{1}{2}$

13  $0.25 - 0.07 = 0.18 = \frac{18}{100} = \frac{108}{600}$

$\frac{2}{3} - \frac{1}{2} = \frac{4}{6} - \frac{3}{6} = \frac{1}{6} = \frac{100}{600}$

0.25 - 0.07 is larger

14  $\frac{3}{5} \times \frac{5}{4} = \frac{3}{4}$

15  $\frac{45}{1000} = \frac{9}{200}$

16  $8.6 \div 100 = 0.086$

$0.086 \times 25 = \text{£}2.15$

17 a 9

b 5

18 a  $3.4 \times 10^9$

b  $3.04 \times 10^{-7}$

19  $37.55 \leq x < 37.65$

20 a 51

b 12, 15, 21, 51, 25, 52

21 a  $200 \times 9 \times 10 = 18000 = \text{£}180.00$

b Underestimate since all numbers were rounded down.

22  $40\% \text{ of } 600 = 240$

$\frac{1}{5} \text{ of } 600 = 120$

$600 - (240 + 120) = 240$

23 More than 33%, less than 50%, multiple of 5. 35%

24 No, since 2 is a prime number and odd + odd + even = even

25  $0.8 \times 349 = \text{£}279.20$

26 a 3.1

b 3.05

27 a 325 000

b 320 000

28  $3 \times 3 \times 3 \times 3 \times 3 \times 3 = 729$

29 a  $26.25 + 18.23 + (4 \times 5.5) = \text{£}66.48$

$\text{£}66.48 \div 4 = \text{£}16.62$

30  $0.19 \times 18\ 000 = 3420$

31 a 2010 and 2011

b  $1.1 \times 102.3 = 112.53$

## Algebra

### Understanding expressions, equations, formulae and identities

1 a  $3a + 6 = 10$  (It can be solved to find the value of  $a$ .)

b  $C = \pi D$  (The value of  $C$  can be worked out if the value of  $D$  is known.)

c  $3(a + 2)$  (It does not have an equals sign.)

d  $3ab + 2ab = 5ab$  (Collecting the like terms on the left-hand side gives  $5ab$  which is equal to the right-hand side.)

2 James is correct.

$4x - 2 = 2x$  can be solved to find the value of  $x$  so it is an equation.

Or, the two sides of  $4x - 2 = 2x$  are not equal for all values of  $x$  so it cannot be an identity. For example, when  $x = 2$ :

(Left-hand side)  $4x - 2 = 4 \times 2 - 2 = 6$

(Right-hand side)  $2x = 2 \times 2 = 4$

$6 \neq 4$

## Simplifying expressions

## Stretch it!

The expressions must all contain algebra, so each part must include  $t$ .

There are four possible combinations that make  $12t^3$ :

$$12t \times t \times t, 2t \times 6t \times t, 2t \times 3t \times 2t, 3t \times 4t \times t.$$

- 1 a  $p^3$   
 b  $4 \times b \times c \times 7 = 4 \times 7 \times b \times c = 28bc$   
 c  $4a \times 3b = 4 \times 3 \times a \times b = 12ab$   
 d  $5x \times 4x = 5 \times 4 \times x \times x = 20x^2$   
 e  $2g \times (-4g) = 2 \times (-4) \times g \times g = -8g^2$   
 f  $2p \times 3q \times r = 2 \times 3 \times p \times q \times r = 6pqr$
- 2 a  $10x \div 2 = \frac{10x}{2} = 5x$   
 b  $\frac{14w}{-2} = -7w$   
 c  $6p \div p = \frac{6p}{p} = 6$   
 d  $8mn \div 2m = \frac{8mn}{2m} = 4n$   
 e  $\frac{12xy}{3y} = 4x$   
 f  $9abc \div bc = \frac{9abc}{bc} = 9a$

## Collecting like terms

- 1 a  $5f$   
 b  $7b$   
 c  $5mn$   
 d  $3d + 4e + d - 6e = 3d + d + 4e - 6e = 4d - 2e$   
 e  $2x + 5y + 3x - 2y - 2 = 2x + 3x + 5y - 2y - 2 = 5x + 3y - 2$   
 f  $2a - b - 5a - 3 = 2a - 5a - b - 3 = -3a - b - 3$   
 g  $2x^2$   
 h  $2t^3 + 4 - t^3 - 4 = 2t^3 - t^3 + 4 - 4 = t^3$   
 i  $7\sqrt{x}$   
 j  $3\sqrt{x}$   
 k  $7\sqrt{x}$

## Using indices

- 1 a  $x^5 \times x^4 = x^{5+4} = x^9$   
 b  $p \times p^4 = p^{1+4} = p^5$   
 c  $2m^4 \times 3m^4 = 2 \times 3 \times m^4 \times m^4 = 6 \times m^{4+4} = 6m^8$   
 d  $3m^4n \times 5m^2n^3$   
 $= 3 \times 5 \times m^4 \times m^2 \times n \times n^3$   
 $= 15 \times m^{4+2} \times n^{1+3} = 15m^6n^4$   
 e  $u^{-2} \times u^5 = u^{-2+5} = u^3$   
 f  $t^7 \times t^{-6} = t^{7+(-6)} = t$
- 2 a  $x^4 \div x^2 = x^{4-2} = x^2$   
 b  $\frac{y^7}{y^3} = y^{7-3} = y^4$   
 c  $\frac{p^9}{p^8} = p^{9-8} = p$   
 d  $8x^6 \div 4x^3 = \frac{8x^6}{4x^3}$   
 $(8 \div 4) \times (x^6 \div x^3) = 2 \times x^{6-3} = 2x^3$   
 e  $m^3 \div m^5 = m^{3-5} = m^{-2} = \frac{1}{m^2}$   
 f  $\frac{5x^8}{15x^4} = \frac{5}{15} \times \frac{x^8}{x^4} = \frac{1}{3} \times x^{8-4} = \frac{x^4}{3}$   
 g  $3x^2 \div 9x = \frac{3x^2}{9x} = \frac{3}{9} \times \frac{x^2}{x} = \frac{1}{3} \times x^{2-1} = \frac{x}{3}$

- 3 a  $(x^2)^3 = x^{2 \times 3} = x^6$   
 b  $(y^4)^4 = y^{4 \times 4} = y^{16}$   
 c  $(p^5)^2 = p^{5 \times 2} = p^{10}$   
 d  $(4m^5)^2 = 4^2 \times (m^5)^2 \times 16 \times m^{5 \times 2} = 16m^{10}$   
 e  $(x^2)^{-3} = x^{2 \times (-3)} = x^{-6} = \frac{1}{x^6}$   
 f  $(n^{-4})^{-2} = n^{-4 \times (-2)} = n^8$
- 4 a  $2x \times 3x^2 = 2 \times 3 \times x \times x^2 = 6 \times x^{1+2} = 6x^3$   
 b  $\frac{x^8}{x} = x^{8-1} = x^7$   
 c  $\frac{2x^3}{8x^2} = \frac{2}{8} \times \frac{x^3}{x^2} = \frac{1}{4} \times x^{3-2} = \frac{x}{4}$   
 d  $(3y^3)^3 = 3^3 \times (y^3)^3 = 27 \times y^{3 \times 3} = 27y^9$   
 e  $y^{-4} = \frac{1}{y^4}$   
 f  $a^2b \times a^3b^2$   
 $= a^2 \times a^3 \times b \times b^2 = a^{2+3} \times b^{1+2} = a^5b^3$

## Expanding brackets

## Stretch it!

- 1  $(x+2)(x+4) = x^2 + 6x + 8$
- 2 a  $2x^2 + 8x + 6$       b  $3x^2 + 10x - 8$   
 c  $6x^2 + 7x - 3$
- 1 a  $3a + 6$       b  $4b - 16$       c  $10c + 25$   
 d  $6 - 2e$       e  $4x + 4y + 8$       f  $-2y - 4$   
 g  $x^2 - 2x$       h  $2a^2 + 10a$
- 2 a  $2(2x + 3) + 4(x + 5) = 4x + 6 + 4x + 20 = 8x + 26$   
 b  $3(3y + 1) + 2(4y - 3) = 9y + 3 + 8y - 6 = 17y - 3$   
 c  $4(2m + 4) - 3(2m - 5) = 8m + 16 - 6m + 15 = 2m + 31$
- 3 a  $(x + 2)(x + 3) = x^2 + 3x + 2x + 6 = x^2 + 5x + 6$   
 b  $(y - 3)(y + 4) = y^2 + 4y - 3y - 12 = y^2 + y - 12$   
 c  $(a + 3)(a - 7) = a^2 - 7a + 3a - 21 = a^2 - 4a - 21$   
 d  $(m - 1)(m - 6) = m^2 - 6m - m + 6 = m^2 - 7m + 6$
- 4 a  $(x + 1)^2 = (x + 1)(x + 1)$   
 $= x^2 + x + x + 1 = x^2 + 2x + 1$   
 b  $(x - 1)^2 = (x - 1)(x - 1)$   
 $= x^2 - x - x + 1 = x^2 - 2x + 1$   
 c  $(m - 2)^2 = (m - 2)(m - 2)$   
 $= m^2 - 2m - 2m + 4 = m^2 - 4m + 4$   
 d  $(y + 3)^2 = (y + 3)(y + 3)$   
 $= y^2 + 3y + 3y + 9 = y^2 + 6y + 9$

## Factorising

## Stretch it!

The width of the rectangle =  $x + 1$ , since  $x^2 + 3x + 2 = (x + 2)(x + 1)$

- 1 a  $3(a + 3)$       b  $5(b - 2)$   
 c  $7(1 + 2c)$       d  $d(d - 2)$
- 2 a  $4(2a + 5)$       b  $4(b - 3)$   
 c  $9(2 + c)$       d  $d(2d - 3)$
- 3 a  $2(2x - 3y)$       b  $m(a + b)$   
 c  $x(4x + 3y)$       d  $n(2 - 9n)$   
 e  $5x(1 + 2y)$       f  $4p(q - 3)$
- 4 a  $(x + 1)(x + 7)$       b  $(x - 1)(x + 5)$   
 c  $(x + 2)(x - 4)$       d  $(x - 2)(x - 3)$   
 e  $(x - 3)(x - 3)$       f  $(x + 3)(x + 4)$   
 g  $(x - 2)(x + 5)$       h  $(x + 4)(x - 5)$

- 5 a  $x^2 - 16 = x^2 - 4^2 = (x + 4)(x - 4)$   
 b  $x^2 - 36 = x^2 - 6^2 = (x + 6)(x - 6)$   
 c  $x^2 - 81 = x^2 - 9^2 = (x + 9)(x - 9)$   
 d  $y^2 - 100 = y^2 - 10^2 = (y + 10)(y - 10)$

### Substituting into expressions

- 1  $c = 5 \times 3 + 2 \times (-2) = 15 + (-4) = 11$   
 2 a  $2 - 2 \times (-4) = 2 - (-8) = 10$   
 b  $3 \times 2 \times (-4) = -24$   
 c  $4 \times (-4) - 3 \times 2 = -16 - 6 = -22$   
 d  $2^2 + (-4)^2 = 4 + 16 = 20$   
 e  $2 \times 2 + 4(2 - (-4)) = 2 \times 2 + 4 \times 6 = 4 + 24 = 28$   
 f  $\frac{1}{2}(2 + (-4)) = \frac{1}{2} \times -2 = -1$
- 3 False.  
 When  $a = 3$ :  $3a^2 = 3 \times 3^2 = 3 \times 9 = 27$
- 4 a  $10 \times \frac{1}{2} \times 4 = 20$   
 b  $8 \times \left(\frac{1}{2}\right)^2 = 8 \times \left(\frac{1}{4}\right) = 2$   
 c  $0.5 \times 4 \times 10 \times 10 \times 10 \times 10 \times 10 = 200\ 000$   
 ( $= 2 \times 10^5$ )  
 d  $(4 \times 3) \div (10 \times 10 \times 10) = 12 \div 1000 = 0.012$   
 e  $2 \times 4 \times (10 \times 10 \times 10 \times 10 \times 10) \times 3 \div (10 \times 10 \times 10)$   
 $2 \times 4 \times 100\ 000 \times 3 \div 1000$   
 $= 2\ 400\ 000 \div 1000$   
 $= 2400$   
 f  $4^2 - 12 \times \frac{1}{2} = 16 - 12 \times \frac{1}{2} = 16 - 6 = 10$

### Writing expressions

- 1 a  $4 - q$                                   b  $n + m$  (or  $m + n$ )  
 c  $8t$     d  $xy$  (or  $yx$ )  
 e  $p^2$     f  $a^3$
- 2  $x + y$   
 3  $100n + 75b$   
 4 Perimeter  $= 3a + 2a + 4 + 4a - 2 = 9a + 2$   
 5 Area  $= \frac{1}{2} \times 4 \times (2a + 5) = 2 \times (2a + 5) = 4a + 10$

### Solving linear equations

- 1 a  $5a = 35$   
 $a = \frac{35}{5}$   
 $a = 7$   
 b  $b - 9 = 8$   
 $b = 8 + 9$   
 $b = 17$   
 c  $\frac{c}{4} = 4$   
 $c = 4 \times 4$   
 $c = 16$   
 d  $d + 4 = 2$   
 $d = 2 - 4$   
 $d = -2$
- 2 a  $2x + 3 = 13$   
 $2x = 10$   
 $x = 5$

- b  $3y - 4 = 11$   
 $3y = 15$   
 $y = 5$   
 c  $2p + 9 = 1$   
 $2p = -8$   
 $p = -4$   
 d  $\frac{f}{3} - 7 = 4$   
 $\frac{f}{3} = 11$   
 $f = 33$   
 e  $\frac{x+5}{2} = 8$   
 $x + 5 = 16$   
 $x = 11$   
 f  $\frac{f-7}{3} = 4$   
 $f - 7 = 12$   
 $f = 19$
- 3 a  $9 - m = 7$   
 $9 = 7 + m$   
 $2 = m$   
 b  $10 - 3x = 1$   
 $10 = 1 + 3x$   
 $9 = 3x$   
 $3 = x$   
 c  $7 - 2x = 2$   
 $7 = 2 + 2x$   
 $5 = 2x$   
 $\frac{5}{2} = x$   
 (Or  $x = 2.5$ , or  $x = 2\frac{1}{2}$ )  
 d  $5 = 1 - 2f$   
 $5 + 2f = 1$   
 $2f = -4$   
 $f = -2$
- 4 Hannah has not subtracted 4 from *both* sides.  
 Correct working:  
 $2x + 4 = 8$   
 $2x = 4$   
 $x = 2$
- 5 a  $3(a + 2) = 15$   
 $3a + 6 = 15$   
 $3a = 9$   
 $a = 3$   
 b  $4(b - 2) = 4$   
 $4b - 8 = 4$   
 $4b = 12$   
 $b = 3$   
 c  $3(4c - 9) = 9$   
 $12c - 27 = 9$   
 $12c = 36$   
 $c = 3$   
 d  $2(d + 3) + 4 = 2$   
 $2d + 6 + 4 = 2$   
 $2d + 10 = 2$   
 $2d = -8$   
 $d = -4$

- e**  $4(2x + 3) - 2 = 6$   
 $8x + 12 - 2 = 6$   
 $8x + 10 = 6$   
 $8x = -4$   
 $x = -\frac{4}{8} = -\frac{1}{2}$   
(Or  $x = -0.5$ )
- 6 a**  $3m = m + 6$   
 $2m = 6$   
 $m = 3$
- b**  $5t - 6 = 2t + 3$   
 $3t - 6 = 3$   
 $3t = 9$   
 $t = 3$
- c**  $4x + 3 = 2x + 8$   
 $2x + 3 = 8$   
 $2x = 5$   
 $x = \frac{5}{2}$   
(Or  $x = 2.5$  or  $x = 2\frac{1}{2}$ )
- d**  $3 - 2p = 6 - 3p$   
 $3 + p = 6$   
 $p = 3$
- e**  $3y - 8 = 5y + 4$   
 $-8 = 2y + 4$   
 $-12 = 2y$   
 $-6 = y$
- 7 a**  $2(x + 5) = x + 6$   
 $2x + 10 = x + 6$   
 $x + 10 = 6$   
 $x = -4$
- b**  $7b - 2 = 2(b + 4)$   
 $7b - 2 = 2b + 8$   
 $5b - 2 = 8$   
 $5b = 10$   
 $b = 2$
- c**  $4(2y + 1) = 3(5y - 1)$   
 $8y + 4 = 15y - 3$   
 $4 = 7y - 3$   
 $7 = 7y$   
 $1 = y$
- d**  $2x - 1 = 8 - 4x$   
 $6x - 1 = 8$   
 $6x = 9$   
 $x = \frac{9}{6} = \frac{3}{2}$   
(Or  $x = 1.5$  or  $x = 1\frac{1}{2}$ )
- 2 a** Angles in a quadrilateral add up to  $360^\circ$  so:  
 $x + 20 + 2x - 15 + x + 65 + 2x - 10 = 360$   
 $6x + 60 = 360$   
 $6x = 300$   
 $x = 50$
- b** Largest angle:  $x + 65 = 50 + 65 = 115^\circ$   
(Other angles:  $x + 20 = 50 + 20 = 70^\circ$ ;  
 $2x - 15 = 2 \times 50 - 15 = 85^\circ$ ;  
 $2x - 10 = 2 \times 50 - 10 = 90^\circ$ )
- 3** Let  $a =$  Karen's age  
Monica is 4 years younger:  $a - 4$   
 $a + a - 4 = 64$   
 $2a - 4 = 64$   
 $2a = 68$   
 $a = 34$   
Karen is 34 years old.  
 $a - 4 = 34 - 4 = 30$   
Monica is 30 years old.
- 4** Let  $n =$  number.  
 $2n + 4 = 16 - n$   
 $3n + 4 = 16$   
 $3n = 12$   
 $n = 4$   
The number is 4.
- 5** Let  $l =$  length of rectangle.  
Width is 2 cm smaller:  $l - 2$   
Perimeter =  $2l + 2(l - 2)$   
 $= 2l + 2l - 4 = 4l - 4$   
 $4l - 4 = 36$   
 $4l = 40$   
 $l = 10$   
Length is 10 cm.  
 $l - 2 = 10 - 2 = 8$   
Width is 8 cm.
- 6** Base angles of an isosceles triangle are equal so:  
 $4a - 20 = 2a + 16$   
 $2a - 20 = 16$   
 $2a = 36$   
 $a = 18$   
When  $a = 18$ :  $4a - 20 = 4 \times 18 - 20 = 52$   
So  $2a + 16 = 52$   
Angles in a triangle add up to  $180^\circ$  so:  
 $4b - 2a + 52 + 52 = 180$   
 $4b - 2a + 104 = 180$   
 $4b - 2a = 76$  (Substitute  $a = 18$ )  
 $4b - 2 \times 18 = 76$   
 $4b - 36 = 76$   
 $4b = 112$   
 $b = 28$

### Writing linear equations



- 1 a** Perimeter =  $4 \times (2s + 3) = 8s + 12$   
(Or, Perimeter =  $2s + 3 + 2s + 3 + 2s + 3 + 2s + 3 = 8s + 12$ )
- b**  $8s + 12 = 84$   
 $8s = 72$   
 $s = 9\text{cm}$

### Linear inequalities

- 1 a**  $x = 3, 4, 5$   
**b**  $x = 2, 3, 4, 5$   
**c**  $x = 0, 1, 2, 3$   
**d**  $x = -3, -2, -1, 0, 1$

2 a  $x < 3$       b  $x \geq -2$


c  $-1 \leq x \leq 5$



4 a  $2x - 2 > 4$

$2x > 6$

$x > 3$



b  $4x + 3 \leq 13$

$4x \leq 10$

$x \leq \frac{10}{4} = \frac{5}{2}$


(Or  $x \leq 2.5$  or  $x \leq 2\frac{1}{2}$ )



c  $4x < 2x - 10$

$2x < -10$

$x < -5$



d  $7x + 2 \geq 3x - 2$

$4x + 2 \geq -2$

$4x \geq -4$

$x \geq -1$



5 Olivia has not multiplied all the terms in the bracket by the term outside.

Correct working:

$3(x + 4) > 22$

$3x + 12 > 22$

$3x > 10$

$x > \frac{10}{3}$

(Or  $x > 3\frac{1}{3}$ )

6 a  $2x + 3 < 4x + 8$

$3 < 2x + 8$

$-5 < 2x$

$-\frac{5}{2} < x$

(Or  $x > -\frac{5}{2}$ )

b Smallest integer value of  $x$  that satisfies the inequality is  $-2$

7 No, Lee is not correct.

$6 \leq 2x + 4 < 16$

$-2 \leq 2x < 12$

$-1 \leq x < 6$

Inequality is not true when  $x = 6$  so  $x = 6$  is not a possible solution.

Alternative method:

When  $x = 6$ :

$6 \leq 2 \times 6 + 4 < 16$

$6 \leq 16 < 16$

Since 16 is not less than 16,  $x = 6$  is not a possible solution.

8 a  $4 - x \leq 1$

$4 \leq x + 1$

$3 \leq x$  (Or  $x \geq 3$ )

Alternative method:

$4 - x \leq 1$

$-x \leq -3$

$x \geq 3$

b  $6 - 3x > 9$

$6 > 3x + 9$

$-3 > 3x$

$-1 > x$  (Or  $x < -1$ )

Alternative method:

$6 - 3x > 9$

$-3x > 3$

$x < -1$

c  $8 - 2x \geq 7$

$8 \geq 2x + 7$

$1 \geq 2x$

$\frac{1}{2} \geq x$  (Or  $x \leq \frac{1}{2}$ )

Alternative method:

$8 - 2x \geq 7$

$-2x \geq -1$

$x \leq \frac{1}{2}$

d  $-2 < -x \leq 3$

$2 > x \geq -3$

### Formulae

1 Pay =  $8 \times 35 + 25 = 280 + 25 = 305$

Pay = £305

2  $P = 2(8 + 5.5) = 2 \times 13.5 = 27$

3  $v = 10 + (-20) \times 5 = 10 + (-100) = -90$

4  $C = 25d + 50$

5  $A = P^2$

6 a  $P = 2a + 2(a + 3) = 2a + 2a + 6 = 4a + 6$

(Or  $P = a + a + a + 3 + a + 3 = 4a + 6$ )

b  $P = 4 \times 6 + 6 = 24 + 6 = 30$

$P = 30 \text{ cm}$

7  $-10 = \frac{D}{6.5}$

$-65 = D$

8 a  $v = u + at$

$v - u = at$

$\frac{v - u}{t} = a$

b  $V = \frac{1}{3}Ah$

$3V = Ah$

$\frac{3V}{A} = h$




c  $y = 3(x - 3)$   
 $y = 3x - 9$   
 $y + 9 = 3x$   
 $\frac{y + 9}{3} = x$   
 Or:  $x = \frac{y}{3} + 3$

d  $v^2 = u^2 + 2as$   
 $v^2 - u^2 = 2as$   
 $\frac{v^2 - u^2}{2a} = s$

e  $T = \sqrt{\frac{2s}{g}}$   
 $T^2 = \frac{2s}{g}$   
 $gT^2 = 2s$   
 $g = \frac{2s}{T^2}$

### Linear sequences

- 1 a i 2, 5, 8, 11, **14, 17**  
 ii 23, 19, 15, 11, **7, 3**  
 iii 3, 9, 15, 21, **27, 33**  
 iv 4, 9, 14, 19, **24, 29**
- b i 2, 5, 8, 11, 14, 17, 20, 23, 26, **29**  
 (Or, 10th term =  $2 + (3 \times 9) = 29$ )  
 ii 23, 19, 15, 11, 7, 3, -1, -5, -9, **-13**  
 (Or, 10th term =  $23 - (4 \times 9) = -13$ )  
 iii 3, 9, 15, 21, 27, 33, 39, 45, 51, **57**  
 (Or, 10th term =  $3 + (6 \times 9) = 57$ )  
 iv 4, 9, 14, 19, 24, 29, 34, 39, 44, **49**  
 (Or, 10th term =  $4 + (5 \times 9) = 49$ )
- 2 a 1st term =  $1 \times 4 - 2 = 2$   
 2nd term =  $2 \times 4 - 2 = 6$   
 3rd term =  $3 \times 4 - 2 = 10$   
 4th term =  $4 \times 4 - 2 = 14$
- b 20th term =  $20 \times 4 - 2 = 78$
- 3 -25, -18, -11, -4, **3, 10**
- 4 a 
- b Number of triangles: 2, 4, 6, 8, 10, 12, 14, 16  
 So 16 triangles in pattern number 8  
 Or,  $2 + 7 \times 2 = 16$  triangles
- c No. The number of triangles forms an even number sequence and 35 is odd.
- 5 a 3, 7, 11, 15, 19  
 Common difference = +4  
 $4 \times$  term number = 4, 8, 12, 16, 20  
 - 1 to get each term in the original sequence  
 So,  $n$ th term is  $4n - 1$
- b  $4n - 1 = 99$   
 $4n = 100$   
 $n = 25$   
 Yes, 99 is a term in the sequence because 25 is an integer.

### Non-linear sequences

- 1 1, 3, 5, 7, 9, ... Arithmetic sequence (Term-to-term rule is add 2)  
 1, 2, 4, 8, 16, ... Geometric sequence (Term-to-term rule is multiply by 2, or double)  
 1, 4, 5, 9, 14, ... Fibonacci-type sequence (Next term of sequence is found by adding the previous two terms together)  
 1, 4, 9, 16, 25, ... Square-number sequence (Sequence of square numbers  $1^2, 2^2, 3^2, 4^2, \dots$ )
- 2 a  $4, 2, 1, \frac{1}{2}, \frac{1}{4}$  ( $\div 2$ )  
 b 5, 0.5, 0.05, **0.005, 0.0005** ( $\div 10$ )  
 c  $\frac{1}{2}, \frac{1}{4}, \frac{1}{8}, \frac{1}{16}, \frac{1}{32}$  ( $\div 2$ )  
 d  $\frac{1}{9}, \frac{1}{3}, 1, \mathbf{3, 9}$  ( $\times 3$ )  
 e -0.1, -0.2, -0.4, **-0.8, -1.6** ( $\times 2$ )  
 f 3, -6, 12, **-24, 48** ( $\times -2$ )
- 3 When  $n = 5$ :  
 $3 \times 5^2 - 4 = 3 \times 25 - 4 = 71$
- 4 a 1st term =  $a$   
 2nd term =  $b$   
 3rd term =  $a + b$   
 4th term =  $b + a + b = a + 2b$   
 5th term =  $a + b + a + 2b = 2a + 3b$
- b  $b = 5$   
 $2a + 3b = 23$  (Substitute  $b = 5$ )  
 $2a + 3 \times 5 = 23$   
 $2a + 15 = 23$   
 $2a = 8$   
 $a = 4$

### Show that...


- 1 a LHS =  $4(x - 3) + 2(x + 5) = 4x - 12 + 2x + 10 = 6x - 2$   
 RHS =  $3(2x - 1) + 1 = 6x - 3 + 1 = 6x - 2$   
 LHS = RHS  
 So  $4(x - 3) + 2(x + 5) \equiv 3(2x - 1) + 1$
- b LHS =  $(x + 2)(x - 2) = x^2 - 2x + 2x - 4 = x^2 - 4$   
 LHS = RHS  
 So  $(x + 2)(x - 2) = x^2 - 4$
- 2 Rod A =  $n$   
 Rod B =  $n + 1$   
 Rod C =  $n + 2$   
 Rod A + Rod C =  $n + n + 2 = 2n + 2 = 2(n + 1)$   
 Rod A + Rod C is 2 times the length of rod B.

### Functions

- 1 a  $10 \times 3 - 3 = 30 - 3 = 27$   
 b  $(9 + 3) \div 3 = 12 \div 3 = 4$   
 c If  $x = y$ ,  
 $3x - 3 = x$   
 $2x - 3 = 0$   
 $2x = 3$   
 $x = \frac{3}{2}$   
 (Or  $x = 1 \frac{1}{2}$ )

## Coordinates and midpoints

- 1 a A(2, 2)  
b and c




- d B(-4, 2), D(2, -3)  
x-coordinate:  $2 + (-4) = -2$   
 $-2 \div 2 = -1$   
y-coordinate:  $-3 + 2 = -1$   
 $-1 \div 2 = -0.5$   
Midpoint is (-1, -0.5)

## Straight-line graphs

### Stretch it!


To solve the equation, you need to find where the graph of  $y = 3x - 2$  intersects the graph of  $y = -x$ .



So the solution to  $3x - 2 = -x$  is  $x = 0.5$ .



1

x	-2	-1	0	1	2
y	4	3	2	1	0



2

x	-2	0	2
y	-7	-3	1



$$\text{Gradient} = \frac{\text{difference in } y \text{ coordinate}}{\text{difference in } x \text{ coordinate}} = \frac{5}{5} = 1$$

$$\text{Gradient, } m = 1$$

$$y\text{-intercept, } c = 1$$

Using the general form of the equation of a line  $y = mx + c$ : the equation of the line is  $y = x + 1$

- 4 Gradient,  $m = 2$

$$y = mx + c \text{ (General form of the equation of a line)}$$

$$y = 2x + c$$

$$\text{For point } (1, -2), x = 1, y = -2:$$

$$-2 = 2 \times 1 + c$$

$$-2 = 2 + c$$

$$-4 = c$$

$$\text{So the equation of the line is } y = 2x - 4$$

- 5 Gradient =  $\frac{\text{difference in } y \text{ coordinate}}{\text{difference in } x \text{ coordinate}} = \frac{4 - 2}{0 - 4} = \frac{-2}{-4} = \frac{1}{2}$


$$\text{Gradient, } m = \frac{1}{2}$$

$$\text{Given the point } (0, 4): y\text{-intercept, } c = 4$$

$$\text{So the equation of the line is } y = \frac{1}{2}x + 4$$


6 a

x	0	6
y	6	0



b


x	0	4
y	2	0



- 7 A:  $y = 4x + 1$   
 B:  $4x + 4y = 4$   
 $4y = -4x + 4$   
 $y = -x + 1$   
 C:  $x - 2y = 2$   
 $-2y = -x + 2$   
 $y = \frac{x}{2} - 1$   
 D:  $2y = 4 + 8x$   
 $y = 2 + 4x$   
 $y = 4x + 2$

Lines A and D are parallel.

- 8  $2y = x - 4$   
 $y = \frac{x}{2} - 2$   
 The y-intercept is  $(0, -2)$



- a  $y = 3.4$       b  $x = 0.5$   
 c  $x \approx 2.5$       d  $x \approx -0.5$

### Solving simultaneous equations

- 1 a  $2x + y = 4$  (1)  
 $3x - y = 1$  (2)  
 (1) + (2):  $5x = 5$   
 $x = 1$   
 Substitute  $x = 1$  in (1)  
 $2 \times 1 + y = 4$   
 $2 + y = 4$   
 $y = 2$   
 Solution:  $x = 1, y = 2$
- b  $x - y = 5$  (1)  
 $2x + y = 4$  (2)  
 (1) + (2):  $3x = 9$   
 $x = 3$   
 Substitute  $x = 3$  into (1)  
 $3 - y = 5$   
 $3 = y + 5$   
 $y = -2$   
 Solution:  $x = 3, y = -2$
- c  $2x + y = 8$  (1)  
 $x + y = 2$  (2)  
 (1) - (2):  $x = 6$   
 Substitute  $x = 6$  into (2)  
 $6 + y = 2$   
 $y = -4$   
 Solution:  $x = 6, y = -4$
- d  $4x - y = 10$  (1)  
 $x + 2y = 7$  (2)  
 (1)  $\times$  2:  $8x - 2y = 20$  (3)  
 (2) + (3):  $9x = 27$   
 $x = 3$   
 Substitute  $x = 3$  into (2):  
 $3 + 2y = 7$   
 $2y = 4$   
 $y = 2$   
 Solution:  $x = 3, y = 2$
- e  $2x + y = 7$  (1)  
 $x - 4y = 8$  (2)  
 (1)  $\times$  4:  $8x + 4y = 28$  (3)  
 (2) + (3):  $9x = 36$   
 $x = 4$   
 Substitute  $x = 4$  into (1):  
 $2 \times 4 + y = 7$   
 $8 + y = 7$   
 $y = -1$   
 Solution:  $x = 4, y = -1$
- f  $2x + 3y = 7$  (1)  
 $3x - 2y = 4$  (2)  
 (1)  $\times$  2:  $4x + 6y = 14$  (3)  
 (2)  $\times$  3:  $9x - 6y = 12$  (4)  
 (3) + (4):  $13x = 26$   
 $x = 2$

Substitute  $x = 2$  into (1)

$$2 \times 2 + 3y = 7$$

$$4 + 3y = 7$$

$$3y = 3$$

$$y = 1$$

Solution:  $x = 2, y = 1$

**2**  $x + y = 21$  (1)

$x - y = 7$  (2)

(1) + (2):  $2x = 28$

$x = 14$

Substitute  $x = 14$  into (1)

$$14 + y = 21$$

$$y = 7$$

The two numbers are 7 and 14.

**3** Let  $b =$  burger and  $c =$  cola.

$3b + 2c = 505$  (1)

$3b + 4c = 725$  (2)

(2) - (1):  $2c = 220$

$c = 110$

Substitute  $c = 110$  into (1)

$$3b + 2 \times 110 = 505$$

$$3b + 220 = 505$$


$$3b = 285$$

$$b = 95$$

A burger costs 95p.


A cola costs £1.10.

**4 a**



$x = 2, y = 8$

**b**




$x = 8, y = 1$

## Quadratic graphs

### Stretch it!

Rearrange  $x^2 - 3x = 3$ , to give  $x^2 - 3x - 2 = 1$


You can solve this graphically by finding where the lines  $y = x^2 - 3x - 2$  and  $y = 1$  intersect.



So the solutions to the equation  $x^2 - 3x = 3$  are  $x = 3.8$  and  $x = -0.8$ . Acceptable readings from the graph would be in the range 3.6 to 3.9 and  $-0.6$  to  $-0.9$ .

**1 a**

$x$	-2	-1	0	1	2	3	4
$y$	6	1	-2	-3	-2	1	6




**b i**  $x = 1$

**ii**  $(1, -3)$

**2**

$x$	-2	-1	0	1	2	3	4
$y$	8	3	0	-1	0	3	8



- a**  $x = 0$  and  $x = 2$   
**b**  $x \approx -1.2$  and  $x \approx 3.2$   
**c**  $x = 1$

### Solving quadratic equations

#### Stretch it!

$$\frac{x^2}{2} = 8$$

$$x^2 = 16$$

$$x = \sqrt{16}$$

$$\text{So } x = 4 \text{ or } x = -4$$

$$2x^2 = 50$$

$$x^2 = 25$$

$$x = \sqrt{25}$$

$$\text{So } x = 5 \text{ or } x = -5$$

**1 a**  $x^2 - 4x = 0$

$$x(x - 4) = 0$$

$$\text{Either } x = 0 \text{ or } x - 4 = 0$$

$$x = 4$$

$$\text{So } x = 0 \text{ or } x = 4$$

**b**  $x^2 + 7x = 0$

$$x(x + 7) = 0$$

$$\text{Either } x = 0 \text{ or } x + 7 = 0$$

$$x = -7$$

$$\text{So } x = 0 \text{ or } x = -7$$

**c**  $x^2 - 16 = 0$  ( $x^2 - 16 = x^2 - 4^2$ , Factorise)

$$(x + 4)(x - 4) = 0$$

$$\text{Either } x + 4 = 0 \text{ or } x - 4 = 0$$

$$x = -4 \quad x = 4$$

$$\text{So } x = -4 \text{ or } x = 4$$

**d**  $x^2 + 10x + 9 = 0$

$$(x + 1)(x + 9) = 0$$

$$\text{Either } x + 1 = 0 \text{ or } x + 9 = 0$$

$$x = -1 \quad x = -9$$

$$\text{So } x = -1 \text{ or } x = -9$$

**e**  $x^2 + x - 12 = 0$

$$(x - 3)(x + 4) = 0$$

$$\text{Either } x - 3 = 0 \text{ or } x + 4 = 0$$

$$x = 3 \quad x = -4$$

$$\text{So } x = 3 \text{ or } x = -4$$

**f**  $x^2 - 6x - 16 = 0$

$$(x + 2)(x - 8) = 0$$

$$\text{Either } x + 2 = 0 \text{ or } x - 8 = 0$$

$$x = -2 \quad x = 8$$

$$\text{So } x = -2 \text{ or } x = 8$$

**2 a**  $y = x^2 - 49$  (Set  $y = 0$ )

$$x^2 - 49 = 0$$
 ( $x^2 - 49 = x^2 - 7^2$ , Factorise)

$$(x + 7)(x - 7) = 0$$

$$\text{Either } x + 7 = 0 \text{ or } x - 7 = 0$$

$$x = -7 \quad x = 7$$

$$\text{So } x = -7 \text{ or } x = 7$$

**b**  $y = x^2 - 3x$  (Set  $y = 0$ )

$$x^2 - 3x = 0$$

$$x(x - 3) = 0$$

$$\text{Either } x = 0 \text{ or } x - 3 = 0$$

$$x = 3$$

$$\text{So } x = 0 \text{ or } x = 3$$

**c**  $y = x^2 + 7x + 6$  (Set  $y = 0$ )

$$x^2 + 7x + 6 = 0$$

$$(x + 1)(x + 6) = 0$$

$$\text{Either } x + 1 = 0 \text{ or } x + 6 = 0$$

$$x = -1 \quad x = -6$$


$$\text{So } x = -1 \text{ or } x = -6$$

### Cubic and reciprocal graphs

**1 a**

$x$	-3	-2	-1	0	1	2	3
$y$	-28	-9	-2	-1	0	7	26


**b**



**2 a**


$x$	-4	-3	-2	-1	$-\frac{1}{2}$	$-\frac{1}{4}$	$\frac{1}{4}$	$\frac{1}{2}$	1	2	3	4
$y$	$\frac{1}{4}$	$\frac{1}{3}$	$\frac{1}{2}$	1	2	4	-4	-2	-1	$-\frac{1}{2}$	$-\frac{1}{3}$	$-\frac{1}{4}$

**b**




### Drawing and interpreting real-life graphs

**1 a**




- b** i €36  
 ii £75  
**c** From the graph: £30 = €36  
 So £90 = €36 × 3 = €108  
 Ring is cheaper in France.

- 2 a** Monthly charge = £10 (cost of 0 minutes from the graph)  
**b** Gradient =  $\frac{30}{240} = 0.125$   
 Charge per minute of calls is 13p.  
**3** A: The temperature is steadily increasing.  
 B: The temperature remains constant.  
 C: The temperature rises steadily for a period and then remains constant.  
**4 a** 30 minutes (Horizontal line on graph)  
**b** 55 km  
**c** Speed before break =  $\frac{\text{distance (km)}}{\text{time (hours)}} = \frac{30}{1} = 30 \text{ km/hr}$   
 Speed after break =  $\frac{\text{distance (km)}}{\text{time (hours)}} = \frac{25}{1.5} = 16.7 \text{ km/hr}$



- 5 a** 6 m/s  
**b** 4 seconds  
**c** 6 seconds  
**d** Acceleration =  $\frac{\text{change in speed (m/s)}}{\text{time (s)}} = \frac{6}{4} = 1.5 \text{ m/s}^2$




**b** Acceleration =  $\frac{\text{change in speed (m/s)}}{\text{time (s)}} = \frac{20-10}{30} = \frac{10}{30} = 0.3 \text{ m/s}^2$

- 7 a** The maximum depth of water in the bath before the person got in was 35cm  
**b** Between C and D, the person was taking their bath.  
**c** Between D and E, the person got out of the bath.

- d** Running water into the bath was quicker. The slope of the line between O and A (filling the bath) is steeper than the slope of the line between E and F (emptying the bath).

### Review it!

- 1 a** P(3, 5)  
**b, c and e**




- d** Q(3, -2), S(-4, 5)  
 x-coordinate:  $-4 + 3 = -1$   
 $-1 \div 2 = -0.5$   
 y-coordinate:  $5 + (-2) = 3$   
 $3 \div 2 = 1.5$   
 Midpoint is (-0.5, 1.5)

- 2 a**  $2x + 8 = 4$   
 $2x = -4$   
 $x = -2$   
**b**  $4 \times 4 - 3 \times 3 = 16 - 9 = 7$   
**c** Millie is correct.  
 When  $x = 4$ ,  $3x^2 = 3 \times 4^2 = 3 \times 16 = 48$   
 (George has worked out  $(3x)^2$  instead.)

- 3 a**  $4(2x + 3)$   
**b**  $m^4 \times m = m^{4+1} = m^5$   
**c**  $\frac{x^8}{x^3} = x^{8-3} = x^5$

**4 a**

x	-1	0	1	2	3
y	-1	1	3	5	7



- c** Compare  $y = 2x + 1$  with  $y = mx + c$  (general form of the equation of a line):  
 Gradient,  $m = 2$

5  $4x + 4 = x + 13$

$3x + 4 = 13$

$3x = 9$

$x = 3$

6 a  $x \leq 2$



c  $x = 1, 2, 3$

d  $4x + 2 \leq 2x + 5$

$2x + 2 \leq 5$

$2x \leq 3$

$x \leq \frac{3}{2}$

(Or  $x \leq 1 \frac{1}{2}$ )

Largest integer value of  $x$  that satisfies the inequality = 1

7 a 3, 9, 15, 21, 27, 33

b No. This is a sequence of odd numbers and 44 is even.

c When  $n = 5$ :

$2n^2 - 3 = 2 \times 5^2 - 3 = 2 \times 25 - 3 = 47$

8  $(x + 3)(x + 4) = x^2 + 4x + 3x + 12 = x^2 + 7x + 12$

9 The opposite sides of a rectangle are equal in length so:

$5x - 8 = 2x + 4$

$3x - 8 = 4$

$3x = 12$

$x = 4$

$14 - 2y = 4y + 2$

$14 = 6y + 2$

$12 = 6y$


$2 = y$

10 a  $12m$

b  $3p \times 4p = 3 \times 4 \times p \times p = 12p^2$

c  $12x \div 2 = \frac{12x}{2} = 6x$

11 a



b i 25 cm

ii From the graph: 25 cm = 10 inches

So 50 cm =  $10 \times 2 = 20$  inches

c From the graph: 10 inches = 25 cm

So 60 inches =  $25 \times 6 = 150$  cm

Cost of beading =  $150 \times 2 = 300$ p

Cost = £3.00

12 a  $T = 12.50x + 10$

b  $72.50 = 12.50x + 10$

$62.50 = 12.50x$

$5 = x$

Suzanne hired the costume for 5 days.

13 a  $4x + 2 \leq 8$

$4x \leq 6$

$x \leq \frac{6}{4} = \frac{3}{2}$

(Or  $x \leq 1 \frac{1}{2}$ )

b  $3x - 4 < 17$

$3x < 21$

$x < 7$

$4x + 2 \geq 22$

$4x \geq 20$

$x \geq 5$

If  $x < 7$  and  $x \geq 5$  then  $x = 6$  and  $x = 5$  satisfy both.

14 Ollie has squared each term inside the brackets rather than squaring the whole bracket.

Correct working:

$(x + 4)^2 = (x + 4)(x + 4) = x^2 + 4x + 4x + 16 = x^2 + 8x + 16$

15  $P = \frac{Q}{4} + R$

$P - R = \frac{Q}{4}$

$4(P - R) = Q$

16 a  $m(m + 8)$

b  $(x + 3)(x + 4)$

17 a 2, 5, 8, 11, 14

Common difference = +3

$3 \times$  term number: 3, 6, 9, 12, 15

- 1 to get each term in the original sequence

So  $n$ th term =  $3n - 1$

b  $2n - 3 = 112$

$2n = 115$

$n = 57.5$

No, Kadena is incorrect.

112 cannot be a term in the sequence because 57.5 is not an integer.

18 a  $4(x + 5) - 3(2x - 1) = 4x + 20 - 6x + 3 = -2x + 23$

b  $4a^3b^2 \times 5a^2b = 4 \times 5 \times a^3 \times a^2 \times b^2 \times b$   
 $= 20 \times a^{3+2} \times b^{2+1} = 20a^5b^3$

19 Perimeter =  $3x - 2 + 2x + 1 + 3x + 5 + 2x = 10x + 4$

$10x + 4 = 49$

$10x = 45$

$x = 4.5$

20 a 1st term:  $a$

2nd term:  $b$

3rd term:  $a + b$

4th term:  $b + a + b = a + 2b$

5th term:  $a + b + a + 2b = 2a + 3b$

6th term:  $a + 2b + 2a + 3b = 3a + 5b$

7th term:  $2a + 3b + 3a + 5b = 5a + 8b$

b  $a + b = 5$  (1)

$5a + 8b = 34$  (2)

(1)  $\times 5$ :  $5a + 5b = 25$  (3)

$$(2) - (3): 3b = 9$$

$$b = 3$$

Substitute  $b = 3$  into (1)

$$a + 3 = 5$$

$$a = 2$$

## Ratio, proportion and rates of change

### Units of measure

- 3000 m
  - 75 mins
  - 13 000 cm<sup>2</sup>
  - 3.52 litres
  - 7200 seconds
  - 14 kg
- $4.5 - 0.325 = 4.175$  kg or  $4500 - 325 = 4175$  g
- $5 \div 2.2 = 2.2\dot{7}$  kg

### Ratio

**Stretch it!**  $31 + 25 = 56$ , fraction male =  $\frac{31}{56}$

- 1:4      b 1:3:4      c 4:5
- $35:5 = 7:1$
- 7:1
  - $800 \div (7 + 1) = 100$   
 $1 \times 100 = 100$  tickets
- 3:2
  - $\frac{3}{5}$  of 200 =  $(200 \div 5) \times 3 = 120$
- $1500 \text{ g} \div (2 + 3) = 300$   
 $2 \times 300 = 600$  g or 0.6 kg
- $9 \div 3 = 3$   
 $4 \times 3 = 12$  cm  
 $5 \times 3 = 15$  cm
- $s = 20t$
- $5 - 2 = 3$   
 $60 \div 3 = 20$  g  
 $2 \times 20 \text{ g} = 40$  g

### Scale diagrams and maps

**Stretch it!** 50 miles on ground =  $\frac{50}{x}$  miles on map

$$1 \text{ mile} = 1610 \text{ m} = 161\,000 \text{ cm}$$

$$50 \text{ miles on ground} = \frac{50}{x} \times 161\,000 \text{ cm}^* \text{ on map}$$

- A, B, F
- $3 \times 12 = 36$  km
  - $15 \div 12 = 1.25$  cm
- $12 \times 1000 = 12\,000$  cm = 120 m
- 2 cm:  $2 \times 50\,000 = 100\,000$  cm = 1 km  
(Any answer within the range of 1 km – 1.1 km is acceptable.)
  - 250°

### Fractions, percentages and proportion

- $\frac{20}{3500} = \frac{1}{175}$
- $2 + 3 + 8 = 13$  hours  
 $24 - 13 = 11$  hours  
 $\frac{11}{24}$  of the day remaining

- $\frac{15}{20} = \frac{3}{4}$
  - $1 - \frac{3}{4} = \frac{1}{4} = 25\%$
- $1 + 2 + 7 = 10$ ,  $\frac{1}{10} = 10\%$
- School A:  $125:145 = 25:29$   
School B:  $100:120 = 5:6$   
No since the ratios are not equivalent.
- $150 \div 100 = 1.5$   
 $1.5 \times 22 \text{ g} = 33 \text{ g}$   
 $33 \text{ g} \div 8 = 4.125 \text{ g}$

### Direct proportion

#### Stretch it!

For two values to be in direct proportion, when one is 0 the other must be 0. Here, when distance is 0 miles, the fee is £2.

- A and E
- 20 meringues = 2 eggs, divide both by 2 to give: 10 meringues = 1 egg  
3 eggs:  $3 \times 10 = 30$  meringues
    - 20 meringues = 120 g of sugar, divide both by 2 to give: 10 meringues = 60 g of sugar. Multiply both by 10 to give 100 meringues
  - 20 meringues = 2 eggs, divide both by 2 to give 10 meringues = 1 egg, multiply both by 7 to give 70 meringues = 7 eggs
- $675 \div 4.5 = 150$  minutes = 2 hours 30 minutes
- A, D

### Inverse proportion

- D
- At 60 miles it takes 15 minutes.  
 $60 \times \frac{2}{3} = 40$   
 $15 \div \frac{2}{3} = 22.5$  mins
- $2 \times 3 = 6$  decorators  
 $5 \div 3 = 1\frac{2}{3}$  of a day
- 2
  - The age of the chicken and the number of eggs it lays are in inverse proportion, this means that as the age of the chicken increases, the number of eggs it lays decreases.

### Working with percentages

**Stretch it!** £128

**Stretch it!** Let percentage rate =  $x$

$$(1 + \frac{x}{100})^5 \times \text{£}100 = \text{£}110$$

$$(1 + \frac{x}{100})^5 = \frac{110}{100}$$

$$1 + \frac{x}{100} = \sqrt[5]{\frac{110}{100}}$$

$$1 + \frac{x}{100} = 1.02$$

$$\frac{x}{100} = 0.02$$

$$x = 2$$

Percentage interest is 2%

\*This answer differs from the one in the Revision Guide due to an error in our first edition. This answer has now been re-checked and corrected.



- 1 a  $1.03 \times 50 = \text{£}51.50$   
 b  $2.48 \times 400 = 992$   
 c  $0.195 \times 64 = 12.48$
- 2  $45 - 40 = 5, \frac{5}{40} \times 100 = 12.5\%$
- 3  $24 \div 115 = 0.209, 0.209 \times 100 = 20.9^\circ\text{C}$
- 4  $15\,000 \times 1.20^3 = 25920$
- 5 20% is  $\frac{1}{5}$  of the price.  
 $30 \times 5 = \text{£}150$

### Compound units

Stretch it!  $\frac{100}{x}$  mph

- 1  $29.50 \div 0.18 = 164$  or  $2950 \div 18 = 164$  units
- 2 Time =  $\frac{80}{120} = \frac{2}{3}$  hour = 40 minutes
- 3 Density =  $\frac{0.72}{3} = 0.24 \text{ g/cm}^3$
- 4 Pressure =  $\frac{12}{2} = 6 \text{ N/m}^2$
- 5  $3 \text{ m/s} = 3 \times 60 \text{ m/minute} = 3 \times 60 \times 60 \text{ m/hour}$   
 $= 10800 \text{ m/hour} = 10.8 \text{ km/hour}$
- 6 0.6 litres per second =  $0.6 \times 60$  litres per minute  
 $= 0.6 \times 60 \times 60$  litres per hour  
 $= 2160$  litres per hour.
- $2160 \div 4.55 = 475$  gallons  
 475 gallons per hour (to the nearest whole number)
- 7 Bolt: 100 m in 9.58 seconds = 10.4 m/s  
 Cheetah: 120 km/h = 120000 m/hour  
 $= 120000 \div 60 \text{ m/min}$   
 $= 2000 \text{ m/min}$   
 $= 2000 \div 60 \text{ m/sec} = 33.3 \text{ m/s}$
- The Cheetah is fastest.

### Review it!



- 1 a  $3.2 \times 1000 = 3200 \text{ m}$   
 b  $9 \times 60 = 540$  seconds  
 c  $0.4 \times 1000 = 400 \text{ ml}$
- 2  $4600 \div 1000 = 4.6 \text{ km}$
- 3  $2.5 \times 60 = 150$  minutes
- 4  $1.1 \times 0.32 = 0.352 \text{ m}^2$  or  $110 \times 32 = 3520 \text{ cm}^2$
- 5  $3 \times 10000 = 30000 \text{ cm}^2$
- 6  $\frac{5}{12}$
- 7  $26:18 = 13:9$
- 8  $100 - 85 = 15, 15 \div 3 = 5$  minutes
- 9 density =  $\frac{345}{0.15} = 2300 \text{ kg/m}^3$
- 10  $10 - 8 = 2 \text{ km}, \frac{2}{8} \times 100 = 25\%$
- 11  $25 - 13 = 12, \frac{12}{25}$  OR 48%
- 12  $15 + 5 + 3 = 23$  mins  
 $\frac{23}{90}$
- 13  $20 \div (\frac{4}{5}) = 25$  hours = 1 day and 1 hour
- 14 a  $50 \div 5 = 10$ , Josie:  $1 \times 10 = 10$  marbles,  
 Charlie:  $4 \times 10 = 40$  marbles, Charlie has 30 more.  
 b  $C = 4J$
- 15  $\frac{100}{360} \times 100 = 28\%$
- 16  $0.8 \times 1200 = \text{£}960$   
 $0.9 \times 960 = \text{£}864$



- 17 1 cm: 50 000 cm  
 $50\,000 \text{ cm} = 0.5 \text{ km}$   
 $3 \text{ km} \div 0.5 = 6$   
 6 cm
- 18  $1.02^3 \times 1500 = \text{£}1591.81$
- 19  $32\,000 \div 4 = 8000$  people
- 20  $393 \div 125 = 3.144$  hours = 3 hours 9 minutes
- 21  $2.50 + 1.90 + (2 \times 5.30) = \text{£}15$   
 $1.05 \times \text{£}15 = \text{£}15.75$
- 22  $37 + 15 + 4 + 19 = 75$   
 $\frac{15}{75} \times 100 = 20\%$
- 23  $0.045 \times 3000 = \text{£}135$   
 $3000 + (5 \times 135) = \text{£}3675$
- 24  $30 \div 3 = 10$   
 boys =  $2 \times 10 = 20$   
 Girls =  $1 \times 10 = 10$   
 Boys =  $20 - 2 = 18$   
 Girls =  $10 + 3 = 13$   
 18:13
- 25 Men to women is 7:6 = 35:30  
 Ratio of women to children is 15:2 = 30:4  
 Ratio of men to women to children is 35:30:4  
 $35 + 30 + 4 = 69$   
 $3450 \div 69 = 50$   
 $35 \times 50 = 1750$  men
- 26 No – for two things to be in direct proportion when one is zero the other must be zero; the graph does not go through the origin so this is not the case.
- 27 Neither, since the time taken to cook increases as the weight increases it is not in indirect proportion. It is not in direct proportion since a graph to illustrate the relationship would not go through the origin.
- 28 speed =  $\frac{\text{distance}}{\text{time}} = \frac{0.05}{17} = \frac{1}{340}$  hours =  $\frac{3}{17}$  mins  
 $= 11$  seconds
- 29 She is incorrect since the ratio of females to males must be the same for them to have equivalent proportions: 35:60 is not equivalent to 12:37.

## Geometry and measures

### Measuring and drawing angles

- 1 a  $43^\circ$       b Acute





**b**  $18^\circ$

### Using the properties of angles

- Angles around a point add up to  $360^\circ$  so:
 
$$a + 112 + 88 + 106 = 360$$

$$a + 306 = 360$$

$$a = 54^\circ$$
- a**
  - $a = (180 - 40) \div 2 = 70^\circ$
  - Base angles of an isosceles triangle are equal.
- Exterior angle of a triangle is equal to the sum of the interior angles at the other two vertices so:
 
$$b = 70 + 40$$

$$b = 110^\circ$$

**Or**, angles on a straight line add up to  $180^\circ$  so:

$$b = 180 - 70 = 110^\circ$$
- Angles around a point add up to  $360^\circ$  so:
 
$$5x + 9x + 108 = 360$$

$$14x + 108 = 360$$

$$14x = 252$$

$$x = 18^\circ$$
- a**
  - $x = 180 - 126 = 54^\circ$
  - Angles on a straight line add up to  $180^\circ$ .
- Angles in a quadrilateral add up to  $360^\circ$  so:
 
$$y + 135 + 54 + 88 = 360$$

$$y + 277 = 360$$

$$y = 83^\circ$$
- a** Angles on a straight line add up to  $180^\circ$  so:
 
$$x = 180 - 84$$

$$x = 96^\circ$$
- i**  $y = 96^\circ$
- Use the fact that corresponding angles are equal, then the fact that vertically opposite angles are equal.
 

**Or**, use the fact that alternate angles are equal, then use angles on a straight line add up to  $180^\circ$ .
- a** Base angles of an isosceles triangle are equal so  $a = 58^\circ$ .
- Angles in a triangle add up to  $180^\circ$  so:
 
$$b = 180 - 58 - 58$$

$$b = 64^\circ$$
- Alternate angles are equal so  $c = 58^\circ$  (since angle  $a =$  angle  $c$ ).

Or, since opposite angles of a parallelogram are equal:

$$b + c = 122$$

$$64 + c = 122$$

$$c = 58^\circ$$

- Angle  $BAD = 62^\circ$  (Opposite angles of a parallelogram are equal)  
 Angle  $ADE = 62^\circ$  (Alternate angles are equal)  
 $x = 180 - 62 - 62$  (Base angles of an isosceles triangle are equal)  
 $x = 56^\circ$
- Angle  $ACB = 36^\circ$  (Base angles of an isosceles triangle are equal)  
 Angle  $ABC = 180 - 36 - 36$  (Angles in a triangle add up to  $180^\circ$ )  
 Angle  $ABC = 108^\circ$   
 $x = 108^\circ$  (Alternate angles are equal)

### Using the properties of polygons

#### Stretch it!

- The angle sum of a triangle is  $180^\circ$ .  
 Sum of interior angles of a hexagon =  $4 \times 180^\circ = 720^\circ$ .

Polygon	Number of sides ( $n$ )	Number of triangles formed	Sum of interior angles
Triangle	3	1	$180^\circ$
Quadrilateral	4	2	$360^\circ$
Pentagon	5	3	$540^\circ$
Hexagon	6	4	$720^\circ$
Heptagon	7	5	$900^\circ$
Octagon	8	6	$1080^\circ$
Decagon	10	8	$1440^\circ$

- $n - 2$
  - $180 \times (n - 2)$
- Stretch it!** Exterior angle of a regular hexagon =  $360 \div 6 = 60^\circ$

Interior angle =  $180 - 60 = 120^\circ$

Three hexagons meet at a point, so  $120 + 120 + 120 = 360^\circ$

Similarly, interior angle of an octagon =  $180 - (360 \div 8) = 135^\circ$

Interior angle of a square =  $90^\circ$ , so  $135 + 135 + 90 = 360^\circ$ .


Regular pentagons have an interior angle of  $108^\circ$ . This does not divide equally into  $360^\circ$ , so these shapes will not fit together at a point in this way.

- Regular decagon has 10 equal sides.  
 Exterior angle =  $360^\circ \div 10 = 36^\circ$
- a** Number of sides =  $360^\circ \div 15^\circ = 24$
- b** Angles on a straight line add up to  $180^\circ$  so:  
 Interior angle + exterior angle =  $180$   
 Interior angle +  $15 = 180$   
 Interior angle =  $165^\circ$   
 Sum of interior angles =  $24 \times 165 = 3960^\circ$

- 3 Exterior angle =  $360^\circ \div 8 = 45^\circ$   
 Angles on a straight line add up to  $180^\circ$  so:  
 $x = 180^\circ - 45^\circ = 135^\circ$

**Using bearings**



- 1 a  $360 - 45$  (acute angle) =  $315^\circ$




- 2 Bearing of P from Q =  $180^\circ + 164^\circ = 344^\circ$   
 3 Kirsty is correct.  
 The bearing is  $314^\circ$  ( $360^\circ - 46^\circ$ ) as it must be measured clockwise from north.

**Properties of 2D shapes**

**Stretch it!**



- b kite  
 2 a 8 possible lines of symmetry:



- b 8

- 3 a A rectangle has rotational symmetry of order 2.  
 b A **rhombus** has all sides equal and rotational symmetry of order 2.  
 c A kite has 1 line of symmetry and **no** rotational symmetry.  
 d The diagonals of a **square** and a **rhombus** bisect each other at  $90^\circ$ .


**Congruent shapes**

- 1 Any accurate copy of shape A, in any orientation.  
 2 a Corresponding angles are equal so  $x = 120^\circ$   
 b Corresponding sides are the same length so  $y = 12$  cm  
 3 a SSS (each triangle has equal sides: 3 cm, 3 cm, 2.5 cm)  
 b ASA (two angles,  $70^\circ$  and  $60^\circ$ , and the included side, 8 cm, are equal)



**Constructions**




**Stretch it!**

A triangle with sides of 5cm with constructions lines indicating the use of compasses





Angle size  $60^\circ$







### Drawing circles and parts of circles


- 1 a A **chord** is a straight line that does not pass through the centre of a circle but touches the circumference at each end.
- b A **tangent** is a straight line that touches the outside of a circle at one point only.
- c A **diameter** is a straight line through the centre of a circle that touches the circumference at each end.
- d An **arc** is part of the circumference of a circle.
- e A **radius** is a straight line from the centre of a circle that is half the length of the diameter.
- f The part of a circle that has a chord and an arc as its boundary is called a **segment**.



### Loci



3 a and b



### Perimeter

- 1  $4 \times 7.2 = 28.8 \text{ cm}$
- 2  $7 + 9 + 9 + 5 + 5 + 7 = 42 \text{ cm}$
- 3 Curved edge  $= 2\pi r \div 2 = (2 \times \pi \times 4) \div 2 = 4\pi$   
Perimeter  $= 4\pi + 2 \times r = 4\pi + 8 \text{ cm}$   
So  $k = 4$  and  $b = 8$
- 4 Perimeter  $= (\pi \times 30) + 100 + 100 = 200 + 30\pi \text{ m}$
- 5 Perimeter  $= \left(\frac{1}{2} \times \pi \times 32\right) + 32 + 32 = 16\pi + 64 \text{ cm}$   
Ribbon  $= 16\pi + 64 + 5 = 16\pi + 69 \text{ cm} = 119.3 \text{ cm}$   
120 cm must be bought  
 $12 \times \text{£}0.15 = \text{£}1.80$

### Area

**Stretch it!** Area of a semicircle =  $\frac{\pi r^2}{2}$ ,  
 area of a quarter circle =  $\frac{\pi r^2}{4}$

- 1 a  $4.5 \times 2 = 9.0 \text{ cm}^2$   
 b  $3 \times 1.5 = 4.5 \text{ cm}^2$   
 c  $\frac{(5+9)}{2} \times 4 = 28.0 \text{ cm}^2$   
 d  $\frac{1}{2} \times 2 \times 5 = 5.0 \text{ cm}^2$   
 e  $\pi \times 4.5^2 = 63.6 \text{ cm}^2$
- 2 Length of side =  $12 \div 4 = 3 \text{ cm}$   
 Area =  $3^2 = 9 \text{ cm}^2$
- 3 Shaded triangles would fit together to form one triangle with base  $10 - 6 = 4$ .  
 So area of shaded triangles =  $\frac{1}{2} \times 4 \times 7 = 14 \text{ cm}^2$   
 Area of trapezium =  $\frac{(6+10)}{2} \times 7 = 56 \text{ cm}^2$   
 Fraction of the shape that is shaded =  $\frac{14}{56} = \frac{1}{4}$
- 4 Area of whole shape =  $6 \times 8 = 48 \text{ cm}^2$   
 Fraction shaded =  $\frac{6}{16} = \frac{3}{8}$   
 Area shaded =  $(\frac{3}{8}) \times 48 = 18 \text{ cm}^2$
- 5 Area of square =  $46 \times 46 = 2116 \text{ cm}^2$   
 Each circle has radius =  $11.5 \text{ cm}$   
 Area of four circles =  $4 \times \pi \times 11.5^2 = 1661.9 \text{ cm}^2$   
 Shaded area =  $2116 - 1661.9 = 454.1 \text{ cm}^2$

### Sectors


- 1 Area =  $\frac{1}{2} \times \pi \times 5^2 = 39.3 \text{ cm}^2$   
 Perimeter =  $\frac{1}{2} \times \pi \times 10 + 10 = 25.7 \text{ cm}$
- 2 Area =  $\frac{3}{4} \times \pi \times 4^2 = 12\pi \text{ cm}^2$
- 3 Area =  $\frac{1}{2} \times \pi \times 3^2 = 14.1 \text{ m}^2$   
 $14.1 \div 2 = 7.05$ , so 8 bags needed.  
 $8 \times 14.99 = \text{£}119.92$

### 3D shapes

**Stretch it!**


3D shape	Faces	Edges	Vertices
Cube	6	12	8
Cuboid	6	12	8
Square-based pyramid	5	8	5
Tetrahedron	4	6	4
Triangular prism	5	9	6
Hexagonal prism	8	18	12

**Stretch it!** There are three planes of symmetry for the first cuboid:




\*This answer differs from the one in the Revision Guide due to an error in our first edition. This answer has now been re-checked and corrected.


There are 5 planes of symmetry for the second cuboid: the same 3 planes as the first cuboid, plus two more planes along the diagonals of the square faces.



**Stretch it!\***








1 a 6 possible rectangular faces:



c Kelli has not counted the hidden edges.

- 2\* Draw a square in the middle with sides of 4 units (1 unit represents 1 cm). Set your compasses to 6 units and draw pairs of intersecting arcs from the corners of the square. These are the apices (top points) of the triangular sides. Draw lines for the sides of the triangles.



### Volume

- $\frac{4}{3} \times \pi \times 4.5^3 = 381.7 = 382 \text{ cm}^3$  (to 3 s.f.)
- $\pi r^2 h + \frac{1}{3} \pi r^2 h = \pi \times 0.5^2 \times 2 + \frac{1}{3} \times \pi \times 0.5^2 \times 1.5$   
 $= 0.625 \pi = 1.96 \text{ m}^3$
- $\frac{1}{3} \times \pi \times 6^2 \times 22 = \frac{1}{3} \times 792 \times \pi = 264\pi \text{ cm}^3$   
 $k = 264$
- Volume of water =  $18 \times 7 \times 7 = 882 \text{ cm}^3$   
 $882 = 7 \times 20 \times h$   
 $882 = 140 \times h$   
 $h = 6.3 \text{ cm}$

### Surface area

- $6 \times (5 \times 5) = 150 \text{ cm}^2$
- $4\pi r^2 = 4 \times \pi \times 3^2 = 36\pi \text{ cm}^2$
- $18 - 4 = 14 \text{ cm}^2$

- sloping surface =  $\pi \times 14 \times 45 = 630\pi \text{ cm}^2$   
Base =  $\pi \times 14^2 = 196\pi \text{ cm}^2$   
Total surface area =  $196\pi + 630\pi = 826\pi$   
Percentage yellow =  $\frac{630}{826} \times 100 = 76.3\%$

### Using Pythagoras' theorem

- Using Pythagoras' theorem  $c^2 = a^2 + b^2$ :

$$AC^2 = AB^2 + BC^2$$

$$15^2 = 11^2 + BC^2$$

$$BC^2 = 15^2 - 11^2 = 104$$

$$BC = \sqrt{104}$$

$$BC = 10.2 \text{ cm (to 3 s.f.)}$$

- $c^2 = a^2 + b^2$

$$6^2 = 3.6^2 + b^2$$

$$b^2 = 6^2 - 3.6^2 = 23.04$$

$$b = \sqrt{23.04}$$

$$b = 4.8$$

The ladder reaches 4.8 m up the wall.

- Using Pythagoras' theorem  $c^2 = a^2 + b^2$ :

$$XZ^2 = XY^2 + YZ^2$$

$$15^2 = XY^2 + 9^2$$

$$XY^2 = 15^2 - 9^2 = 144$$

$$XY = \sqrt{144}$$

$$XY = 12 \text{ cm}$$

$$\text{Area} = \frac{1}{2} bh = \frac{1}{2} \times 9 \times 12$$

$$\text{Area} = 54 \text{ cm}^2$$

- If the triangle is right-angled,  $PQ^2 = PR^2 + RQ^2$

$$PQ^2 = 13^2 = 169$$


$$PR^2 + RQ^2 = 8^2 + 5^2 = 64 + 25 = 89^*$$

$$PQ^2 \neq PR^2 + RQ^2$$

Claudia is not correct.

Notice that  $PR + RQ = 8 + 5 = 13 \text{ cm} = \text{length of } PQ$ , so PQR isn't a triangle at all, it is just a straight line!

- $P(-2, 5), Q(5, -3)$




$$PQ^2 = 8^2 + 7^2 = 113$$

$$PQ = \sqrt{113}$$

$$PQ = 10.63 \text{ (to 2 d.p.)}$$

\*This answer differs from the one in the Revision Guide due to an error in our first edition. This answer has now been re-checked and corrected.


6 A(2, 4), B(5, 6)



$$AB^2 = 2^2 + 3^2 = 13$$

$$AB = \sqrt{13}$$

7



Using Pythagoras' theorem  $c^2 = a^2 + b^2$ :

$$AB^2 = AX^2 + BX^2$$

$$AB^2 = 7^2 + 20^2 = 449$$

$$AB = \sqrt{449}$$

$$AB = 21.2 \text{ (to 3 s.f.)}$$

$$\text{Perimeter of field ABCD} = 15 + 20 + 8 + 21.2 = 64.2 \approx 65\text{m}$$

$$\text{Cost of fencing} = 65 \times \text{£}14 = \text{£}910$$

### Trigonometry

#### Stretch it!

Opposite could have been 1 m, hypotenuse could have been 2 m. They could be any lengths that keep opposite and hypotenuse in the ratio 1:2.

- 1 a 0.4      b 0.6      c 1.0  
 d 26.6      e 48.6      f 54.7

2  $\cos 72^\circ = \frac{MN}{15}$        $MN = 15 \cos 72^\circ = 4.6 \text{ cm}$

3  $\tan ABC = \frac{6}{7}$   
 $ABC = \tan^{-1}\left(\frac{6}{7}\right)$   
 $ABC = 40.6^\circ$

4 Let  $x$  be the depth of water.  
 $\sin 15^\circ = \frac{x}{10}$   
 $x = 10 \sin 15^\circ$   
 $x = 2.6 \text{ m}$

### Exact trigonometric values

- 1 a 0.5      b 0      c 0  
 d  $\frac{1}{\sqrt{2}}$       e  $\sqrt{3}$

2  $\tan 45^\circ = 1 = \frac{\text{opposite}}{\text{adjacent}} = \frac{4}{AC}$

Therefore  $AC = 4 \text{ cm}$

$\cos 45^\circ = \frac{1}{\sqrt{2}} = \frac{4}{BC}$

$BC = 4\sqrt{2}$

Therefore  $BC = 4\sqrt{2} \text{ cm}$

- 3 Since:  $\tan 30^\circ = \frac{1}{\sqrt{3}}$  one angle must be  $30^\circ$  and therefore the other is  $60^\circ$


4  $\sin 30^\circ = \frac{1}{2}$  therefore  $ABC = 30^\circ$

5  $\cos 30^\circ = \frac{\sqrt{3}}{2} = 0.866$  (3 d.p.)       $\tan 45^\circ = 1$   
 Smallest to largest =  $0.5, \frac{3}{4}, \cos 30^\circ, \tan 45^\circ$


### Transformations

#### Stretch it!

Yes. Reflection in the  $x$  axis followed by reflection in the  $y$  axis (or vice versa) will always produce a rotation of  $180^\circ$ .




- 2 Translation by vector  $\begin{pmatrix} -4 \\ -2 \end{pmatrix}$   
 3 a and b



- 4 Reflection in the  $y$ -axis  
 5 Enlargement by scale factor  $\frac{1}{2}$ , centre (3, 3)

6 a and b



c Rotation of 90° clockwise about (0, 0)

Similar shapes

Stretch it!

Perimeter of ABC = 3 + 6 + 5 = 14 cm






Perimeter of DEF = 6 + 12 + 10 = 28 cm

The perimeter of a shape enlarged by scale factor 2 will also be enlarged by scale factor 2.

In general, all lengths on an enlarged shape, including the perimeter, are enlarged by the same scale factor.


- 1 a Angle DFE = 30° (Corresponding angles are the same)
- b Scale factor of enlargement =  $\frac{\text{enlarged length}}{\text{original length}} = \frac{12}{3} = 4$   
Length of EF = 4 cm × 4 = 16 cm
- c Length of AB = 8 cm ÷ 4 = 2 cm
- 2 a Angle MLO = 80° (Corresponding angles are the same: angle MLO = angle QPS)
- b Scale factor of enlargement =  $\frac{\text{enlarged length}}{\text{original length}} = \frac{9}{3} = 3$   
Length of QR = 4.4 cm × 3 = 13.2 cm
- c Length of LO = 12 cm ÷ 3 = 4 cm

Vectors




- c  $\mathbf{a} = \begin{pmatrix} 4 \\ 3 \end{pmatrix}$   
So  $-2\mathbf{a} = -2 \times \begin{pmatrix} 4 \\ 3 \end{pmatrix} = \begin{pmatrix} -8 \\ -6 \end{pmatrix}$
- 3 a  $\mathbf{a} + \mathbf{b} = \begin{pmatrix} 2 \\ 6 \end{pmatrix} + \begin{pmatrix} -5 \\ -3 \end{pmatrix} = \begin{pmatrix} -3 \\ 3 \end{pmatrix}$
- b  $\mathbf{a} + 2\mathbf{b} = \begin{pmatrix} 2 \\ 6 \end{pmatrix} + 2 \times \begin{pmatrix} -5 \\ -3 \end{pmatrix} = \begin{pmatrix} 2 \\ 6 \end{pmatrix} + \begin{pmatrix} -10 \\ -6 \end{pmatrix} = \begin{pmatrix} -8 \\ 0 \end{pmatrix}$
- c  $\mathbf{a} - \mathbf{b} = \begin{pmatrix} 2 \\ 6 \end{pmatrix} - \begin{pmatrix} -5 \\ -3 \end{pmatrix} = \begin{pmatrix} 7 \\ 9 \end{pmatrix}$
- d  $\mathbf{b} - 2\mathbf{a} = \begin{pmatrix} -5 \\ -3 \end{pmatrix} - 2 \times \begin{pmatrix} 2 \\ 6 \end{pmatrix} = \begin{pmatrix} -5 \\ -3 \end{pmatrix} - \begin{pmatrix} 4 \\ 12 \end{pmatrix} = \begin{pmatrix} -9 \\ -15 \end{pmatrix}$
- 4 No, only Jordan is correct.  
 $\mathbf{a} = \begin{pmatrix} 2 \\ 3 \end{pmatrix}, \mathbf{d} = \begin{pmatrix} 20 \\ 30 \end{pmatrix}$   
Vectors  $\mathbf{a}$  and  $\mathbf{d}$  are parallel since  $10\mathbf{a} = 10 \times \begin{pmatrix} 2 \\ 3 \end{pmatrix} = \begin{pmatrix} 20 \\ 30 \end{pmatrix}$   
So Jordan is correct.  
 $\mathbf{a} = \begin{pmatrix} 2 \\ 3 \end{pmatrix}, \mathbf{b} = \begin{pmatrix} 4 \\ -6 \end{pmatrix}$   
Vectors  $\mathbf{a}$  and  $\mathbf{b}$  are not parallel as  $-2\mathbf{a} = \begin{pmatrix} -4 \\ -6 \end{pmatrix}$   
So Jenna is not correct.

Review it




- b BC = 3.8 cm
- c  $x = 50^\circ$
- 2 a 5 faces
- b 6 vertices



\*This answer differs from the one in the Revision Guide due to an error in our first edition. This answer has now been re-checked and corrected.



4 Area A =  $7 \times 6 = 42 \text{ cm}^2$



Area B =  $10 \times 3 = 30 \text{ cm}^2$


Total area =  $42 + 30 = 72 \text{ cm}^2$

5 Area of parallelogram =  $3 \times 12 = 36 \text{ cm}^2$

Length of side of square =  $\sqrt{36} = 6 \text{ cm}$

Perimeter of square =  $4 \times 6 = 24 \text{ cm}$

6



7 Rotation of  $180^\circ$  about  $(1, 0)$

8 Angle CFE =  $112^\circ$  (Corresponding angles are equal)

Angle CFG =  $180 - 112 = 68^\circ$  (Angles on a straight line add up to  $180^\circ$ )

Angle GCF = angle CFG (Base angles of an isosceles triangle are equal)

$x = (180 - 68 - 68) = 44^\circ$  (Angles in a triangle add up to  $180^\circ$ )

9 Shaded Area =  $(10 \times 12) - ((\frac{1}{2} \times 12 \times 3) + (\frac{1}{2} \times 8 \times 7) + (\frac{1}{2} \times 10 \times 4))$   
 $= 120 - (18 + 28 + 20)$   
 $= 54 \text{ cm}^2$

Proportion =  $\frac{54}{120} = \frac{9}{20} = 45\%$


10 If triangle ABC is right-angled,  $c^2 = a^2 + b^2$

$c^2 = 8^2 = 64$

$a^2 + b^2 = 4^2 + 6^2 = 16 + 36 = 52$

$c^2 \neq a^2 + b^2$  so triangle ABC is not right-angled.

11 a



Scale is 1 : 200

b Distance from A to YZ = 2.9 cm

$2.9 \times 200 = 5800 \text{ cm} = 5.8 \text{ m}$

Distance from A to YX = 3.3 cm

$3.3 \times 200 = 6600 \text{ cm} = 6.6 \text{ m}$

Difference in distance =  $6.6 - 5.8 = 0.8^* \text{ m}$

12 a  $\cos 45^\circ = \frac{1}{\sqrt{2}}$


b Ratio of adjacent to hypotenuse is 1:2

Therefore AB = 3 cm


13 a  $\mathbf{a} + 2\mathbf{b} = \begin{pmatrix} 4 \\ -5 \end{pmatrix} + 2 \times \begin{pmatrix} 2 \\ 3 \end{pmatrix}$   
 $= \begin{pmatrix} 4 \\ -5 \end{pmatrix} + \begin{pmatrix} 4 \\ 6 \end{pmatrix}$   
 $= \begin{pmatrix} 8 \\ 1 \end{pmatrix}$

b  $\mathbf{b} - 2\mathbf{a} = \begin{pmatrix} 2 \\ 3 \end{pmatrix} - 2 \times \begin{pmatrix} 4 \\ -5 \end{pmatrix}$   
 $= \begin{pmatrix} 2 \\ 3 \end{pmatrix} - \begin{pmatrix} 8 \\ -10 \end{pmatrix}$   
 $= \begin{pmatrix} -6 \\ 13 \end{pmatrix}$

14



15



\*This answer differs from the one in the Revision Guide due to an error in our first edition. This answer has now been re-checked and corrected.

- 16 a** i  $35^\circ$   
 ii Triangle WYZ is isosceles, and base angles of an isosceles triangle are equal.  
**b** Angles in a triangle add up to  $180^\circ$  so:  
 $b = 180 - 35 - 35$   
 $= 110^\circ$   
**c** Triangle XYZ is isosceles, and base angles of an isosceles triangle are equal so:  
 $c = (180 - 70) \div 2 = 55^\circ$


**17** Using Pythagoras' theorem  $c^2 = a^2 + b^2$ :

$$\begin{aligned} AC^2 &= AB^2 + BC^2 \\ 14^2 &= 6^2 + BC^2 \\ BC^2 &= 14^2 - 6^2 = 160 \\ BC &= \sqrt{160} \\ BC &= 12.6 \text{ cm (to 1 d.p.)} \end{aligned}$$

**18** Interior angle of a square =  $90^\circ$

Sum of interior angles of an octagon (with  $n = 8$ )  
 $= 180 \times (n - 2) = 180 \times (8 - 2) = 1080^\circ$   
 Interior angle of a regular octagon =  $1080^\circ \div 8 = 135^\circ$   
 (Or, exterior angle of a regular octagon =  $360^\circ \div 8 = 45^\circ$ .  
 Then interior angle =  $180^\circ - 45^\circ = 135^\circ$ )  
 Angles around a point add up to  $360^\circ$  so:  
 $x = 360 - 90 - 135$   
 $x = 135^\circ$

**19**



- 20** Divide the trapezium into a rectangle and a triangle. Draw a line DX parallel to AB, with X on the line BC.  $BX = 5 \text{ cm}$ ,  $CX = 7 \text{ cm}$ .  
 Using Pythagoras' theorem  $c^2 = a^2 + b^2$ :  
 $DC^2 = CX^2 + DX^2$   
 $DC^2 = 7^2 + 4^2 = 65$   
 $DC = \sqrt{65}$   
 $DC = 8.06$  (to 2 d.p.)  
 Perimeter of ABCD =  $4 + 5 + 8.06 + 12 = 29.06 \text{ cm}$   
**21** Length of arc =  $\frac{1}{4} \times 2 \times \pi \times 4 = 2\pi$   
 Perimeter =  $4 + (2 \times 9) + 4 + 2\pi = 32.3 \text{ cm}$   
**22** Area of square =  $6 \times 6 = 36 \text{ cm}^2$   
 Area of circle =  $\pi \times 3^2 = 9\pi \text{ cm}^2$   
 Shaded area =  $36 - 9\pi = 7.7 \text{ cm}^2$   
**23** Volume of cylinder =  $\pi \times 3^2 \times 15 = 135\pi \text{ cm}^3$   
 $2 \text{ litres} = 2000 \text{ ml} = 2000 \text{ cm}^3$   
 $2000 \div 135\pi = 4.7$   
 Glass can be completely filled 4 times.

**24** Using Pythagoras' theorem  $c^2 = a^2 + b^2$ :

$$\begin{aligned} PR^2 &= PQ^2 + RQ^2 \\ PR^2 &= 10^2 + 6^2 = 136 \\ PR^2 &= PS^2 + SR^2 \\ 136 &= 11^2 + x^2 \\ x^2 &= 136 - 11^2 = 136 - 121 = 15 \\ x &= \sqrt{15} \\ x &= 3.87 \text{ cm (to 3 s.f.)} \end{aligned}$$

**25 a** Curved surface area =  $\pi \times 6 \times 10 = 60\pi \text{ cm}^2$

$$\text{Base area} = \pi \times 6^2 = 36\pi \text{ cm}^2$$

$$\text{Total surface area} = 60\pi + 36\pi = 96\pi = 300 \text{ cm}^2 \text{ to 2 s.f.}$$

$$\text{b Volume} = \frac{1}{3} \times \pi \times 6^2 \times 8 = 96\pi = 300 \text{ cm}^3$$

**26**  $\tan x = \frac{8}{6}$   
 $x = \tan^{-1}\left(\frac{8}{6}\right)$   
 $x = 53.1^\circ$


**27** Translation by vector  $\begin{pmatrix} -7 \\ -6 \end{pmatrix}$

## Probability

### Basic probability

**Stretch it!** No – each time the probability of getting an even number is  $\frac{1}{2}$ . You would expect to get even numbers approximately 50 times but cannot guarantee it.

**1**  $\frac{4}{10}$



**2 a** Total number of sweets =  $12 + 3 + 10 = 25$

$$\frac{3}{25}$$

$$\text{b } \frac{(3+10)}{25} = \frac{13}{25}$$

- 3** Pair **a**, because when you flip a coin, you can't get both a head and a tail at the same time. (Prime numbers on a dice are 2, 3, 5 and odd numbers are 1, 3, 5, so events **b** are **not** mutually exclusive because 3 is in both groups.)  
**4** Pair **b**, because the first sweet chosen is replaced, so the possible outcomes of the second choice remain the same. (If the first sweet chosen is eaten, the possible outcomes of the second choice are altered, and so events **a** are **not** independent.)  
**5**  $P(6) = 1 - (0.1 + 0.15 + 0.1 + 0.02 + 0.2)$   
 $= 1 - 0.57 = 0.43$   
**6**  $P(\text{green or red}) = 1 - 0.4 = 0.6$   
 $P(\text{green}) = 2 \times P(\text{red})$   
 $P(\text{red}) = \frac{0.6}{3} = 0.2$   
 $P(\text{green}) = 2 \times 0.2 = 0.4$

### Two-way tables and sample space diagrams

**1**

	Chicken	Beef	Vegetarian
Fruit	12	6	4
Cake	5	3	8
Total	17	9	12

- a 12 (this is worked out by using the numbers in the 'Total' row, which must add up to 38)  
 b As shown in the table.

2 a


		Dice 1					
		1	2	3	4	5	6
Dice 2	1	2	3	4	5	6	7
	2	3	4	5	6	7	8
	3	4	5	6	7	8	9
	4	5	6	7	8	9	10
	5	6	7	8	9	10	11
	6	7	8	9	10	11	12

- b i  $\frac{2}{36} = \frac{1}{18}$   
 ii  $\frac{3}{36} = \frac{1}{12}$   
 iii 0

- 3 To score 6, the player must pick two cards showing 3. To score 2, the player must pick two cards showing 1. Since the probability of getting 3 and 3 is more than 0, and the probability of getting 1 and 1 is more than 0, there must be at least 2 of each of those numbers. So the cards must be 1, 1, 3, 3.

### Sets and Venn diagrams

Stretch it! None




- b  $A \cap B = \{\text{multiples of 6 less than 20}\}$  because these numbers are multiples of both 2 and 3.

- 2 a  $C \cap T$  is the set of students who travel by car AND train

$C' \cap B$  is the set of students who do NOT travel by car AND travel by bus.\*


- b i  $P(C) = \frac{(14 + 11 + 11 + 2)}{(14 + 11 + 11 + 2 + 17 + 19 + 26)} = \frac{38}{100} = \frac{19}{50}$   
 ii  $P(B \cup T) = \frac{(19 + 11 + 2 + 0 + 11 + 17)}{100} = \frac{60}{100} = \frac{3}{5}$   
 iii  $P(B' \cap T) = \frac{(11 + 17)}{100} = \frac{28}{100} = \frac{7}{25}$

- 3  $P(A \cap B) = \frac{3}{20}$  so there must be 3 elements in the intersection.  
 $P(A) = \frac{3}{10} = \frac{6}{20}$  so there must be a total of 6 elements in A.  
 The total number of elements must sum to 20.




### Frequency trees and tree diagrams

- 1 a Apple = 123  
 Pear =  $347 - 123 = 224$   
 Apple fruiting = 102  
 Apple not fruiting =  $123 - 102 = 21$   
 Pear not fruiting = 34  
 Pear fruiting =  $224 - 34 = 190$



- 2 a  $\frac{190}{347}$




b  $\frac{3}{5} \times \frac{3}{5} = \frac{9}{25}$

c  $\frac{3}{5} \times \frac{3}{5} = \frac{9}{25}$

$\frac{2}{5} \times \frac{3}{5} = \frac{6}{25}$

$\frac{3}{5} \times \frac{2}{5} = \frac{6}{25}$

$P(\text{at least one prime}) = 1 - P(\text{no primes})$   
 $= 1 - \frac{2}{5} \times \frac{2}{5} = 1 - \frac{4}{25} = \frac{21}{25}$



b  $P(\text{two glass marbles}) = \frac{3}{7} \times \frac{2}{6} = \frac{6}{42}$

$P(\text{glass then plastic}) = \frac{3}{7} \times \frac{4}{6} = \frac{12}{42}$

\*This answer differs from the one in the Revision Guide due to an error in our first edition. This answer has now been re-checked and corrected.

$$P(\text{plastic then glass}) = \frac{4}{7} \times \frac{3}{6} = \frac{12}{42}$$

$$\begin{aligned} P(\text{at least one glass}) &= 1 - P(\text{both plastic}) \\ &= 1 - \frac{4}{7} \times \frac{3}{6} = 1 - \frac{12}{42} \\ &= 1 - \frac{2}{7} = \frac{5}{7} \end{aligned}$$

### Expected outcomes and experimental probability

**Stretch it!** The dice has not been rolled enough times to decide if it is biased. More tests need to be carried out.

1  $0.45 \times 300 = 135$

2 Red =  $\frac{2}{10} = \frac{1}{5}$   
 $\frac{1}{5} \times 100 = 20$  red sweets


3  $\frac{1}{2} \times 100 = 50$  primes

4 a Charlie – he has carried out the most tests.

b  $\frac{(112 + 10 + 28)}{(112 + 10 + 28 + 74 + 7 + 19)} \times 10 = 6$

### Review it!

1  $0.12 \times 250 = 30$



3  $1 - 0.3 = 0.7$

4 B, C

5 a  $\frac{3}{5}$

b  $(\frac{1}{5}) \times 25 = 5$

6

	Pizza	Pasta	Risotto	Total
Cake	12	6	1	19
Ice Cream	10	11	10	31
Total	22	17	11	50

7  $0.2 + 5x + 0.2 + x = 1$

$$6x + 0.4 = 1$$

$$x = 0.1$$

$$P(\text{white}) = 5x + 0.2 = 0.7$$

8 a No, he has not tested his dice enough times.

b  $P(2) = \frac{9}{(12 + 9 + 16 + 7 + 6 + 0)} = \frac{9}{50}$

$$\frac{9}{50} \times 100 = 18$$

9  $P(R, R) = 0.1 \times 0.5 = 0.05$

$$P(R, G) = 0.1 \times 0.5 = 0.05$$

$$P(G, R) = 0.9 \times 0.5 = 0.45$$

$$0.05 + 0.05 + 0.45 = 0.55$$

Or  $P(\text{at least one red}) = 1 - P(\text{green, green})$

$$= 1 - (0.9 \times 0.5)$$

$$= 1 - 0.45$$

$$= 0.55$$

10 a

		Dice					
		1	2	3	4	5	6
Coin	Heads	2	4	6	8	10	12
	Tails	3	4	5	6	7	8


b i  $\frac{2}{12} = \frac{1}{6}$

ii  $\frac{2}{12} = \frac{1}{6}$

11 a i 6      ii 1      iii 5

b  $\frac{4}{8} = \frac{1}{2}$

12 a



b  $\frac{7}{9} \times \frac{6}{8} = \frac{42}{72} = \frac{7}{12}$

13 a Possible fractions:  $\frac{1}{2}, \frac{1}{3}, \frac{1}{4}, \frac{2}{3}, \frac{2}{4}, \frac{3}{4}$

Less than  $\frac{1}{2}$  are  $\frac{1}{3}$  and  $\frac{1}{4}$ .

$$P(\text{less than } \frac{1}{2}) = \frac{2}{6} = \frac{1}{3}$$

$$\frac{1}{3} \times 30 = 10$$

b  $\frac{1}{3}$  is only a theoretical probability and therefore will not necessarily be accurate in real life.

14 45% of 300 = 135

135 boys and 165 girls.

$$\frac{2}{3} \text{ of } 135 = 90$$

$$\frac{4}{5} \text{ of } 165 = 132$$

Total playing sport = 222

$$\text{Probability} = \frac{222}{300} = \frac{37}{50} = 0.74$$

15  $P(\text{hooking a winning duck}) = \frac{5}{20} = 0.25$

If 100 people play, expected number of winners =  $0.25 \times 100 = 25$  people.

The game makes  $\text{£}1 \times 100$  people =  $\text{£}100$ .

The money paid out in prizes = 25 winners  $\times$   $\text{£}2 = \text{£}50$

Profit =  $\text{£}100 - \text{£}50 = \text{£}50$

16 a Milo will have the better estimate as he has surveyed a greater number of people.

b Number of left-handed students =  $5 + 4 + 7 + 7 = 23$

Number of right-handed students =  $23 + 18 + 51 + 60 = 152$

$$P(\text{left-handed}) = \frac{23}{23 + 152} = \frac{23}{175}$$

$$\frac{23}{175} \times 2000 = 262.8$$

You would expect to find 263 left-handed students in a school with 2000 students.

## Statistics

### Data and sampling

**Stretch it!** A random sample could be taken; you could allocate a number to each pupil and randomly generate the numbers to survey. Any method is acceptable as long as each person in the school has an equally likely chance of being chosen. Alternatively a stratified sample could be taken.

- Primary source: Recording the data by measuring it yourself.  
Secondary source: Any sensible source, e.g. the Meteorological Office, local paper etc.
- Qualitative data.
- It is cheaper and quicker than surveying the whole population.
- The people working for an animal charity are more likely to be opposed to wearing real fur; every member of the population does not have an equal chance of being chosen.
  - Surveying people in the street, a random telephone survey, any sensible method that ensures that any member of the population has an equal chance of being chosen.
- $\frac{400}{2000} = \frac{1}{5}$
  - $\frac{1}{5} \times 50 = 10$  bottles
- $\frac{3}{200} \times 800\,000 = 12\,000$
  - The sample is relatively small, the sample is not a random sample as it is taken on one day in a year.

### Frequency tables

1

Number of people on the bus	Frequency
0*–9	4
10–19	12
20–29	3
30–39	1

2 a

Number of courgettes	Frequency
0	1
1	0
2	1
3	1
4	9
5	3
6	0


- $(0 \times 1) + (1 \times 0) + (2 \times 1) + (3 \times 1) + (4 \times 9) + (5 \times 3) = 56$
- 3 There are gaps between his groups – where would he record someone who spent 15.5 hours training? His groups do not have the same width.

### Bar charts and pictograms

- $15 + 4 + 1 = 20$
  - $4 + 1 = 5$   
 $\frac{5}{20} \times 100 = 25\%$

- $11 - 7 = 4$
  - Total number of people surveyed =  $18 + 18 + 12 + 3 = 51$   
Total number of boys =  $11 + 6 + 3 = 20$   
 $\frac{20}{51} \times 100 = 39.2\%$
  - Proportion of boys who played two sports =  $\frac{6}{18} = \frac{1}{3}$   
Proportion of boys who played three sports =  $\frac{3}{12} = \frac{1}{4}$   
 $\frac{1}{3} > \frac{1}{4}$  so Jasmine is incorrect, and you can't prove that what she says is true.\*

- $50 - (11 + 15) = 24$   
 $24 \div 3 = 8$   
 Therefore:  $2 \times 8 = 16$  size 7 shoes  
 $1 \times 8 = 8$  size 8 shoes




- $90 - 20 = 70$
  - Total number of bikes =  $50 + 50 + 20 + 90 = 210$   
 $\frac{50}{210} = \frac{5}{21}$

### Pie charts

#### Stretch it!

Round appropriately – but check the angles sum to  $360^\circ$

- $27 + 42 + 21 = 90$   
 $360^\circ \div 90 = 4^\circ$   
 French =  $27 \times 4 = 108^\circ$   
 Spanish =  $42 \times 4 = 168^\circ$   
 German =  $21 \times 4 = 84^\circ$



- $\frac{1.5}{360} \times 240 = 1$  student earned more than £40 000.
  - $\frac{288 + 63}{360} \times 100 = 97.5\%$  of students earned less than £30 000.
- $18 + 10 = 28$
  - The bar chart, since the frequency is easy to read from the bar chart.

\*This answer differs from the one in the Revision Guide due to an error in our first edition. This answer has now been re-checked and corrected.

## Stem and leaf diagrams

- 1 **a** 7  
**b**  $4.3 - 4.1 = 0.2$  kg
- 2 Age of people using a dentist

2	0 0 0 0 1 1 1
3	2 5 5 7
4	1 2 2 6

The leaves were not in ascending order, the spaces between leaves were not regular.

- 3 **a** Stem and leaf diagram – you can see the smallest number of passengers was 3; however, on the bar chart you only know it is between 0 and 9.  
**b** Both since the shape of the data is preserved in both.

## Measures of central tendency: mode

- 1 The other three must be 12.2.  
 2  $1 < t \leq 2$   
 3 17

## Measures of central tendency: median

- 1 Ordering the data gives; 2.9, 3.1, 4.3, 6.5, 8.7, 9.2  
 Median =  $\frac{4.3 + 6.5}{2} = 5.4$
- 2  $29 + 28 + 30 + 3 + 10 = 100$   
 $\frac{(100 + 1)}{2} = 50.5$  – median term is between the 50th and 51st terms.  
 Both these lie in the  $2 \leq b < 4$  class.
- 3 **a** Group A =  $\frac{(82 + 85)}{2} = 83.5$   
 Group B =  $\frac{(75 + 79)}{2} = 77$   
**b** Group A has a higher median, so they did better on the test.

## Measures of central tendency: mean

**Stretch it!** **a** mode **b** mean/median **c** mean/median

- 1 **a** Total frequency =  $12 + 3 + 5 = 20$   
 Mean =  $\frac{(2 \times 12) + (6 \times 3) + (10 \times 5)}{20} = 4.6$   
**b** You are using the midpoint of the groups as an estimate of the actual value for each group.
- 2  $\frac{(5 \times 9) + 6}{5 + 1} = 8.5$
- 3 No – they could be any pair of numbers which sum to 10.

## Range

- 1  $9.5 - 0.7 = 8.8$   
 2 Girls =  $18 - 15 = 3$   
 Boys =  $18 - 16 = 2$


\*This answer differs from the one in the Revision Guide due to an error in our first edition. This answer has now been re-checked and corrected.

- 3 Range for Athlete A =  $15.2 - 13.0 = 2.2$   
 Range for Athlete B =  $15.2 - 14.3 = 0.9$   
 Athlete A has the greatest range.
- 4  $45\% - 10 = 35\%$  or  $45\% + 30 = 75\%$


## Comparing data using measures of central tendency and range

- 1 **a** **i** Mean =  $\frac{(32 + 29 + 18 + 41 + 362 + 19)}{6} = \frac{501}{6} = 83.5$  minutes  
**ii** Ordered data: 18, 19, 29, 32, 41, 362  
 Median =  $\frac{(29 + 32)}{2} = 30.5$  minutes
- b** The extreme value (362 mins) affects the mean but not the median.
- 2 All the data is used to find the mean.
- 3 Either as long as suitably justified:  
 Car A – although the mean time is higher, it is more consistent in performance since the range is smaller.  
 Car B – the acceleration is quicker on average.
- 4 **a** and **b** The mode or median since the mean will not be a whole number and therefore not meaningful.

## Time series graphs


- 1 
- 2 **a**  $67^\circ\text{C}$   
**b** Approx.  $27^\circ\text{C}$   
**c** No, since it is extrapolation (beyond the limits of the data).
- 3 **a** 17 000 **b** **i** April **ii** August  
**c** The number of tourists peaks in April and again in December. The low seasons are February/March and July/August/September/October.

## Scatter graphs

- 1 **a** and **b**\* 
- c** Positive

- d This will vary according to the line of best fit: approximately 4.7 kg. A range of 4.6kg to 4.8kg would be acceptable.
- e This is beyond the limits of the data and therefore extrapolation.

2 a and b




- c The seeds failed to germinate or the seedling died.
  - d The further the seedling is from the light source the shorter its height.
- 3 No, although the two things correlate one does not cause another. There may be many reasons why the crime rate is high in the area, perhaps there is poverty and inequality causing social tension.

Review it!


- 1 The sample is too small and he only asked his friends. His data is therefore not representative of the population of TV viewers.
- 2 a Margherita  
b Total frequency =  $11 + 2 + 6 + 1 = 20$   
 $\frac{1}{20} = \frac{5}{100} = 5\%$   
c  $360^\circ \div 20 = 18^\circ$   
Pepperoni =  $1 \times 18^\circ = 18^\circ$   
(or  $5\%$  of  $360^\circ = 18^\circ$ )
- 3 a  $\frac{90}{360} = \frac{1}{4}$   
b  $45^\circ = \frac{1}{8}$  of  $360^\circ$   
Therefore  $\frac{1}{8}$  of the pie chart represents 60 cars.  
The whole pie chart =  $8 \times 60 = 480$  cars  
c  $(\frac{105}{360}) \times 480 = 140$  cars
- 4 a The number of people doing their grocery shopping online is increasing.  
b Any sensible answer, approximately 75%  
c No – it is outside the limits of the data therefore extrapolation.
- 5 a Outside: i Mode = 21 and 31 ii Median =  $\frac{(28 + 29)}{2} = 28.5$  iii Range =  $41 - 20 = 21$   
Greenhouse: i Mode = 47 ii Median =  $\frac{(47 + 47)}{2} = 47$  iii Range =  $51 - 37 = 14$

- b The seedlings are taller in the greenhouse since both mode and median is larger, the range of data is smaller in the greenhouse so the height the seedlings reach is more consistent.
- c Range =  $51 - 20 = 31$

6 a Comparative bar chart or compound bar chart:



Or:



- b Total number of students =  $(3 + 5 + 1 + 0 + 8 + 2 + 2 + 4) = 25$   
Number of cats =  $3 + 8 = 11$   
 $\frac{11}{25}$
- 7 a Total frequency =  $17 + 2 + 32 + 23 + 9 = 83$   
Median value =  $\frac{(83 + 1)}{2} = 42$ nd term  
42nd term is in group 40–59  
Median class = 40–59  
b The youngest person is between 0 and 19, the youngest may be any age in this range and the oldest is between 80 and 99 therefore any age in this range.
- 8 a 7  
b Size 5  
c Mean =  $\frac{(3 \times 2) + (4 \times 1) + (5 \times 7) + (6 \times 5) + (7 \times 3)}{2 + 1 + 7 + 5 + 3} = 5.3$   
d Mode – the mean is not an actual shoe size.
- 9 a Time for 800 m (seconds)  
11 | 2 2 5 8 9  
12 | 0 1 9  
13 | 1 2  
Key: 11|2 = 112 seconds  
b  $\frac{4}{10} = \frac{2}{5}$ \*


\*This answer differs from the one in the Revision Guide due to an error in our first edition. This answer has now been re-checked and corrected.

10 a  $\frac{50}{150} = \frac{1}{3}$

b  $60 - 40 = 20$

c Biology

11 a\* and c



b Negative

d Approximately 40 minutes: it depends on line of best fit.

e This is outside the limits of the data and therefore extrapolation.

f As the age of the customer increases the time spent on the phone decreases.

12 a  $\frac{(65 \times 3) + (75 \times 5) + (85 \times 2)}{3 + 5 + 2} = 74 \text{ kg}$

b The midpoint of the class is used as the age of each of the patients rather than the actual age.


13 Male  $< 50 = \left(\frac{12\,201}{36\,579}\right) \times 120 = 40$

Female  $< 50 = \left(\frac{10\,678}{36\,579}\right) \times 120 = 35$

Male  $\geq 50 = \left(\frac{5\,699}{36\,579}\right) \times 120 = 19$

Female  $\geq 50 = \left(\frac{8\,001}{36\,579}\right) \times 120 = 26$

14 Annual income for surveyed population



15 Mean  $= \frac{(10.3 \times 10) + 9.5}{11} = 10.2$  (1 d.p.)

16 Mean is 3.8 so the sum of the scores is  $3.8 \times 5 = 19$

Mode is 3 so she must roll at least two 3s.

Range is 4.

If the range is 4 then the lowest and highest must be either 1 and 5 or 2 and 6.

The numbers are: 2, 3, 3, 5 and 6



# Edexcel Foundation Mathematics Exam Practice Book

## Full worked solutions

### Number

#### Factors, multiples and primes

- 1  $18 = 2 \times 3 \times 3$   
 $24 = 2 \times 2 \times 2 \times 3$   
Find the factors that they both share (2 and 3) and multiply together:  
 $2 \times 3 = 6$ , so HCF is 6.
- 2 17, 19, 23 are the only numbers in this range with only 2 factors (1 and the number itself).
- 3  $60 = 20 \times 3$   
 $= 2 \times 2 \times 5 \times 3$   
 $= 2^2 \times 3 \times 5$
- 4 Drummer 1 hits her drum at: 6 12 18 24 30 36 42 48 54 60 seconds  
Drummer 2 hits his drum at: 8 16 24 32 40 48 56 seconds  
They hit their drums at the same time twice (two times): after 24 seconds and after 48 seconds.

#### Ordering integers and decimals

- 1 Negative numbers are smaller than zero  
-12 is further left on the number line than -8, -1 is larger than -8 (and -12) so it appears next.  
Then comes 0, then 2.  
So the order is:  
-12, -8, -1, 0, 2
- 2 First look at the place value for 10ths: 0.32 and 0.3 have the higher number of 10ths.  
Now compare their 100ths. 0.32 has 2 100ths but 0.3 doesn't have any, so it's smaller.  
Similarly, 0.23 and 0.203 both have 2 10ths, but 0.23 is bigger than 0.203 because it has 3 100ths while 0.203 only has 3 1000ths.  
So the order is:  
0.32, 0.3, 0.23, 0.203
- 3 a  $-4 < 0.4$  (the negative number is smaller)  
b  $4.200 < 4.3$  (the larger number has more 10ths)  
c  $-0.404 > -0.44$  (because they are both negative, the one with 'more' 100ths is smaller)  
d  $0.33 < 0.4$  (the larger number has more 10ths)

#### Calculating with negative numbers

- 1 a  $-7 + -3 = -7 - 3 = -10$   
b  $-7 - -3 = -7 + 3 = -4$   
c  $8 + -5 - -2 = 8 - 5 + 2 = 5$   
d  $-4 - -6 + -1 = -4 + 6 - 1 = 1$
- 2 a -18  
b  $-12 \div -3 = 12 \div 3 = 4$   
c  $-4 \times -2 \times 5 = 4 \times 2 \times 5 = 40$   
d  $(-24 \div 3) \times 2 = -8 \times 2 = -16$
- 3 At 10pm,  $2 - 4 = -2^\circ\text{C}$ . At 2am,  $-2 - 5 = -7^\circ\text{C}$
- 4 Let  $a$  = number of correct answers,  $b$  = number of incorrect answers  
 $3a - 2b = -5$  (1)  
There are five questions, so  $a + b = 5$  and  $b = 5 - a$  (2)  
Substituting this for  $b$  in (1):  $3a - 2(5 - a) = -5$   
 $3a - 10 + 2a = -5$   
 $5a = 5$   
 $a = 1$   
Substituting this in (2):  
 $b = 5 - 1$   
 $b = 4$   
Sally got 1 correct answer and 4 incorrect answers in the test.

### Multiplication and division

- 1 a 
$$\begin{array}{r} 357 \\ \times 6 \\ \hline 2142 \end{array}$$
- b 
$$\begin{array}{r} 261 \\ \times 43 \\ \hline 783 \\ 10440 \\ \hline 11223 \end{array}$$
  
 $(= 261 \times 3)$   
 $(= 261 \times 40)$
- c 
$$\begin{array}{r} 092 \\ 6 \overline{)5512} \\ \underline{30} \\ 251 \\ \underline{18} \\ 712 \\ \underline{42} \\ 292 \\ \underline{18} \\ 112 \end{array}$$
  
92
- d 
$$\begin{array}{r} 052 \\ 13 \overline{)676} \\ \underline{39} \\ 286 \\ \underline{26} \\ 26 \end{array}$$
  
52
- 2 a 
$$\begin{array}{r} 012 \text{ remainder } 12 \\ 24 \overline{)3060} \end{array}$$
  
So 12 boxes are filled.
- b  $24 \times 12 = 20 \times 12 + 4 \times 12 = 240 + 48 = 288$   
 $300 - 288 = 12$   
There are 12 books left over.
- 3  $12500 - 440 = 12060$
- $$\begin{array}{r} 00335 \\ 36 \overline{)121060} \\ \underline{108} \\ 1216 \\ \underline{108} \\ 0180 \end{array}$$
- Each repayment is £335.
- 4  $52 - 6 = 46$  weeks     $46 \times 26 = 1196$  hours

#### Calculating with decimals

- 1 First note the combined number of decimal places in both numbers (2).  
Remove the decimal points to do the calculation:
- $$\begin{array}{r} 92 \\ \times 83 \\ \hline 276 \\ 7360 \\ \hline 7636 \end{array}$$
- Now you've got the digits right put the decimal point back, counting in from the right 2 places, to give a number with 2 decimal places:  
76.36
- 2 
$$\begin{array}{r} 22.50 \\ + 19.99 \\ \hline 42.49 \end{array}$$
- $$\begin{array}{r} 50.00 \\ - 42.49 \\ \hline 07.51 \end{array}$$
- She should get £7.51 change.
- 3 
$$\begin{array}{r} 038.29 \\ 6 \overline{)229.74} \end{array}$$

4 Kirsty raises  $\frac{172.50}{5+1}$

Kirsty raises  $\pounds 28.75 \times 5 = 28.75 \times 10 \div 2 = 287.5 \div 2 = 143.7$

$$\begin{array}{r} 028.75 \\ 6 \overline{)172.50} \end{array}$$

Kirsty raises  
 $\pounds 143.75$

$$\begin{array}{r} 61141 \\ 172.50 \\ -143.75 \\ \hline 028.75 \end{array}$$

Flo raises  $\pounds 28.75$

### Rounding and estimation

- a The first non-zero digit is 7, so round the digit after the 8. This is below 5, so the 8 doesn't change.  
0.798

b Look at the digit in the third decimal place. It is 5 or above (8), so round the 9 up to 10 and the 7 up to 8.  
0.80
- $\frac{9.74 \times 4.02}{7.88} \approx \frac{10 \times 4}{8} = 5$
- a  $40 \times 500 = 20\,000$   
 $20\,000 - 12\,500 = \pounds 7\,500$

b Overestimate, because the concert ticket price and number of tickets sold were rounded up, and so the amount of income was estimated more than it really is.

### Converting between fractions, decimals and percentages

- a There are  $0 \times 10^{\text{ths}}$ ,  $7 \times 100^{\text{ths}}$  and  $1 \times 1000^{\text{ths}}$   
 $= 071 \times 1000^{\text{ths}}$  so:  $\frac{71}{1000}$

b  $63 \div 100 = 0.63$

c  $0.4 \times 100 = 40\%$

d  $32\% = \frac{32}{100} = \frac{8}{25}$
- a  $5 \div 16 = 0.3125$

b To convert a number to a percentage, multiply its decimal value by 100.  
 $0.3125 \times 100 = 31.25\%$
- $\frac{5}{8} = 0.625$ ,  $60\% = 0.6$ , so 0.65 is the largest.

### Ordering fractions, decimals and percentages

- a  $\frac{1}{2} = \frac{5}{10} = 0.5$ , so  
 $\frac{1}{2} < 0.6$

b  $\frac{3}{4} = 3 \div 4 = 0.75$ , so  
 $\frac{3}{4} > 0.7$

c  $\frac{-3}{10} = -0.3$ , so  
 $\frac{-3}{10} < 0.2$
- a LCM of 12, 15 and 20 is 60  
 $\frac{5}{12} = \frac{25}{60}$   
 $\frac{7}{15} = \frac{28}{60}$   
 $\frac{9}{20} = \frac{27}{60}$

So order from lowest to highest is  $\frac{5}{12}$ ,  $\frac{9}{20}$ ,  $\frac{7}{15}$

b  $45\% = \frac{45}{100} = 0.45$   
 $\frac{1}{25} = \frac{4}{100} = 0.04$   
 $0.04 < 0.4 < 0.45$   
 So order is:  
 $\frac{1}{25}$ , 0.4, 45%

- Shop C is cheapest ( $\frac{2}{5} = 40\%$ ), then Shop A ( $\frac{1}{3} = 33.3\% \dots$ ), and Shop B offers the least discount at 30%.

4  $\frac{5}{9} = 0.5$

$38.5\% = 0.385$

$\frac{3}{10} = 0.3$

So the order is  $\frac{5}{9}$ , 38.5%, 0.38,  $\frac{3}{10}$

### Calculating with fractions

- $\frac{1}{5} + \frac{4}{9} = \frac{9}{45} + \frac{20}{45} = \frac{9+20}{45} = \frac{29}{45}$
- $2\frac{3}{4} - 2\frac{2}{3} = \frac{11}{4} - \frac{8}{3} = \frac{33}{12} - \frac{32}{12} = \frac{1}{12}$
- $1\frac{5}{6} \times \frac{2}{7} = \frac{11}{6} \times \frac{2}{7} = \frac{22}{42} = \frac{11}{21}$
- $6 \div \frac{3}{5} = 6 \times \frac{5}{3} = \frac{30}{3} = 10$

Jo can make 10 necklaces.

### Percentages

- $\frac{40}{100} \times 25 = 10$
- $16 \times 0.85 = \pounds 13.60$
- $12450 \times 1.14 = 14193$
- $40 \times 7 \times 3 = \pounds 840$   
 $840 \times 1.2 = \pounds 1008$

### Order of operations

- $3^2 - 6 \div (2 + 1) = 9 - \frac{6}{3} = 9 - 2 = 7$
- $2^3 + 3x\sqrt{25} = 8 + (3 \times 5) = 8 + 15 = 23$
- $(1.7 - 0.12)^2 + \sqrt[3]{4.096} = 4.0964$

### Exact solutions

- Area of triangle =  $\frac{1}{2} \times \text{base} \times \text{vertical height} = 0.5 \times 0.76 \times 0.35 = 0.133$
- $(1\frac{1}{3})^2 = (\frac{4}{3})^2 = \frac{16}{9} = 1\frac{7}{9} \text{ m}^2$
- $\sqrt{2} \times \sqrt{6} = \sqrt{12} = 2\sqrt{3}$
- Area of a circle =  $\pi r^2$   
 The fraction of the circle shown =  $\frac{3}{4}$   
 The area of the circle shown =  $\frac{3}{4} \times \pi r^2$   
 The radius = 2 cm  
 So area of shape shown =  $\frac{3}{4} \times \pi \times 2^2 = 3\pi \text{ cm}^2$

### Indices and roots

- a  $7 \times 7 \times 7 \times 7 = 7^4$

b  $\frac{1}{5 \times 5 \times 5} = \frac{1}{5^3} = 5^{-3}$
- a  $2^4 = 2 \times 2 \times 2 \times 2 = 16$

b  $10^{-2} = \frac{1}{10^2} = \frac{1}{100}$
- $2^3 = 2 \times 2 \times 2 = 8$   
 $3^{-2} = \frac{1}{9}$   
 $\sqrt[3]{27} = 3$  or  $-3$   
 $\sqrt{25} = 5$  or  $-5$

Assuming the square roots are positive, the answer is:  
 $3^{-2}$ ,  $\sqrt[3]{27}$ ,  $\sqrt{25}$ ,  $2^3$

If they were negative, the answer would be:  
 $\sqrt{25}$ ,  $\sqrt[3]{27}$ ,  $3^{-2}$ ,  $2^3$
- $\frac{9^5}{9^3 \times 9^2} = \frac{9^5}{9^5} = 1$

### Standard form

- 2750
- $1.5 \times 10^8$

- 3 Move the decimal point three places to the right to give  $6.42 \times 10^{-3}$
- 4  $(1.4 \times 10^{-5}) \times 20 = (2.8 \times 10^{-5}) \times 2 \times 10 = 2.8 \times 10^{-4}$  km

### Listing strategies

- 1 259, 295, 529, 592, 925, 952

2 a

		4-sided spinner			
		0	1	2	3
3-sided spinner	1	1	2	3	4
	2	2	3	4	5
	3	3	4	5	6

b 3

3

		Dice					
		1	2	3	4	5	6
Coin	H	H1	H2	H3	H4	H5	H6
	T	T1	T2	T3	T4	T5	T6

- 4 spj; spi; sfj; sfi ; bpj; bpi; bfj; bfi

## Algebra

### Understanding expressions, equations, formulae and identities

- 1 a identity    b equation    c expression
- 2 a Equation, because it has an equals sign and can be solved.  
b Formula, because it has an equals sign and cannot be solved.  
c Expression, because it has letter terms and no equals sign.  
d Formula, because it has an equals sign and cannot be solved.
- 3 a Any of:  $2x + 10$  or  $10x + 2$  or  $x + 210$  or  $x + 102$   
b Any of:  $2x = 10$  or  $10x = 2$

### Simplifying expressions

- 1  $8x$
- 2 a  $6a \times 8a = (6 \times 8) \times (a \times a) = 48 \times a^2 = 48a^2$   
b  $2p \times 3p \times 5p = (2 \times 3 \times 5) \times (p \times p \times p) = 30 \times p^3 = 30p^3$
- 3  $35yz \div 7z = (35 \div 7) \times (yz \div z) = 5 \times y = 5y$
- 4  $\frac{32uv}{4v} = \frac{32}{4} \times \frac{uv}{v} = 8 \times u = 8u$

### Collecting like terms

- 1 a  $7m + 6n - 4m - 2n = (7 - 4)m + (6 - 2)n = 3m + 4n$   
b  $9q - 5r - 12q + 3r = (9 - 12)q + (3 - 5)r = -3q - 2r$
- 2 a  $11a + 5b - 10a + 8b = (11 - 10)a + (5 + 8)b = a + 13b$   
b  $6c - 4d - 7c + 5d = (6 - 7)c + (5 - 4)d = -c + d$
- 3 a  $9p^3 + p - 4p^3 = (9 - 4)p^3 + p = 5p^3 + p$   
b  $12 - 5x^2 + 3x - 2x^2 = 12 - (5 + 2)x^2 + 3x = -7x^2 + 3x + 12$
- 4  $3\sqrt{5} - f - 8\sqrt{5} + 2f = (3 - 8)\sqrt{5} + (2 - 1)f = -5\sqrt{5} + f$

### Using indices

- 1 a  $p^3 \times p = p^{(3+1)} = p^4$   
b  $4y^2 \times 3y^3 = (4 \times 3) \times y^{(2+3)} = 12 \times y^5 = 12y^5$   
c  $2a^4b \times 5ab^2 = (2 \times 5) \times a^{(4+1)} \times b^{(1+2)} = 10 \times a^5 \times b^3 = 10a^5b^3$
- 2 a  $q^{-2} \times q^{-4} = q^{(-2-4)} = q^{-6}$   
b  $(u^{-3})^2 = u^{(-3 \times 2)} = u^{-6}$   
c  $x^{-1} \times x = x^{-1} \times x^1 = x^{(-1+1)} = x^0 = 1$
- 3 a  $b^4 \div b^3 = b^{(4-3)} = b^1 = b$   
b  $\frac{f^5}{f^2} = f^{(5-2)} = f^3$   
c  $\frac{xy^3}{x^2y} = x^{(1-2)} \times y^{(3-1)} = x^{-1} \times y^2 = \frac{1}{x} \times y^2 = \frac{y^2}{x}$

- 4 Let the first box =  $x$  and the second box =  $y$   
 $(xm^3)^y = x^y m^{3y} = 8m^9$   
comparing terms,  $3y = 9$   
 $y = 3$   
Substitute in the  $y$  value:  $(xm^3)^3 = 8m^9$   
 $x^3 = 8$   
 $x = \sqrt[3]{8} = 2$   
Therefore, the completed expression is  $(2m^3)^3$

### Expanding brackets

- 1 a  $4(m + 3) = (4 \times m) + (4 \times 3) = 4m + 12$   
b  $2(p - 1) = (2 \times p) + (2 \times -1) = 2p - 2$   
c  $10(3x - 5) = (10 \times 3)x + (10 \times -5) = 30x - 50$
- 2 a  $3(m + 2) + 5(m + 1) = 3m + 6 + 5m + 5 = 8m + 11$   
b  $6(x - 1) - 2(x - 4) = 6x - 6 - 2x + 8 = 4x + 2$
- 3 a  $(y + 3)(y + 7) = y^2 + 7y + 3y + 21 = y^2 + 10y + 21$   
b  $(b + 2)(b - 4) = b^2 - 4b + 2b - 8 = b^2 - 2b - 8$   
c  $(x - 4)(x - 6) = x^2 - 6x - 4x + 24 = x^2 - 10x + 24$
- 4 a  $(q + 1)^2 = (q + 1)(q + 1) = q^2 + q + q + 1 = q^2 + 2q + 1$   
b  $(z + 2)^2 = (z + 2)(z + 2) = z^2 + 2z + 2z + 4 = z^2 + 4z + 4$   
c  $(c - 3)^2 = (c - 3)(c - 3) = c^2 - 3c - 3c + 9 = c^2 - 6c + 9$

### Factorising

- 1 Divide the expression by the highest common factor (HCF) of both terms to find the bracket, and then multiply the bracket by the HCF to give the full factorisation.  
a  $(4x + 8) \div 4 = x + 2$   
factorisation:  $4(x + 2)$   
b  $(3d - 15) \div 3 = d - 5$   
factorisation:  $3(d - 5)$   
c  $(8y - 12) \div 4 = 2y - 3$   
factorisation:  $4(2y - 3)$
- 2 Divide the expression by the common term to find the bracket, and then multiply the bracket by the common term to give the full factorisation.  
a  $(q^2 + q) \div q = q + 1$   
factorisation:  $q(q + 1)$   
b  $(a^2 + 6a) \div a = a + 6$   
factorisation:  $a(a + 6)$   
c  $(10z^2 + 15z) \div 5z = (2z + 3)$   
factorisation:  $5z(2z + 3)$
- 3 Find which factors of the number term add together to give the coefficient of the  $x$  term.  
a  $12 = 3 \times 4$   
 $7 = 3 + 4$   
factorisation:  $(x + 3)(x + 4)$   
b  $-16 = (-2) \times 8$   
 $6 = -2 + 8$   
factorisation:  $(x - 2)(x + 8)$   
c  $24 = (-6) \times (-4)$   
 $-10 = (-6) + (-4)$   
factorisation:  $(a - 6)(a - 4)$
- 4 a Write  $y^2 - 4$  in the form of  $a^2 - b^2$ :  
 $y^2 - 2^2$   
Using the formula for the difference of two squares, the factorisation is  
 $(y + 2)(y - 2)$   
b Write  $x^2 - 9$  in the form of  $a^2 - b^2$ :  
 $x^2 - 3^2$   
Using the formula for the difference of two squares, the factorisation is  
 $(x + 3)(x - 3)$

c Write  $p^2 - 100$  in the form of  $a^2 - b^2$ :

$$p^2 - 10^2$$

Using the formula for the difference of two squares, the factorisation is

$$(p + 10)(p - 10)$$

### Substituting into expressions

1  $4x + 5y = 4 \times 3 + 5 \times (-2) = 12 - 10 = 2$

2  $s = ut + \frac{1}{2}at^2$   
 $= 12 \times 2 + \frac{1}{2} \times 10 \times 2^2$   
 $= 12 \times 2 + \frac{1}{2} \times 40$   
 $= 24 + 20$

$$s = 44$$

3 a  $f = 3c - 2(c - d)$   
 $= 3 \times 7 - 2 \times (7 - (-5))$   
 $= 21 - 2 \times (12)$   
 $= 21 - 24$

$$f = -3$$

b  $f = -c(d^2 - 3c)$   
 $= -7 \times ((-5)^2 - 3 \times 7)$   
 $= -7 \times (25 - 21)$   
 $= -7 \times 4$

$$f = -28$$

c  $f^2 = 7c - 3d$   
 $= 7 \times 7 - 3 \times (-5)$   
 $= 49 + 15$   
 $= 64$

$$f = \sqrt{64}$$

$$f = \pm 8$$

### Writing expressions

1 a  $n + 3$       b  $(n \times 2) - 9 = 2n - 9$

2 a  $x + y$       b  $5 \times x = 5x$

c  $(12 \times x) + (11 \times y) = 12x + 11y$

3  $2 \times 9p + 2(5p + 2) = 18p + 10p + 4 = 28p + 4$

4 The area of the rectangle is given by height  $\times$  length, which is  $s \times (5s + 1) = s(5s + 1)$ .

### Solving linear equations

1 a  $x = 12 - 5$

$$x = 7$$

b  $x = 10 + 3$

$$x = 13$$

c  $x = \frac{20}{4}$

$$x = 5$$

d  $x = 6 \times 3$

$$x = 18$$

2 a  $2x + 3 = 15$

$$2x = 12$$

$$x = 6$$

b  $3x - 5 = 16$

$$3x = 21$$

$$x = 7$$

c  $\frac{x}{5} + 3 = 8$

$$\frac{x}{5} = 5$$

$$x = 25$$

d  $7 - 2x = 1$

$$7 = 2x + 1$$

$$6 = 2x$$

$$x = 3$$

3 a  $3(x + 9) = 30$

$$3x + 27 = 30$$

$$3x = 3$$

$$x = 1$$

b  $5(p - 2) = 10$

$$5p - 10 = 10$$

$$5p = 20$$

$$p = 4$$

c  $2(10 - 3m) = 8$

$$20 - 6m = 8$$

$$12 = 6m$$

$$m = 2$$

d  $4(8 - 2q) = 0$

$$32 - 8q = 0$$

$$32 = 8q$$

$$q = 4$$

4 a  $4x - 6 = x + 9$

$$3x - 6 = 9$$

$$3x = 15$$

$$x = 5$$

b  $2y + 5 = 4y - 3$

$$5 = 2y - 3$$

$$8 = 2y$$

$$y = 4$$

c  $4(2x + 3) = 11x + 3$

$$8x + 12 = 11x + 3$$

$$9 = 3x$$

$$x = 3$$

d  $3(n + 4) = 2(2n + 3)$

$$3n + 12 = 4n + 6$$

$$12 = n + 6$$

$$n = 6$$

### Writing linear equations

1 Sum of the angles in a triangle are  $180^\circ$

$$(2x + 3) + 81 + (3x - 4) = 180$$

$$5x + 80 = 180$$

$$5x = 100$$

$$x = 20^\circ$$

2 Let Jamie's age =  $x$  years. Sophie's age =  $\frac{x}{2}$

$$x + \frac{x}{2} = 18$$

$$\frac{3x}{2} = 18$$

$$3x = 36$$

$$x = 12$$

Jamie is 12 years old

3 Let width =  $x$  so length =  $x + 3$

$$\text{Perimeter} = 2x + 2(x + 3)$$

$$46 = 4x + 6$$

$$x = 10$$

Length = 10cm and width = 13cm

$$\text{Area} = 10 \times 13 = 130\text{cm}^2$$

4 Opposite angles are equal so  $3x + 10 = 5x - 20$

$$30 = 2x \text{ giving } x = 15^\circ$$

$$\text{Also } 3x + 10 + 7x + 5y = 180$$


$$10x + 10 + 5y = 180$$

$$\text{Now } x = 15 \text{ so } 150 + 10 + 5y = 180$$

$$\text{Solving this gives } y = 4^\circ$$


### Linear inequalities

1 a  $-1, 0, 1, 2, 3, 4, 5$



2 a  $4x > 20$


$x > 5$



b  $3x - 8 \leq 13$

$3x \leq 21$

$x \leq 7$




c  $2(x - 3) < 10$

$2x - 6 < 10$

$2x < 16$

$x < 8$



3 a  $2 \leq 3x + 5$

$-3 \leq 3x$

$-1 \leq x$

$3x + 5 < 11$

$3x < 6$

$x < 2$

Hence  $-1 \leq x < 2$

b  $-4 > 5x + 6$

$-10 < 5x$

$-2 < x$

$5x + 6 \leq 6$

$5x \leq 0$

$x \leq 0$

Hence  $-2 < x \leq 0$

4  $n + n + 3 < 15$

$2n + 3 < 15$

$2n < 12$

$n < 6$

$n = 1, 2, 3, 4, 5$

**Formulae**

1 a  $t = (40 \times 2) + 20 = 100$  minutes = 1 hour 40 minutes

b  $t = (40 \times 1.5) + 20 = 80$  minutes = 1 hour 20 minutes

The chicken should be put in the oven 1 hour and 20 minutes earlier than 1.30 pm, which is a time of 12.10 pm.

2 a  $C = l + kn$

b  $C = 90 + 6.5 \times 3$

$C = \text{£}109.50$

3 a  $p = \frac{qs}{3}$

$3p = qs$

$q = \frac{3p}{s}$

b  $p = \frac{q}{r} + t$

$p - t = \frac{q}{r}$

$q = rp - rt = r(p - t)$

c  $p = 3(q + r)$

$\frac{p}{3} = q + r$

$q = \frac{p}{3} - r = \frac{p - 3r}{3}$

d  $p = \sqrt{2q}$

$p^2 = 2q$

$q = \frac{p^2}{2}$

**Linear sequences**

1 a The term in position 1 is  $1 \times 5 + 1 = 6$

The term in position 2 is  $2 \times 5 + 1 = 11$

The term in position 3 is  $3 \times 5 + 1 = 16$

The term in position 4 is  $4 \times 5 + 1 = 21$

b The term in position 50 is  $50 \times 5 + 1 = 251$

2 a Each pattern has 2 more dots than the last, so pattern 7 will have 8 more dots than pattern 3. Pattern 7 will have 19 dots.

b Rachel is wrong. Each pattern adds one triangle, so in pattern 4 there will be one more triangle than in pattern 3, giving 6 triangles. The rule for the number of triangles is  $n + 2$ .

3 a Common difference = 11, so  $11n$  is in the sequence.

When  $n = 1$ :

$11n = 11$ , but the 1<sup>st</sup> term is 3.

$3 = 11n - 8$

So the expression for the sequence is  $11n - 8$

b Assume 100 is in the sequence. Then:

$11n - 8 = 100$

$11n = 108$

$n = 108 \div 11 = 9$  remainder 9

But  $n$  must be a whole number, and it is not; so 100 is not in this sequence.

**Non-linear sequences**

1 a Rule is multiply by 2.

$8 \times 2 = 16$

$16 \times 2 = 32$

So terms are 16, 32

b Rule is divide by 10.

$1 \div 10 = 0.1$

$0.1 \div 10 = 0.01$

So terms are 0.1, 0.01

c Rule is multiply by  $-2$ .

$-12 \times -2 = 24$

$24 \times -2 = -48$

So terms are 24,  $-48$

d They involve multiplying and dividing, not adding and subtracting, so they are geometric.

2 a Next term =  $6 + 9 = 15$

b 5<sup>th</sup> term =  $6 + 9 = 15$

6<sup>th</sup> term =  $9 + 15 = 24$

7<sup>th</sup> term =  $15 + 24 = 39$

8<sup>th</sup> term =  $24 + 39 = 63$

9<sup>th</sup> term =  $39 + 63 = 102$

The 9<sup>th</sup> term is the first term in the sequence over 100

3 a

Day	Mon	Tue	Wed	Thu	Fri
Number of ladybirds	2	8 (= 2 × 4)	32 (= 8 × 4)	32 × 4 = 128	128 × 4 = 512

The gardener is correct. There will be more than 500 ladybirds.

b Saturday:  $512 \times 4 = 2048$

4 a First term:  $\frac{1}{2} \times 1^2 = \frac{1}{2}$

Second term:  $\frac{1}{2} \times 2^2 = 2$

Third term:  $\frac{1}{2} \times 3^2 = \frac{9}{2} = 4\frac{1}{2}$

b If 32 is in the sequence, then:

$\frac{1}{2}n^2 = 32$

$n^2 = 64$

$n = 8$

This gives  $n$  as a whole number, 8, so 32 is the 8<sup>th</sup> term in the sequence.

**Show that...**

- LHS =  $2x + 1$ ; RHS =  $2x + 1$ ; LHS = RHS. Therefore,  $2(x + \frac{1}{2}) \equiv x + x + 1$
- LHS =  $x^2 - 25 + 9 = x^2 - 16$ ; RHS =  $x^2 - 16$
- Let the three consecutive numbers be  $n, n + 1$  and  $n + 2$ .  
 $n + n + 1 + n + 2 = 3n + 3 = 3(n+1)$ . Therefore, the sum of three consecutive numbers is a multiple of 3.
- a** Width of pond =  $x - y + x + x + x - y = 4x - 2y$   
Length of pond =  $4x$   
Perimeter =  $4x - 2y + 4x - 2y + 4x + 4x = 16x - 4y$   
**b** Yes Sanjit is correct, because  $16x - 4y = 4(4x - y)$ , showing that when  $x$  and  $y$  are whole numbers, the perimeter is always a multiple of 4.

**Functions**


- a** when  $x = 3, y = 3 \times 4 - 1 = 11$   
**b** when  $y = 23, 4x - 1 = 23$ , therefore  $x = (23 + 1) \div 4 = 6$   
**c** To get  $y$  you multiply  $x$  by 4 and subtract 1, so  $y = 4x - 1$

$x$	Operations	$y$
-2	$(-2) \times 2 + 3$	-1
0	$0 \times 2 + 3$	3
3	$(9 - 3) \div 2$	9

$x$	Operations	$y$
-2	$(-2) \div 2 + 1$	0
1	$(1) \div 2 + 1$	$1\frac{1}{2}$
8	$(5 - 1) \times 2$	5


**Coordinates and midpoints**

- a** 1 along the  $x$ -axis and 2 up the  $y$ -axis: (1, 2)



- c**  $4\frac{1}{2}$  along the  $x$ -axis and 1 up the  $y$ -axis:  $(4\frac{1}{2}, 1)$

- a** 1 along the  $x$ -axis and -4 'up' the  $y$ -axis: (1, -4)



S = (6, -2) to make a parallelogram

- a**  $x$  coordinate =  $\frac{4 + (-2)}{2} = 1$   
 $y$  coordinate =  $\frac{5 + 1}{2} = 3$   
Midpoint of  $XY$  is (1, 3).


**b** Midpoint of  $XZ = (\frac{4+4}{2}, \frac{5+(-4)}{2}) = (4, \frac{1}{2})$

**c** Midpoint of  $YZ = (\frac{(-2)+4}{2}, \frac{1+(-4)}{2}) = (1, -1\frac{1}{2})$

**Straight line graphs**


- a** For  $y = 2x + 3$

$x$	-1	0	1	2
Operations	$2 \times (-1) + 3$	$2 \times (0) + 3$	$2 \times (1) + 3$	$2 \times (2) + 3$
$y$	1	3	5	7



- a** For  $y = 3x$

$x$	-2	-1	0	1	2
Operations	$3 \times (-2)$	$3 \times (-1)$	$3 \times (0)$	$3 \times (1)$	$3 \times (2)$
$y$	-6	-3	0	3	6



- b** Rearrange the equation to give  $y = -2x - \frac{1}{2}$

$x$	Operations	$y$
-3	$-2 \times (-3) - \frac{1}{2}$	$5\frac{1}{2}$
-2	$-2 \times (-2) - \frac{1}{2}$	$3\frac{1}{2}$
-1	$-2 \times (-1) - \frac{1}{2}$	$1\frac{1}{2}$
0	$-2 \times (0) - \frac{1}{2}$	$-\frac{1}{2}$
1	$-2 \times (1) - \frac{1}{2}$	$-2\frac{1}{2}$
2	$-2 \times (2) - \frac{1}{2}$	$-4\frac{1}{2}$

- c** Line C goes through points (0, -3), (1, 1) and (2, 5)  
The  $y$  intercept is -3.

The gradient is  $\frac{\text{difference in } y \text{ coordinates}}{\text{difference in } x \text{ coordinates}} = \frac{5 - 1}{2 - 1} = 4$

The equation of line C is  $y = 4x - 3$ .

- a** B and C, because they have the same gradient of 2.  
**b** A and B, because they both have a  $y$ -intercept at (0, 1).
- The gradient is  $\frac{2 - (-6)}{3 - (-1)} = \frac{8}{4} = 2$

Using point (3, 2) and gradient  $m = 2$ :


$$y = 2x + c$$

$$2 = 2 \times 3 + c$$

$$c = -4$$


Equation of the line is  $y = 2x - 4$

5 a




Draw a vertical line up from  $x = 3$  to the graph, and then a horizontal line to the  $y$ -axis to read off the result:  $y = 2$ .

b




Draw a horizontal line across from  $y = -2$  to the graph, and then a vertical line up to the  $x$ -axis to read off the result:  $x = 1$ .

6 a



Draw a horizontal line across from  $y = -2$  to the graph, and then a vertical line up to the  $x$ -axis to read off the result:  $x = 2.5$ .


b



Draw a vertical line up from  $x = -0.5$  to the graph, and then a vertical line across to the  $y$ -axis to read off the result:  $y = 4$ .


c Reading up from  $x = -1.2$  and then across to  $y$  axis gives  $y = 0.6$ . Any value from 0.6 to 0.75 is acceptable.

7 a, b



b Where the graphs cross, draw a vertical line up to the  $x$ -axis. It meets the axis two thirds of the way between  $x = -1$  and  $x = 0$ , so the solution is approximately  $x = -0.67$ . Any value from  $-0.6$  to  $-0.7$  is acceptable.

8



Compare the equations  $y = 4x - 3$  and  $4x - 3 = 2$ .  $y$  has been replaced with 2, so add line  $y = 2$  to the graph.

The intersection point of the two graphs gives the solution to the equation  $4x - 3 = 2$

$x = 1.25$ . Any answer between 1.2 and 1.3 is acceptable.

## Solving simultaneous equations

- 1 a Substituting  $y = 2x$  into the first equation gives  
 $3x + 2x = 15$   
 $5x = 15$   
 $x = 3$

When  $x = 3$ ,  $y = 2(3) = 6$

- b  $3x = 12$   
 $x = 4$

Substituting  $x = 4$  into the first equation gives  
 $8 + y = 9$   
 $y = 1$

- c  $5x = 10$   
 $x = 2$

Substituting  $x = 2$  into the first equation gives  
 $6 + y = 4$   
 $y = -2$

- 2 a  $2x + 2y = 14$  (1)  
 $3x + y = 11$  (2)  
 $(2) \times 2 \quad 6x + 2y = 22$   
 $(3) - (1) \quad 4x = 8$   
 $x = 2$

Substitute into (1)  $4 + 2y = 14$   
 $2y = 10$   
 $y = 5$

Solution is  $x = 2, y = 5$

- b  $4x - 2y = 2$  (1)  
 $2x - 3y = 7$  (2)  
 $(2) \times 2 \quad 4x - 6y = 14$  (3)  
 $(3) - (1) \quad -4y = 12$   
 $y = -3$

Substitute into (1)  $4x + 6 = 2$   
 $4x = -4$   
 $x = -1$

Solution is  $x = -1, y = -3$

- c  $2x + 3y = 20$  (1)  
 $3x + 2y = 15$  (2)  
 $(1) \times 2 \quad 4x + 6y = 40$  (3)  
 $(2) \times 3 \quad 9x + 6y = 45$  (4)  
 $(4) - (3) \quad 5x = 5$   
 $x = 1$


Substituting  $x = 1$  into equation (1)  $2 + 3y = 20$   
 $3y = 18$   
 $y = 6$

Solution is  $x = 1, y = 6$


- 3 a  $x + y = 2$        $2x - y = 1$

x	0	2
y	2	0

x	0	$\frac{1}{2}$
y	-1	0



From the intersection of the two lines,  $x = 1, y = 1$ .




From the intersection of the two lines,  $x = 2, y = 3$ .

## Quadratic graphs


- 1 a C and D – they are straight lines so they are linear.  
 b E – it is quadratic, with a positive multiplier for  $x^2$ , and is symmetrical about the origin.  
 c A – it is quadratic, with a negative multiplier for  $x^2$ , and is symmetrical about the origin.  
 d D – the  $x$  coordinates are all different, but all the  $y$  coordinates on this line are 1.  
 e B – it is the same as E except that it has been moved 1 unit up the  $y$ -axis.

2 a


x	-3	-2	-1	0	1	2	3
y	6	1	-2	-3	-2	1	6



- c  $x = 0$       d  $(0, -3)$



- b  $x = -1$       c  $(-1, -2)$




- b  $x = -2$  or  $x = 1$

- 5  $x = -3$  or  $x = 1$

6 a

x	-3	-2	-1	0
$x^2 + 3x + 1$	$(-3)^2 + 3 \times (-3) + 1$	$(-2)^2 + 3 \times (-2) + 1$	$(-1)^2 + 3 \times (-1) + 1$	$(0)^2 + 3 \times (0) + 1$
y	1	-1	-1	1





- b** Compare the equations  $y = x^2 + 3x + 1$  and  $x^2 + 3x + 1 = -1$ .  $y$  has been replaced with  $-1$ , so the solutions to the equation  $x^2 + 3x + 1 = -1$  are where  $y = -1$ .  
 $x = -2$  and  $x = -1$
- c** Compare the equations  $y = x^2 + 3x + 1$  and  $x^2 + 3x + 1 = 0$ .  $y$  has been replaced with  $0$ , so the solutions to the equation  $x^2 + 3x + 1 = 0$  are where the graph crosses the  $x$ -axis.  
 $x = -2.6$  and  $x = -0.38$  (any answer close to  $-0.4$  is acceptable)

**Solving quadratic equations**


- 1 **a**  $x(x + 6) = 0$   
 $x = 0$  or  $-6$
- b**  $y(y - 11) = 0$   
 $y = 0$  or  $11$
- c**  $3d(d - 3) = 0$   
 $d = 0$  or  $3$
- 2 **a**  $(x + 4)(x - 4) = 0$   
 $x = 4$  or  $-4$
- b**  $(a + 9)(a - 9) = 0$   
 $a = 9$  or  $-9$
- c**  $(z - 10)(z + 10) = 0$   
 $z = 10$  or  $-10$
- 3 a**  $(x + 3)(x + 2) = 0$   
 $x = -2$  or  $-3$
- b**  $(x + 5)(x - 2) = 0$   
 $x = -5$  or  $2$
- c**  $(x - 7)(x - 2) = 0$   
 $x = 2$  or  $7$
- 4 a**  $0 = x(x - 3)$   
 $x = 0$  or  $3$
- b**  $0 = (x - 5)(x + 5)$   
 $x = 5$  or  $-5$
- c**  $0 = (x + 6)(x - 3)$   
 $x = -6$  or  $3$

**Cubic and reciprocal graphs**

- 1 **a** A, C and D – they are not continuous curves with two turning points (s-shaped curves).  
**b** B – it has two turning points and has rotational symmetry about the origin.  
**c** E – it has two turning points and is a reflection of B, raised up one unit on the  $y$ -axis.  
**d** D – this is the form for a reciprocal graph.

2 **a**


$x$	-2	-1	0	1	2
$y$	-4	3	4	5	12



- 3 **a** cubic      **b** (0, -8)      **c** (2, 0)


4 **a**

$x$	-3	-2	-1	$-\frac{1}{2}$	$\frac{1}{2}$	1	2	3
$y$	$\frac{2}{3}$	$\frac{1}{2}$	0	-1	3	2	$1\frac{1}{2}$	$1\frac{1}{3}$




**Drawing and interpreting real-life graphs**

- 1 **a** The initial charge is the value at 0 miles, where the graph cuts the  $y$ -axis (the  $y$ -intercept): \$3.  
**b** The charge per mile is given by the gradient of the graph.  
 Gradient =  $\frac{\text{difference in } y \text{ coordinates}}{\text{difference in } x \text{ coordinates}} = \frac{23 - 3}{8 - 0} = \frac{20}{8} = 2.5$   
 Charge per mile is \$2.50



- d** From the conversion graph, £15 = \$18.  
 From the graph of the cost of a taxi in New York, \$18 allows you to travel 6 miles.
- 2 **a** You can read this from the highest point of the graph: 16 m.  
**b** Read from the highest point down to the value on the horizontal axis: 4 seconds.  
**c** This is when the ball reaches 0 height for the second time: 8 seconds.  
**d** Read from 12 on the vertical axis across to the curve: 2 seconds and 6 seconds.
- 3 **a** This is the highest value on the vertical axis: 10 m/s.  
**b** The cyclist is travelling at a constant speed of 10 m/s.  
**c** The cyclist is decelerating, so it will be a negative value.  
 Acceleration =  $\frac{-10}{20} = -\frac{1}{2} \text{ m/s}^2$   
**d** The speed returns to 0: the cyclist stops.
- 4 **a** Assuming that water pours into the container at a constant rate,  
 A: depth goes up increasingly slowly as the container widens, so 2.  
 B: depth rises steadily and fairly slowly in a broad container of consistent diameter, so 4.  
 C: depth rises quickly at first, then more slowly as the container widens, then more quickly as it gets narrow towards the top, so 3.  
 D: depth rises steadily and quickly in a narrow container of consistent diameter, so 1.



## Ratio, proportion and rates of change

### Units of measure

- $4(\text{m}) \times 100 = 400 \text{ cm}$
  - $500(\text{g}) \div 1000 = 5 \text{ kg}$
  - $1.5(\text{l}) \times 1000 = 1500 \text{ ml}$
  - $8250(\text{m}) \div 1000 = 8.25 \text{ km}$
- $6(\text{litres}) \times 1000 = 6000 \text{ ml}$   
 $6000 - 3500 = 2500$   
Sally had **2500 ml** of lemonade left.
- Luke: 240 seconds; Adam:  $3 \times 60 + 47 = 227$  seconds.  
Adam arrived first.  
or Luke:  $240 \div 60 = 4$  minutes; Adam: 3 minutes 47 seconds. Adam arrived first.
  - $4 \text{ minutes} - 3 \text{ minutes } 47 \text{ seconds} = 13 \text{ seconds}$ .  
Adam waited **13 seconds** for Luke to arrive at school.
- Ben =  $1.25 \text{ m} = 3.2 + 0.8 \text{ feet} = 4 \text{ feet}$ . Tom is taller.  
or Tom =  $4.8 \text{ feet} = 3.2 + 1.6 \text{ feet} = 1 + 0.5 \text{ metres} = 1.5 \text{ metres}$ . Tom is taller.

### Ratio

- 12 stories and 8 colouring  
Ratio of story: colouring =  $12 : 8 = 3 : 2$
- Total parts =  $1 + 2 = 3$   
 $1 \text{ part} = \frac{15}{3} = 5$   
Spent =  $2 \times 5 = \text{£}10$
- Total parts =  $1 + 2 + 3 = 6$   
 $1 \text{ part} = \frac{60}{6} = 10$   
Amount given to charity =  $3 \times 10 = \text{£}30$
- Ratio of blue to yellow required is  $3 : 7$ .  
There are  $3 + 7 = 10$  parts. He needs to make 5 litres.  
 $10 \text{ parts} = 5000 \text{ ml}$   
 $1 \text{ part} = 500 \text{ ml}$   
Phil needs  $3 \times 500 \text{ ml} = 1500 \text{ ml} = 1.5 \text{ litres}$  of blue paint.  
He has 2 litres of blue paint.  
Phil needs  $7 \times 500 \text{ ml} = 3500 \text{ ml} = 3.5 \text{ litres}$  of yellow paint.  
He has 3 litres of yellow paint.  
Phil has enough blue paint, but does not have enough yellow paint.

### Scale diagrams and maps

- 1 cm on the map is **10000 cm** in real life.  
This means 1 cm on the map is **100 m** in real life.
- 1 cm on the map represents 50 m in real life.  
 $3 \times 50 \text{ m} = 150 \text{ m}$ , so the bus stop is **3 cm** from the village shop on the map.
- Measure the distance between the trees on the diagram = 5 cm  
1 cm on the diagram represents 4 m in real life.  
 $5 \times 4 = 20$   
The trees are **20 m** apart.
- A scale of  $1 : 400$  means 1 cm on the model represents 4 m (= 400 cm) in real life.  
 $96 \div 4 = 24$   
The scale model is **24 cm** tall.

### Fractions, percentages and proportion

- $1 + 3 = 4$  parts so Bess receives  $\frac{3}{4}$
- $1 : 3 : 6$
  - $1 + 3 + 6 = 10$  items in the basket  
Fruit =  $\frac{3}{10}$
  - Tins =  $\frac{6}{10} = 60\%$
- $\frac{50}{4000} = \frac{1}{80}$

- Total parts =  $3 + 8 + 14 = 25$   
 $\frac{8}{25} = \frac{32}{100} = 32\%$

### Direct proportion

- One ticket costs  $\text{£}80 \div 5 = \text{£}16$
  - Nine tickets cost  $9 \times \text{£}16 = \text{£}144$
- Read up from 6 packs on the horizontal axis, to the line, then across to the vertical axis to find the cost:  $\text{£}1.20$
  - There are 10 pencils in a pack, so 1 pencil is 0.1 of a pack. Reading off the graph using this value, the price is 2p.
  - It is a straight-line graph; the graph passes through the origin (0, 0).
- Sally needs to make  $28 \div 4 = 7$  lots of the recipe.  
She will need  $1 \times 7 = 7$  teaspoons of turmeric,  
 $2 \times 7 = 14$  teaspoons of chilli powder and  
 $2\frac{1}{2} \times 7 = 17\frac{1}{2}$  teaspoons of cumin.
  - Sally has 75 g of chilli powder. That is  $75 \div 3 = 25$  teaspoons.  
Sally needs 14 teaspoons to make the curry for her class.  
She does have enough.

### Inverse proportion

- Start from 5 on the  $x$ -axis, read up to the graph, then left to the scale on the  $y$ -axis.  
5 winners will each get **£400**.
  - Start from 200 on the  $y$ -axis, read right to the graph, then down to the scale on the  $x$ -axis. 10 winners each get  $\text{£}200$ , so there are **9** other winners.
  - Multiply the number of winners by the amount each one gets to find the total prize money. e.g.  $5 \times \text{£}400 = \text{£}2000$ .
  - Compare 2 points on the graph. (Use your answers to parts **a** and **b**.) If the number of winners goes up, the prize money they each receive goes down. The prize money and the number of winners are in inverse proportion.
- The total time needed to decorate the room is  $3 \times 2 = 6$  hours.
  - $6 \text{ hours} \div 12 \text{ people} = 0.5 \text{ hours} = \text{30 minutes}$ .
- The printer can print  $240 \div 4 = 60$  pages per minute.  
 $600 \div 60 = 10$ , so it would take **10 minutes** to print the larger document.

### Working with percentages

- $125 - 75 = 50$   
 $\frac{50}{125} \times 100 = 40\%$
- Amount of increase =  $24 - 15 = \text{£}9$  million  
 $\frac{9}{15} \times 100 = \text{60\%}$  increase in sales for Company X
  - $125\% = \text{£}35$  million  
 $35 \div 125 \times 100 = \text{£}28$  million sales in 2006.
- $\frac{2}{100} \times 265 = 5.30$  and  $3 \times 5.30 = 15.90$   
 $15.90 + 265 = \text{£}280.90$
  - $265 \times (1.02)^3 = \text{£}281.22$


### Compound units

- $\frac{3 \text{ km}}{\text{minute}} = \frac{3000 \text{ m}}{\text{minute}} = \frac{3000 \text{ m}}{60 \text{ seconds}} = 50 \text{ m/s}$
- $20 \div 5 = 4$  minutes to fill the tank.
- Pressure =  $\frac{300}{0.05} = 6000$  Newtons/m<sup>2</sup>
- On Saturday Sami drove  $4 \times 50 = 200$  miles; on Sunday Sami drove  $356 \div 8 \times 5 = 222.5$  miles. Sami drove further on Sunday.

## Geometry and measures

### Measuring and drawing angles

- 1 a  $123^\circ$   
b  $42^\circ$   
c  $331^\circ$



- 3 a  $100^\circ, 120^\circ, 140^\circ, 160^\circ$   
b First angle + second angle =  $87^\circ$ . This means both angles are less than  $87^\circ$ , and so they both must be acute.

### Using the properties of angles

- 1 a Angle  $ACB = 52^\circ$  (Angles in a triangle add up to  $180^\circ$ )  
 $x = 128^\circ$  (Angles on a straight line add up to  $180^\circ$ )  
b Angle  $ADC = 86^\circ$  (Angles in a quadrilateral add up to  $360^\circ$ )  
 $x = 94^\circ$  (Angles on a straight line add up to  $180^\circ$ )
- 2 a Angle  $BED = 39^\circ$  (Alternate angles are equal)  
Angle  $BDE = 39^\circ$  (Base angles in an isosceles triangle are equal)  
 $x = 102^\circ$  (Angles in a triangle add up to  $180^\circ$ )  
b Angle  $DCF = 98^\circ$  (Vertically opposite angles are equal)  
 $x = 98^\circ$  (Corresponding angles are equal)
- 3 Angle  $CFG = 62^\circ$  (Co-interior angles add up to  $180^\circ$ )  
 $x = 66^\circ$  (Angles on a straight line add up to  $180^\circ$ )
- 4  $x + 40 + 3x + 5x - 40 = 180^\circ$   
 $9x = 180^\circ$   
 $x = 20^\circ$   
Angle  $BAC = x + 40 = 20 + 40 = 60^\circ$   
Angle  $ACB = 3x = 3 \times 20 = 60^\circ$   
Angle  $ABC = 5x - 40 = 5 \times 20 - 40 = 60^\circ$   
Triangle  $ABC$  has equal angles of  $60^\circ$ . Therefore, it is an equilateral triangle.


### Using the properties of polygons

- 1 a  $180^\circ \times (6 - 2) = 720^\circ$   
b  $720^\circ \div 6 = 120^\circ$   
c  $180^\circ - 120^\circ = 60^\circ$  or  $360^\circ \div 6 = 60^\circ$
- 2 a It is an octagon because it has eight sides.  
b All angles are equal; all sides are equal.  
c  $180^\circ \times (8 - 2) = 1080^\circ$   
 $1080^\circ \div 8 = 135^\circ$   
or  $180^\circ - (360^\circ \div 8) = 135^\circ$
- 3 exterior angle =  $180^\circ - 144^\circ = 36^\circ$   
number of sides =  $360 \div 36 = 10$   
Therefore it is a decagon.

### Using bearings


- 1 a This is the angle measured clockwise from North at  $A$ :  $065^\circ$ .  
b  $180 - 138 = 42^\circ$  This is the acute angle at  $C$ .  
Bearing of  $B$  from  $C = 360 - 42^\circ = 318^\circ$   
c  $180 - 65 = 115^\circ$  This is the angle between the north line at  $B$  and  $AB$ , measured anticlockwise.  
Bearing of  $A$  from  $B = 360 - 115 = 245^\circ$
- 2 To find a reciprocal bearing, subtract 180 from the original bearing (or add 180 to it).  
The bearing of  $O$  from  $X = 276 - 180 = 096^\circ$

- 3 a Draw a North line at  $P$ , then join  $P$  to  $Q$  and measure the angle between the North line and this line:  $060^\circ$  (any value  $058^\circ$  to  $062^\circ$  accepted).




### Properties of 2D shapes

- 1 a



- b trapezium  
c one pair of parallel sides


- 2 a



- b 4  
c square  
d Two from: all sides equal in length; all angles are  $90^\circ$ ; diagonals are equal; diagonals bisect each other at  $90^\circ$

3 a rectangle, rhombus

b




### Congruent shapes


- 1 D and F are exactly the same as A – they are the same size and shape (it doesn't matter that they are rotated). E is similar to A, not congruent – it is smaller.
- 2 If the triangles are congruent, all three angles must be the same in both. You know that two of the angles are  $35^\circ$  and  $82^\circ$ , so  $x = 180 - 35 - 82 = 63^\circ$ .
- 3 a Identify what values match: SAS (side, angle, side – two sides and the angle between them).  
b Identify what values match: ASA (angle, side angle – two angles and a corresponding side).
- 4 They have the same angles, and they **look** congruent, but we cannot be sure of this. We are not given measurements to prove the corresponding sides are the same length – one triangle could be a slight enlargement of the other. So the answer is no.

### Constructions


1 a and b



2 a and b



3 a



b Distance on the diagram = 2.5 cm

$$2.5 \times 100 = 250 \text{ cm in real life} = 2.5 \text{ m}$$

### Drawing circles and parts of circles

1 a-d




Diagram not to scale

2 a and b





Diagram not to scale


- 3 No, Donald is not correct. A segment of a circle is the area enclosed by a chord and an arc; a sector of a circle is the area enclosed by two radii and the arc between them.





### Loci

1 a



b






**Perimeter**

- A hexagon has 6 sides so perimeter =  $6 \times 9 = 54$  cm
- Missing vertical length =  $20 - 5 - 5 - 4 = 6$  mm  
Missing horizontal lengths are all equal =  $25 - 13 = 12$  mm each  
Perimeter =  $20 + 25 + 6 + 12 + 4 + 12 + 5 + 12 + 5 + 13 = 114$  mm  
 $114 \div 10 = 11.4$  cm
- Perimeter of cushion =  $\frac{1}{2} \times 2 \times \pi \times 24 + 30 + 48 + 30 = 183$  cm (to nearest cm)  
= 1.83 m. So no, Greta does not have enough lace.

**Area**

- Area =  $12 \times 6 = 72$  cm<sup>2</sup>
  - Area =  $\frac{1}{2} \times (3 + 8) \times 4 = 22$  cm<sup>2</sup>
  - Area of rectangle =  $2 \times 10 = 20$  cm<sup>2</sup>  
Area of trapezium =  $\frac{1}{2} (a + b)h = \frac{1}{2} (2.5 + 10)10 = 62.5$  cm<sup>2</sup>  
Area of shape =  $20 + 62.5 = 82.5$  cm<sup>2</sup>
- First draw a diagram. Two sides are equal, and are 6 cm. The two other sides are equal, and are  $x$  cm.




- $$x + x + 6 + 6 = 16$$
- $$2x + 12 = 16$$
- $$x = \frac{16 - 12}{2} = 2.$$
- Area =  $6 \times 2 = 12$  cm<sup>2</sup>
- Radius of circle = 3.5 cm  
Area of circle =  $\pi r^2 = \pi \times 3.5^2 = 38.48$  cm<sup>2</sup>  
Area of square = 49 cm<sup>2</sup>  
Area of shaded part =  $\frac{49 - 38.48}{4} = 2.63$  cm<sup>2</sup>

**Sectors**

- $\frac{120}{360} = \frac{1}{3}$
  - Area =  $\frac{1}{3} \times \pi \times 3^2 = 3\pi$  cm<sup>2</sup>
- Area =  $\frac{1}{4} \times \pi \times 2.8^2 = 6.2$  cm<sup>2</sup> (1 d.p.)
  - Perimeter =  $2.8 + 2.8 + \frac{1}{4} \times 2 \times \pi \times 2.8 = 10.0$  cm (1 d.p.)
- Area =  $\frac{40}{360} \times \pi \times 5^2 = 8.73$  cm<sup>2</sup> (2 d.p.)
  - Arc AB =  $\frac{40}{360} \times 2 \times \pi \times 5 = 3.49$  cm (2 d.p.)





**3D shapes**



b triangular prism

c

Number of faces	Number of edges	Number of vertices
5	9	6



3 b The front elevation shows 5 cubes and the side shows that the shape is 2 cubes deep.  $5 \times 2 = 10$ , so 10 cubes make up the shape.

### Volume

1 The front elevation shows 5 cubes and the side shows that the shape is 4 cubes deep. Volume =  $5 \times 4 = 20 \text{ cm}^3$

2 a Volume = area of cross-section  $\times$  length =  $\pi \times 5^2 \times 12 = 942 \text{ cm}^3$  (to 3 s.f.)

b Volume =  $\frac{1}{3} \times$  area of cross-section  $\times$  length  
 $= \frac{1}{3} \times \pi \times 7^2 \times 15 = 770 \text{ cm}^3$  (to nearest cm)

3 Volume =  $\frac{1}{2} \times \frac{4}{3} \pi r^3 = \frac{2}{3} \times \pi \times 8^3 = 1072.33 \text{ cm}^3$  (to 2 d.p.)

4 Volume of tank =  $40 \times 40 \times 60 = 96\,000 \text{ cm}^3$

Volume of water in tank, 80% full =  $0.8 \times 96\,000 = 76\,800 \text{ cm}^3$

Height of water in pond (1st fill) =  $76\,800 \div (80 \times 60) = 16 \text{ cm}$

Height of water in pond (2nd fill) =  $16 \times 2 = 32 \text{ cm}$

Height of water in pond (3rd fill) =  $16 \times 3 = 48 \text{ cm}$

Three tanks of water are needed to fill the pond.

Alternative method: divide volume of pond by volume of water in tank.

$$\frac{80 \times 60 \times 48}{76\,800} = 3$$

### Surface area

1 a 6 faces

b Surface area =  $60 + 60 + 5 + 5 + 3 + 3 = 136 \text{ cm}^2$

2 surface area = area of triangular side  $\times 4$  + area of square base

$$= \left(\frac{1}{2} \times 6 \times 5\right) \times 4 + 6^2$$

$$= (15 \times 4) + 36$$

$$\text{Surface area} = 96 \text{ cm}^2$$

3 a Surface area of a sphere =  $4\pi r^2 = 4 \times \pi \times 14^2 = 2463.01 \text{ cm}^2$  (to 2 d.p.)

b Surface area of a cone =  $\pi r l + \pi r^2 = \pi \times 6 \times 10 + \pi \times 6^2 = 301.59 \text{ cm}^2$  (to 2 d.p.)

4 Area of cylinder =  $2\pi r h = 2\pi \times 6 \times 1.5 = 56.55 \text{ cm}^2$

Area of circular base =  $\pi r^2 = \pi \times 6^2 = 113.10 \text{ cm}^2$

Area of curved surface area of cone =  $\pi r l = \pi \times 6 \times 11.5 = 216.77 \text{ cm}^2$

Total surface area =  $56.55 + 113.10 + 216.77 = 386.42 \text{ cm}^2$  (to 2 d.p.)

### Using Pythagoras' theorem

1  $x^2 = 3^2 + 4^2$

$$= 9 + 16$$

$$= 25$$

$$x = \sqrt{25}$$

$$= 5 \text{ cm}$$

$$15^2 = y^2 + 12^2$$

$$225 = y^2 + 144$$

$$81 = y^2$$

$$y = \sqrt{81}$$

$$= 9 \text{ cm}$$

2  $6^2 = 4.5^2 + w^2$

$$36 = 20.25 + w^2$$

$$w = \sqrt{15.75}$$

$$= 3.97 \text{ cm}$$

$$\text{Area} = l \times w = 4.5 \times 3.97 = 17.9 \text{ cm}^2$$

3  $AB^2 = 2^2 + 4^2$

$$= 4 + 16$$

$$= 20$$

$$AB = \sqrt{20}$$

$$= \sqrt{4 \times 5}$$

$$= 2\sqrt{5} \text{ cm}$$

4 Square of diagonal of doorway =  $70^2 + 190^2 = 41\,000$

$$\text{Diagonal of doorway} = \sqrt{41\,000}$$

$$= 202.48 \text{ cm} = 2.0248 \text{ m} = 2.02 \text{ m} \text{ (2 d.p.)}$$

Yes, the artwork will fit through the diagonal of the doorway.

### Trigonometry

1  $\tan x = \frac{8}{13}$

$$x = 31.6^\circ$$

2  $\cos 42 = \frac{17}{AC}$

$$AC = \frac{17}{\cos 42} = 22.9 \text{ cm}$$

3  $\sin 49 = \frac{h}{6}$

$$h = 6 \sin 49$$

$$= 4.53 \text{ m}$$

### Exact trigonometric values

1 a  $\tan x = \frac{\text{opposite}}{\text{adjacent}} = \frac{1}{1} = 1$

b  $x = \tan^{-1}(1) = 45^\circ$

2  $\cos 30 = \frac{\sqrt{3}}{PR}$

$$\frac{\sqrt{3}}{2} = \frac{\sqrt{3}}{PR}$$

$$PR = 2 \text{ cm}$$

3  $\frac{YZ}{20} = \sin 30$

$$YZ = 20 \sin 30$$

$$= 20 \times \frac{1}{2}$$



$$= 10 \text{ cm}$$

4  $\sin 45^\circ = \cos 45^\circ = \frac{1}{\sqrt{2}}$


So the second angle in the triangle must be  $45^\circ$ .

The third angle will be  $180 - 90 - 45 = 45^\circ$ .

**Transformations**



3 Rotation 90° clockwise about (1, -1); or rotation 270° anticlockwise about (1, -1).



c Reflection in  $y = 1$


**Similar shapes**

- 1 a  $YZW$   
 b Scale of enlargement =  $\frac{\text{enlarged length}}{\text{original length}} = \frac{6.3}{2.1} = 3$   
 c  $WZ = 4 \times 3 = 12 \text{ cm}$   
 2 a  $37.5^\circ$   
 b Scale of enlargement =  $\frac{\text{enlarged length}}{\text{original length}} = \frac{5}{2.5} = 2$   
 Length of  $AB = 8 \div 2 = 4 \text{ cm}$   
 c They are isosceles, because they have two equal sides and two equal angles. Note that the diagrams are not drawn to scale, as is common practice in maths questions – you have to go by the numbers.  
 d Length of  $BC = \text{length of } AC = 2.5 \text{ cm}$   
 3 a Scale of enlargement =  $\frac{\text{enlarged length}}{\text{original length}} = \frac{4}{6} = \frac{2}{3}$   
 b Length of  $RT = 4.5 \times \frac{2}{3} = 3 \text{ cm}$   
 4 a Scale of enlargement =  $\frac{\text{enlarged length}}{\text{original length}} = \frac{2.2}{4.4} = \frac{1}{2}$   
 b Length of  $CE = \text{length of } CD \div \frac{1}{2} = 2.8 \times 2 = 5.6 \text{ cm}$   
 c Angle  $ACE = 180 - 70 - 64 = 46^\circ$

**Vectors**

- 1 a  $\mathbf{a} = \begin{pmatrix} 3 \\ 2 \end{pmatrix}$  b  $\mathbf{b} = \begin{pmatrix} -3 \\ 3 \end{pmatrix}$  c  $\mathbf{c} = \begin{pmatrix} 2 \\ -4 \end{pmatrix}$  d  $\mathbf{d} = \begin{pmatrix} -4 \\ -2 \end{pmatrix}$   
 2 a
- 

- b  $-\mathbf{p} = \begin{pmatrix} 6 \\ -1 \end{pmatrix}$   
 c  $2\mathbf{p} = \begin{pmatrix} -12 \\ 2 \end{pmatrix}$   
 d  $2\mathbf{p} + \mathbf{p} = \begin{pmatrix} -12 \\ 2 \end{pmatrix} + \begin{pmatrix} -6 \\ 1 \end{pmatrix} = \begin{pmatrix} -18 \\ 3 \end{pmatrix}$   
 $3\mathbf{p} = 3 \times \begin{pmatrix} -6 \\ 1 \end{pmatrix} = \begin{pmatrix} -18 \\ 3 \end{pmatrix}$   
 Therefore,  $2\mathbf{p} + \mathbf{p} = 3\mathbf{p}$   
 3 a  $\mathbf{a} + \mathbf{b} = \begin{pmatrix} 3 \\ 4 \end{pmatrix} + \begin{pmatrix} -5 \\ -1 \end{pmatrix} = \begin{pmatrix} 3-5 \\ 4-1 \end{pmatrix} = \begin{pmatrix} -2 \\ 5 \end{pmatrix}$   
 b  $\mathbf{c} = \mathbf{a} - \mathbf{b} = \begin{pmatrix} 3 \\ 4 \end{pmatrix} - \begin{pmatrix} -5 \\ -1 \end{pmatrix} = \begin{pmatrix} 3-(-5) \\ 4-(-1) \end{pmatrix} = \begin{pmatrix} 8 \\ 3 \end{pmatrix}$



**Probability**

**Basic probability**

- 1 Probability =  $\frac{\text{number of successful outcomes}}{\text{total number of possible outcomes}} = \frac{1}{10}$   
 2  $P(\text{not rain}) = 1 - P(\text{rain}) = 1 - 0.6 = 0.4$   
 3 a b c
- 
- a  $P(\text{a number from 1 to 8}) = \frac{8}{8} = 1$   
 b  $P(\text{a multiple of 3}) = \frac{2}{8} = \frac{1}{4}$   
 c  $P(\text{a number greater than 3}) = \frac{5}{8}$   
 4  $2p - 0.1 + 2p + 0.1 + p = 1$   
 $5p = 1$   
 $p = 0.2$

Outcome	Red	Blue	Green
Probability	$2p - 0.1$ $= 2 \times 0.2 - 0.1$ $= 0.3$	$2p + 0.1$ $= 2 \times 0.2 + 0.1$ $= 0.5$	$p = 0.2$

Blue is most likely.

**Two-way tables and sample space diagrams**

- 1 Work out the missing values one by one, for example in the order shown from first to seventh. (There is more than one order you can do it in.)

	Single	Double	King	Totals
Oak	2	Fourth: $42 - 12 =$ $14 = 16$	Fifth: $30 - 16 =$ $2 = 12$	30
Pine	First: $54 - 14 =$ $17 = 23$	14	17	54
Walnut	1	12	Sixth: $32 - 12 =$ $17 = 3$	Seventh: $1 + 12 +$ $3 = 16$
Totals	Second: $2 + 23 +$ $1 = 26$	Third: $100 - 26$ $= 32 = 42$	32	100

2 a

		Spinner			
		1	2	3	4
Coin	Heads	1, H	2, H	3, H	4, H
	Tails	1, T	2, T	3, T	4, T

- b  $P(\text{tail}, 1) = \frac{1}{8}$   
 c  $P(\text{head, number greater than 1}) = \frac{3}{8}$

3 a


	Study sciences	Do not study sciences	Totals
Boys	$\frac{4}{5} \times 120 = 24$	$45 - 24 = 21$	$\frac{3}{8} \times 120 = 45$
Girls	$75 - 40 = 35$	$\frac{4}{3} \times 120 = 40$	$120 - 45 = 75$
Totals	$24 + 35 = 59$	$21 + 40 = 61$	120

- b 21 boys do not study science, so probability =  $\frac{21}{120} = \frac{7}{40}$   
 c There are 75 girls and 35 of them study science, so probability =  $\frac{35}{75} = \frac{7}{15}$

### Sets and Venn diagrams

- 1 a  $\xi = \{21, 22, 23, 24, 25, 26, 27, 28, 29\}$   
 b  $A = \{21, 24, 27\}$   
 c  $B = \{24, 28\}$   
 d  $A \cup B = \{21, 24, 27, 28\}$  – A 'union' B means **all** the values in A and **all** the values in B  
 e  $A \cap B = \{24\}$  – A 'intersect' B means only those value that are in **both** A and B.


2 a



- b There are 7 adults who only go to art class, so probability =  $\frac{7}{20}$   
 c 8 adults only go to yoga class and 7 adults only go to art class.  
 Probability =  $\frac{8+7}{20} = \frac{3}{4}$   
 3 a Total number of respondents =  $\zeta$   
 $= 25 + 11 + 2 + 12 + 0 + 7 + 3 = 60$   
 $P(\text{supermarket only}) = \frac{25}{60} = \frac{5}{12}$   
 b  $P(F) = \frac{12+2+7+3}{60} = \frac{24}{60} = \frac{2}{5}$   
 c  $P(L \cap F) = \frac{2+7}{60} = \frac{9}{60} = \frac{3}{20}$   
 d  $P(S') = \frac{0+7+3}{60} = \frac{10}{60} = \frac{1}{6}$


### Frequency trees and diagrams

1 a




- b  $P(W, B) = \frac{3}{4} \times \frac{1}{4} = \frac{3}{16}$   
 c  $P(B, B) = \frac{1}{4} \times \frac{1}{4} = \frac{1}{16}$

2 a



- b  $P(A, O) = \frac{4}{6} \times \frac{2}{3} = \frac{8}{9}$   
 c  $P(O, O) = \frac{2}{6} \times \frac{3}{4} = \frac{1}{2}$   
 d  $P(O, O) + P(A, A) = \frac{2}{30} + \left(\frac{4}{6} \times \frac{3}{5}\right) = \frac{2}{30} + \frac{12}{30} = \frac{14}{30} = \frac{7}{15}$

3 a



- b  $P(R, R) = \frac{3}{10} \times \frac{1}{2} = \frac{3}{20}$   
 c  $P(R, R') + P(R', R) = \left(\frac{3}{10} \times \frac{1}{2}\right) + \left(\frac{7}{10} \times \frac{1}{2}\right)$   
 $= \frac{3}{20} + \frac{7}{20} = \frac{10}{20} = \frac{1}{2}$   
 The probability of one day having rain and one day having no rain is 50%.

### Expected outcomes and experimental probability

- 1 a The spinner was spun  $12 + 13 + 10 + 15 = 50$  times.  
 b Estimated probability of blue =  $\frac{13}{50}$   
 c Estimated probability of yellow =  $\frac{15}{50} = \frac{3}{10}$   
 d You would expect  $\frac{10}{15} \times 100 = 20$  green outcomes from 100 spins.  
 2  $0.75 \times 20 = 15$  students would be expected to pass the exam.  
 3 a Total number of customers =  $26 + 20 + 6 + 5 + 3 = 60$   
 Estimated probability that someone will buy stamps =  $\frac{20}{60} = \frac{1}{3}$   
 b  $\frac{1}{3} \times 450 = 150$  customers buy stamps each day  
 c  $\frac{6}{60} \times 450 = 45$  customers buy foreign currency each day  
 d  $450 \times 6 = 2700$  customers each week  
 $\frac{5}{60} \times 2700 = 225$  customers use the post office for banking each week

## Statistics

### Data and sampling

- 1 65, because that is 10% of 650 (the entire population).  
 2  $\frac{50}{25000} \times 100 = 0.2\%$ . The sample is not big enough.  
 People in the town centre may not be the only ones using buses. For example, some people may take buses to the local train station, school or hospital.  
 3 a  $\frac{9}{45} \times 400 = 80$  people like carrot cake  
 Sam needs to make 80 cakes.



**b Assumptions**

Assumed the sample is representative of the population; this could affect the answer because not all the 400 people who have accepted the invitation may turn up.

Assumed that these are individual carrot cakes. If instead each cake is a large one divided into 8 slices, then only  $80 \div 8 = 10$  carrot cakes would be needed.

**Frequency tables**

- 1 **a**  $1 + 11 + 9 + 6 + 1 + 2 = 30$  tables  
**b**  $(1 \times 1) + (2 \times 11) + (3 \times 9) + (4 \times 6) + (5 \times 1) + (6 \times 2) = 91$  people

2

Number of electronic devices	Tally	Frequency
0–1		3
2–3		10
4–5		5
6–7		5
8–9		1


- 3 **a** Continuous

**b**

Mass, $m$ (kg)	Tally	Frequency
$50 \leq m < 60$		3
$60 \leq m < 70$		5
$70 \leq m < 80$		4
$80 \leq m < 90$		5
$90 \leq m < 100$		3

**Bar charts and pictograms**

- 1 **a**  $9 - 4 = 5$  more boys than girls prefer squash  
**b**  $15 + 6 + 9 + 7 + 4 = 41$  girls were surveyed




- 3 **a**  $3 + 3 + 3 + 3 + 2 = 14$  hours  
**b** Yorkshire gets 9 hours, Inverness-shire gets 7 hours. Yorkshire gets 2 more hours of sunshine each day than Inverness-shire.

**Pie charts**

- 1 **a**  $\frac{150}{360} \times 300 = 125$  people last saw an action movie  
**b** Romance =  $90^\circ$ , so  $\frac{3}{4} \times 300 = 225$  people did not see a romance movie.

2

	Children	Adults	Senior citizens
Calculation	$\frac{80}{250} \times 360$ $= 115.2$	$\frac{155}{250} \times 360$ $= 223.2$	$\frac{15}{250} \times 360$ $= 21.6$
Angle	$115^\circ$	$223^\circ$	$22^\circ$



- 3 Angle for 8:30–9pm =  $72^\circ$   
 $\frac{360}{72} \times 12 = 60$  people were surveyed

**Stem and leaf diagrams**

- 1 **a** 71 cm – this is the highest stem value combined with the highest leaf value  
**b** 20 – count the leaf values  
 2 **a** Longest – shortest =  $4.26 - 2.03 = 2.23$  m  
**b** 3 girls jumped more than 3.5 m (3.56 m, 3.74 m and 4.06 m)  
 3 Write all the amounts in order, and convert them to pounds.

Year 9s	0.50	0.80	0.95	1.30	1.75	2.00	2.65	3.05
Year 10s	0.65	0.75	1.50	1.85	2.70	3.10		

Year 9s	Year 10s
95 80 50	0 65 75
75 30	1 50 85
65 00	2 70
05	3 10

Key
Year 9s
50 0 means £0.50
Year 10s
0 65 means £0.65

**Measures of central tendency: mode**

- 1 There are two modes: 3 minutes and 4 minutes, since each appears twice. An alternative correct answer is to say that there is no mode.  
 2 This is the one with the highest frequency:  $12 < a \leq 13$ .  
 3 This is the class with the biggest slice of the pie: £10–£20.  
 4 This is the one represented the most times in the diagram: 25 kg.

**Measures of central tendency: median**

- 1 Ages in order: 11 11 12 13 13 13 13 (14 15) 15 16 16 17 17 18 18  
 Median =  $14\frac{1}{2}$  years old  
 2 Total frequency =  $5 + 12 + 17 + 10 + 6 = 50$   
 Median =  $\frac{50 + 1}{2} = 25.5$ th person  
 Median class =  $12 < a \leq 13$   
 3 Total frequency = 40  
 Median =  $\frac{40 + 1}{2} = 20.5$ th person  
 Median age =  $\frac{28 + 29}{2} = 28\frac{1}{2}$  years old

**Measures of central tendency: mean**

- 1 Mean age =  $\frac{6 + 7 + 11 + 13 + 18}{5} = 11$  years old  
 2 Number of bedrooms =  $(1 \times 4) + (2 \times 7) + (3 \times 13) + (4 \times 17) = 125$   
 Number of houses =  $4 + 7 + 13 + 17 = 41$   
 Mean number of bedrooms =  $125 \div 41 = 3.05 \approx 3$  bedrooms

- 3 Number of holidays =  $(0 \times 4) + (1 \times 21) + (2 \times 9) + (3 \times 2) = 45$   
 Number of employees =  $4 + 21 + 9 + 2 = 36$   
 Mean number of holidays =  $45 \div 36 = 1.25 \approx 1$  holiday

4

Age of patients, $a$	Midpoint	Frequency	Midpoint $\times$ frequency
$0 < a \leq 10$	5	3	15
$10 < a \leq 20$	15	18	270
$20 < a \leq 30$	25	6	150
$30 < a \leq 40$	35	11	385
$40 < a \leq 50$	45	10	450
$50 < a \leq 60$	55	19	1045
$60 < a \leq 70$	65	16	1040
$70 < a \leq 80$	75	17	1275
		Total = 100	Total = 4630

Mean age of patients =  $\frac{4630}{100} = 46.3 \approx 46$  years old

### Range


- 1 a Range of boys' ages =  $4 - 2 = 2$  years  
 b Range of boys' ages =  $4 - 1 = 3$  years
- 2 a Range in temperatures for Resort A =  $25 - 16 = 9^\circ\text{C}$   
 b Range in temperatures for Resort B =  $28 - 13 = 15^\circ\text{C}$
- 3 a Business A: Range =  $45\,816 - 23\,561 = \pounds 22\,255$   
 Mean profit =  $\frac{23\,561 + 30\,485 + 39\,210 + 45\,816}{4} = \pounds 34\,768$   
 b Business B: Range =  $63\,248 - 17\,894 = \pounds 45\,354$   
 Mean profit =  $\frac{32\,820 + 40\,328 + 17\,894 + 63\,248}{4} = \pounds 38\,572.50$   
 c **Either** Business A because its range in profit is lower and the profit is increasing each year, and so it shows a more consistent performance.  
**or** Business B because its mean profit is higher, and its most recent profit (in Year 4) is  $\pounds 17\,432$  more than Business A.

### Comparing data using measures of central tendency and range

- 1 a Mean time for bus journey =  $\frac{32 + 30 + 39 + 32 + 43 + 31}{6} = 34.5$  minutes  
 b Range for bus journey =  $43 - 30 = 13$  minutes  
 c Mean time for train journey =  $\frac{16 + 24 + 18 + 26 + 70 + 17}{6} = 28.5$  minutes  
 d Range for train journey =  $70 - 16 = 54$  minutes  
 e **Either** The bus is better because although it takes longer (on average), the range is lower, and so you can predict the time it takes for the journey.  
**or** The train is better because it is quicker than the bus (on average), although the range suggests it may be less reliable.
- 2 a Mean = 13. This does not represent the age of the people using the playground. In fact, those using the playground are small children (under 10) and their parents (over 25).  
 b There are five modes (3, 4, 5, 7, 8), and so the mode does not represent the age of the people using the playground.  
 c Ages in order: 3 3 4 4 5 5 7 7 8 8 26 30 33 39  
 Median position =  $\frac{14 + 1}{2} = 7.5$ th value  
 Median age = 7 years old
- 3 Mode = 0; Median = 0; Mean = 2 days. Mode or median are the best averages to use, because the mean is skewed by the student who is absent due to sickness for 24 days.


### Time series graphs

- 1 a Start at Day 3, read up to the graph line then left to the weight scale: 190 g  
 b Start at 225 g, read across to the graph line then down: 5 days old  
 c Mean weight gained =  $\frac{\text{total weight gained}}{\text{number of days}} = \frac{270 - 130}{7} = 20$  g per day



- b There is an increase in campers in May. This may be due to May bank holidays, or May half-term, or perhaps there was some very sunny weather.


- 3 a
- Sales figures for two different branches of a DIY store




- b Branch 1 had a steady increase in sales for the first four months. Then sales levelled off to stay at around  $\pounds 25\,000$ . Branch 2 had a slow start to its sales in the first three months. Then perhaps it had a promotion, because sales increased a lot in month 4. Sales have been increasing ever since.

### Scatter graphs

- 1 a Positive correlation. This means as the temperature rises, more pairs of flip flops are sold.  
 b Negative correlation. This means as the temperature rises, fewer wellington boots are sold.



- b** The scatter diagram shows a positive correlation between students' maths and physics test percentages. Therefore, the students who got a low percentage in the maths test got the lower percentages in the physics test; the students who got a high percentage in the maths test got the higher percentages in the physics test.
- c** The outlier is the point marked at (0, 58).
- d** The student was absent for the maths test.



The black line shows the line of best fit.  
 The grey line shows the line where a laptop loses £150 every 6 months.  
 The shop owner is not correct. The line of best fit shows on average a laptop loses approximately £159/£160 every 6 months.

- b** The line of best fit cannot make a prediction outside the available data. The data only goes as far as 18 months.


### Practice papers

#### Non-calculator

- It is in the 1000s column, so 7000.
- $\frac{40}{100} \times 50 = 20$
- No, Sandeep is not correct:  
 $\frac{2}{5} = \frac{4}{10} = 0.4$   
 But  $4\% = 0.04$
- $45 = 5 \times 9 = 5 \times 3 \times 3$   
 So prime factors are 3 and 5.

- Distance on diagram between lamp post A and lamp post B = 6 cm  
 $6 \times 20 = 120$  m
- There are 5 sections on the spinner; 2 are labelled r.  
 $P(r) = \frac{2}{5}$
- Yes. Fun run + music festival = £9689 + £9689 + £6370 = £25 748
- $3 + 7 = 10$  parts in the ratio  
 Milk chocolates =  $\frac{7}{10}$
- Prize A =  $8 \times 4 = 32$  tickets  
 Prize B =  $2 + 4 + 8 + 16 = 30$  tickets  
 Prize A gives more tickets.
- a** There are 2 triangles in Pattern 1, and 4 more are added for each pattern.  
 Number of triangles =  $4p - 2$   
 In Pattern 8, there are  $4 \times 8 - 2 = 30$  triangles  
**b** No, Harry is incorrect. The number of triangles is not the pattern number multiplied by 4. Rather, it is add 4 triangles each time.
- a** £3.50  
**b** £5.00 – appears the most times
- Cost before VAT =  $45 + 3 \times 35 = £150$   
 Cost with VAT =  $150 \times 1.2 = £180$
- Bus 2A will stop at: 10:00, 10:15, 10:30, 10:45, 11:00  
 Bus 2B will stop at: 10:00, 10:12, 10:24, 10:36, 10:48, 11:00  
 They next arrive at the bus stop together at 11 am.
- Total number of patients =  $10 + 45 + 5 = 60$   
 Angle per patient =  $360 \div 60 = 6^\circ$

	Children	Adults	Senior citizens
Number	10	45	5
Angle	$10 \times 6 = 60^\circ$	$45 \times 6 = 270^\circ$	$5 \times 6 = 30^\circ$




- Cost of T shirts =  $8 \times 12 = £96$  (for 48 shirts)  
 Saturday:  $\frac{3}{8} \times 48 = 18$   $18 \times 5 = £90$   
 Sunday: 30 shirts for £3 each  $30 \times 3 = £90$   
 Profit =  $90 + 90 - 96 = £84$
- a**  $\frac{5}{8} - \frac{7}{12} = \frac{15}{24} - \frac{14}{24} = \frac{1}{24}$   
**b**  $2\frac{2}{3} \div \frac{4}{9} = \frac{8}{3} \times \frac{9}{4} = 2 \times 3 = 6$
- a**  $13 - 4x$   
**b**  $(x + 3)(x - 4) = x^2 - 4x + 3x - 12 = x^2 - x - 12$
- $30\% = £45$   
 $10\% = \frac{45}{3} = £15$   
 $100\% = £15 \times 10 = £150$   
 Original price = £150
- a**  $(-3, 0)$   
**b**  $B = (1, 2), C = (3, -4)$   
 Midpoint of BC =  $(\frac{1+3}{2}, \frac{2-4}{2}) = (2, -1)$

- 20 Count the total number of decimal places in both numbers (3). Do the calculation without the decimal points.


$$\begin{array}{r} 815 \\ \times 34 \\ \hline 3260 \\ 24450 \\ \hline 27710 \end{array}$$

Now put 3 decimal places back into your answer: 27.710  
= 27.71

21



22



23  $\frac{8.02 \times 3.76}{15.98} \approx \frac{8 \times 4}{16} \approx 2$

24 a  $K = \frac{1}{2} \times 11 \times 3^2$

$K = 49.5$

b Rearrange the formula:

$$K = \frac{1}{2}mv^2$$

$$v^2 = \frac{2K}{m}$$

$$v = \sqrt{\frac{2K}{m}}$$

Substitute values for  $K$  and  $m$  to find  $v$

$$v = \sqrt{\frac{2 \times 180}{10}}$$

$v = 6$  or  $-6$

25  $x = 180 - (90 + 36) = 54^\circ$

26 a Average speed =  $\frac{\text{total distance travelled}}{\text{time taken}} = \frac{140}{2} = 70$  km/h

b 15 minutes or  $\frac{1}{4}$  hour – when the line is horizontal and no distance is covered

c Fastest speed is where the graph has its highest gradient (i.e. 1.75 to 2 hours)

$$\text{Gradient} = \frac{40}{0.25} = 160$$

Fastest speed = 160 km/h

27  $5x + 3 = 6x - 7$  (as base angles of an isosceles triangle are equal)

$10 = x$

Hence base angles are  $53^\circ$


Other angle =  $180 - (53 + 53) = 74^\circ$

So  $74^\circ = 7y - 10$

Solving gives  $y = 12^\circ$

Hence  $x = 10^\circ$  and  $y = 12^\circ$

28 a and b



c Reflection in  $x = y$ , or rotation of  $90^\circ$  anticlockwise about the point  $(1.5, 1.5)$

29  $b - 3a = \begin{pmatrix} 7 \\ 7 \end{pmatrix} - 3\begin{pmatrix} -1 \\ 2 \end{pmatrix} = \begin{pmatrix} 7 \\ 7 \end{pmatrix} + \begin{pmatrix} 3 \\ -6 \end{pmatrix} = \begin{pmatrix} 10 \\ 1 \end{pmatrix}$

### Calculator

1 On your calculator:  $7 \div 8 = 0.875$ .

2  $350 \div 100 \times 140 = \text{£}490$

3  $\text{£}83.75 \div 5 = \text{£}16.75$

$\text{£}16.75 \times 37 = \text{£}619.75$

4  $255 \times 1.16 = 295.8$  so the price in France is  $\text{€}295.80$ . The games console is cheaper in the UK.

5 On your calculator:  $\sqrt[3]{13.824} + (4.5 - 0.38)^2 = 19.4$  (to 3 s.f.)

6  $4.5^2 = 3.8^2 + XY^2$

$20.25 = 14.44 + XY^2$

$XY^2 = 5.81$

$XY = 2.4$  cm (to 1 d.p.)

7 a Friday

b On Thursday Jeff received 18 work emails and 2 personal emails.

$\frac{2}{20} \times 100 = 10\%$  of his emails on Thursday were personal.

c Total number of emails received =  $8 + 16 + 4 + 15 + 5 + 16 + 2 + 18 + 11 + 10 = 105$

Total number of work emails received =  $16 + 15 + 16 + 18 + 10 = 75$

$\frac{75}{105} = \frac{5}{7}$  of his emails that week were about work.

8 Factors of 12: 1, 2, 3, 4, 6, 12

Multiples of 9: 9, 18, 27, 36, 45, 54, 63, 72, 81, 90...

$1 \times 81 = 81$ ;  $3 \times 27 = 81$

9  $-1, 0, 1, 2, 3, 4$  – the range is less than 5 so this is not included

10 Area to be painted =  $16 \times 1.8 \times 1.8 = 51.84 \text{ m}^2$

2 tins are enough for  $2 \times 20 = 40 \text{ m}^2$  and 3 tins are enough for  $3 \times 20 = 60 \text{ m}^2$ .

$40 < 51.84 < 60$

So Geraldine needs 3 tins, at a price of  $3 \times 22.50 = \text{£}67.50$ .

11  $6 \times 9 \times 7 + \frac{1}{2} \times 6 \times 8 \times 7 = 546 \text{ cm}^3$

12  $84 \text{ km/h} = \frac{84 \times 1000}{60 \times 60} \text{ m/s} = 23.3 \text{ m/s}$  (to 1 d.p.)

13 1 part of the ratio =  $\text{£}30$

Jessie gets  $2 \times 30 = \text{£}60$

14  $PQ^2 = 2^2 + 4^2$

$PQ^2 = 20$

$PQ = \sqrt{20}$

$= \sqrt{4 \times 5}$

$= 2\sqrt{5}$

15  $\frac{2}{3}$  are the 42 women so  $\frac{1}{3}$  is 21 women and

$\frac{3}{3} = 21 \times 3 = 63$

So there are 63 men and women and this represents 50% of the total.

Hence there are 126 people in the audience.

16  $58 \text{ million} = 58 \times 10^6 = 5.8 \times 10^7 \text{ km}$


17 Let number by Seren =  $x$

Number by Tom =  $x + 3$

Number by Rohan =  $x + 3 + 2 = x + 5$

Total number =  $x + x + 3 + x + 5 = 3x + 8$

18 a



b  $P(\text{late and not late}) + P(\text{not late and late}) = \left(\frac{1}{8} \times \frac{7}{8}\right) + \left(\frac{7}{8} \times \frac{1}{8}\right) = \frac{14}{64} = \frac{7}{32}$

19  $\tan x = \frac{8}{5}$   
 $x = \tan^{-1} \frac{8}{5}$   
 $x = 58^\circ$

$\cos 59^\circ = \frac{y}{11}$

$y = 11 \cos 59^\circ$   
 $= 5.67 \text{ cm}$

20 Mean for Team A =  $\frac{(0 \times 3) + (1 \times 4) + (2 \times 4) + (3 \times 2) + (4 \times 1) + (5 \times 2)}{16}$

$= \frac{32}{16} = 2$

Mean for Team B =  $\frac{(0 \times 3) + (1 \times 4) + (2 \times 8) + (3 \times 4) + (4 \times 1)}{20}$

$= \frac{36}{20} = 1.8$

Team A scored more goals on average.

21 a Option A is best value if you plan to visit the gym once per week, as in one month 4 visits cost £20 (or 5 visits cost £25). This is less than the monthly fee of £30.

b The graphs intersect at 6, so the cost is the same for 6 visits.

It is cheaper to pay the monthly fee if you visit the gym more than 6 times.

22 Area of semicircle =  $\frac{1}{2}\pi r^2$

Area of large semicircle =  $\frac{1}{2} \times \pi \times 7.5^2 = \frac{225\pi}{8}$


Area of small semicircles =  $3 \times \left(\frac{1}{2} \times \pi \times 2.5^2\right) = 3 \times \frac{25\pi}{8} = \frac{75\pi}{8}$

Area of lawn =  $\frac{225\pi}{8} - \frac{75\pi}{8} = \frac{150\pi}{8} = 58.9 \text{ m}^2$  (to 3 s.f.)

23 a

x	Operation	y
-2	$2 \times (-2)^2 - 3 =$	5
-1	$2 \times (-1)^2 - 3 =$	-1
0	$2 \times (0)^2 - 3 =$	-3
1	$2 \times (1)^2 - 3 =$	-1
2	$2 \times (2)^2 - 3 =$	5

b



24 a Draw a line of best fit to show that, yes, the car dealer is correct.

b Start at 55 000 miles and read up to your line of best fit, then read across to the price scale. The car dealer should charge £3500 to £4000. (Value varies with students' own graphs.)

25  $(n + 5)(n - 3) = n^2 + 2n - 15$