

AQA Foundation Mathematics

Revision Guide

Full worked solutions


Number

Factors, multiples and primes

1 a 5 b 1, 12 c 1, 5, 45 d 1, 5

2 $70 = 2 \times 5 \times 7$
 $150 = 2 \times 3 \times 5 \times 5$
 HCF = 10, LCM = 1050

3 $2 \times 3^2 \times 5$



a $2 \times 5 = 10$ b $2 \times 2 \times 2 \times 3 \times 5 \times 7 = 840$

5 12 and 18

Ordering integers and decimals

1 a false c true e true*
 b true d true*

Note: parts d and e differ from the answers in the Revision Guide due to an error in our first edition. Although two different numbers can never be equal, the sign \geq means 'greater than or equal to'; similarly, \leq means 'less than or equal to', so the statements are true.

2 $-0.3, -1.5, -2.5, -4.2, -7.2$

3 0.049, 0.124, 0.412, 0.442, 1.002

4 a < b < c >

Calculating with negative numbers

Stretch it! Multiplying three negative numbers together always gives a negative answer.

1 a $-8 - 3 = -11$ d $14 + 4 = 18$
 b 99 e 0
 c -6 f $12 + 15 - 2 = 25$

2 -8 and 9

3 32°C

Multiplication and division

Stretch it!

$$\begin{array}{r} 621 \\ \times 239 \\ \hline 5589 \\ 18630 \\ 124200 \\ \hline 148419 \end{array}$$

1 a
$$\begin{array}{r} 235 \\ \times 9 \\ \hline 2115 \\ \hline \end{array}$$

 2115

b
$$\begin{array}{r} 924 \\ \times 61 \\ \hline 924 \\ 55440 \\ \hline 56364 \end{array}$$
 56364

2 a
$$\begin{array}{r} 047 \\ 5 \overline{)235} \\ \underline{5} \\ 23 \\ \underline{23} \\ 5 \end{array}$$
 47

b
$$\begin{array}{r} 0516 \\ 8 \overline{)4128} \\ \underline{8} \\ 412 \\ \underline{412} \\ 8 \end{array}$$
 516

c
$$\begin{array}{r} 0126 \\ 17 \overline{)2146} \\ \underline{17} \\ 44 \\ \underline{34} \\ 106 \\ \underline{102} \\ 4 \end{array}$$

126 remainder 4, or $126 \frac{4}{17}$

3
$$\begin{array}{r} 033 \\ 8 \overline{)265} \\ \underline{8} \\ 26 \\ \underline{24} \\ 25 \end{array}$$
 remainder 1

a 33 boxes

b 1 pencil

4
$$\begin{array}{r} 091.25 \\ 4 \overline{)365.00} \\ \underline{4} \\ 365 \\ \underline{365} \\ 00 \end{array}$$
 £91.25

5
$$\begin{array}{r} 32 \\ \times 9 \\ \hline 288 \\ \hline \end{array}$$
 £288

6
$$\begin{array}{r} 307.66 \\ 3 \overline{)923.00} \\ \underline{9} \\ 230 \\ \underline{21} \\ 200 \\ \underline{18} \\ 200 \\ \underline{18} \\ 20 \end{array}$$
 $307.\dot{6} = 307\frac{2}{3}$

*This answer differs from the one in the Revision Guide due to an error in our first edition. This answer has now been re-checked and corrected.

$$\begin{array}{r}
 823 \\
 \times 35 \\
 \hline
 4115 \times 5 \\
 24690 \times 30 \\
 \hline
 28805
 \end{array}$$

28 805

$$\begin{array}{r}
 037 \\
 12 \overline{)450} \\
 \underline{36} \\
 90 \\
 \underline{84} \\
 6
 \end{array}$$

37 remainder 6, so
37 boxes

9 He has not placed a zero in the ones column before multiplying through by 5. The $\times 50$ line should have 5 digits: 36300, so his final three rows of working should look like this:

$$\begin{array}{r}
 1452 \times 2 \\
 36300 \times 50 \\
 \hline
 37752
 \end{array}$$

Calculating with decimals

Stretch it! $3.2 + 7.5 \times 2 = 3.2 + 15 = 18.2$

1 a
$$\begin{array}{r}
 3.40 \\
 -1.07 \\
 \hline
 2.33
 \end{array}$$
 2.33

c $5 \times 7 = 35$
 $0.05 \times 0.7 = 0.035$

b
$$\begin{array}{r}
 19.300 \\
 + 5.091 \\
 \hline
 24.391
 \end{array}$$

24.391

d
$$\begin{array}{r}
 321 \\
 \times 19 \\
 \hline
 2889 \times 9 \\
 3210 \times 10 \\
 \hline
 6099
 \end{array}$$

$3.21 \times 1.9 = 6.099$

e
$$\begin{array}{r}
 1.563 \\
 5 \overline{)7.815}
 \end{array}$$
 1.563

2
$$\begin{array}{r}
 64 \\
 \times 24 \\
 \hline
 256 \\
 1280 \\
 \hline
 1536
 \end{array}$$
 $24 \times 64 = 1536$
 $24 \times \text{£}0.64 = \text{£}15.36$
 $\text{£}20 - \text{£}15.36 = \text{£}4.64$

3
$$\begin{array}{r}
 27.46 \\
 3 \overline{)82.38} \\
 \hline
 27.46
 \end{array}$$
 $\text{Erica } 2 \times 27.46 = \text{£}54.92$
 $\text{Freya } 82.38 - 54.92 = \text{£}27.46$

Rounding and estimation

Stretch it! a 1.0 b 1.00 c 1.000
All the answers are 1

Stretch it! $6.5 \times 8.5 = 55.25 \text{ m}^2$ – an overestimate.

1 a 0.35 c 32.6
b 10 d 33100

2 a $150 \leq x < 250$ c $3.15 \leq x < 3.25$
b $5.5 \leq x < 6.5$ d $5.055 \leq x < 5.065$

3 $\frac{30}{0.5 \times 6} = 10$

4 b is false since $18 \times 1 = 18$ so 18×0.9 cannot be 1.62
c is false because if you divide by a number smaller than 1, the answer will be larger.

5 Night-time low tariff: $2.32 \text{ units} \times 1.622\text{p} = 3.76\text{p}$
 $7.151 \text{ units} \times 2.315\text{p} = 16.55\text{p}$
20.31p

One tariff: $2.320 + 7.151 = 9.471 \text{ units}$
 $9.471 \text{ units} \times 1.923\text{p} = 18.21\text{p}$
Tarik should choose One tariff, since it is cheaper.

Converting between fractions, decimals and percentages

Stretch it! $0.\dot{1}$, $0.\dot{2}$, $0.\dot{3}$, ... $0.\dot{4}$, $0.\dot{5}$. The number of nines is the same as the digit that recurs. The exception is $\frac{9}{9}$ which is the same as 1.

1 a $\frac{32}{100} = \frac{8}{25}$ c $\frac{33}{100}$
b $1\frac{24}{100} = 1\frac{6}{25}$ d $\frac{95}{100} = \frac{19}{20}$

2 a
$$\begin{array}{r}
 0.41666\dots \\
 12 \overline{)5.00000}
 \end{array}$$
 0.416

b
$$\begin{array}{r}
 0.375 \\
 8 \overline{)3.0000}
 \end{array}$$
 0.375

c 0.49
d 0.185

e
$$\begin{array}{r}
 0.42857142\dots \\
 7 \overline{)3.00000000}
 \end{array}$$
 0.428571

3 a $\frac{91}{100} = 91\%$
b $\frac{3}{10} = \frac{30}{100} = 30\%$
c $\frac{4}{5} = \frac{80}{100} = 80\%$
d $\frac{9}{15} = \frac{3}{5} = \frac{6}{10} = \frac{60}{100} = 60\%$

4 $\frac{3}{8} \frac{0.375}{8 \overline{)3.0000}} = 0.375 = 37.5\%$

5 $0.35 \frac{2}{5} = \frac{4}{10} = 0.4$ $30\% = 0.3$
 $30\%, 0.35, \frac{2}{5}$

6 $\frac{15}{20} = \frac{75}{100} = 75\%$ – Amy
Rudi's mark was higher.

Ordering fractions, decimals and percentages

1 $\frac{1}{3} = \frac{8}{24}$ $\frac{3}{8} = \frac{9}{24}$ $\frac{7}{12} = \frac{14}{24}$
 $\frac{7}{12}, \frac{3}{8}, \frac{1}{3}$

2 $-2.2, 7, \frac{1}{5} = 0.2, -\frac{1}{10} = -0.1, 15\% = 0.15,$
 $1\% = 0.01, 0.1$
In order, this is: $-2.2, -\frac{1}{10}, 1\%, 0.1, 15\%, \frac{1}{5}, 7.$
The middle value is 0.1

3 Yes. If the numerator of a fraction is half the denominator then the fraction is equivalent to $\frac{1}{2}$.
If the numerator is smaller than this the fraction must be less than $\frac{1}{2}$.

Standard form

- 1 a 45 000 000 b 0.091
 2 a 6.45×10^8 b 7.9×10^{-8}
 3 $350000 - 4200 = 345800$
 4 $3.2 \times 10^2 = 320$ $3.1 \times 10^{-2} = 0.031$
 $3.09 \times 10 = 30.9$ $3 + (2.1 \times 10^2) = 213$
 In order, this gives: 3.1×10^{-2} 3.09×10
 $3 + (2.1 \times 10^2)$ 3.2×10^2
 5 3×10^8 m/s
 6 $200 \times 1.1 \times 10^{-4} = 2.2 \times 10^{-2} = 0.022$ m = 2.2 cm

Listing strategies

Stretch it!

red + small, red + medium, red + large,
 green + small, green + medium, green + large,
 blue + small, blue + medium, blue + large.

- 1 111,
 112, 121, 211, 113, 131, 311,
 222
 221, 212, 122, 223, 232, 322
 333
 331 313 133 332 323 233
 123 132 213 231 312 321
 2 444 446 449
 464 466 469
 494 496 499
 3 Small A, Small B, Small C, Small D
 Medium A, Medium B, Medium C, Medium D
 Large A, Large B, Large C, Large D.

Review it!

- 1 7 and 6 (or 11 and 2, where both are prime and 2 is also a factor of 12)
 2 $620 = 2 \times 2 \times 5 \times 31 = 2^2 \times 5 \times 31$
 3 $18 = 2 \times 3 \times 3$
 $36 = 2 \times 2 \times 3 \times 3$
 $40 = 2 \times 2 \times 2 \times 5$
 HCF = 2
 4 -11.5, -8.3, -3.5, -3.2, 1.4
 5 a
$$\begin{array}{r} 32.99 \\ +18.74 \\ \hline 51.73 \end{array}$$
 £51.73
 b
$$\begin{array}{r} 18.33 \\ 3 \overline{)54.99} \end{array}$$
 £18.33
 c (for a) $51.73 - 32.99 = 18.74$
 or $51.73 - 18.74 = 32.99$
 (for b) $18.33 \times 3 = 54.99$

6 a
$$\begin{array}{r} 23 \\ \times 14 \\ \hline 92 \times 4 \\ 230 \times 10 \\ \hline 322 \end{array}$$
 $23 \times 0.14 = 3.22$

b
$$\begin{array}{r} 149 \\ \times 27 \\ \hline 1043 \times 7 \\ 2980 \times 20 \\ \hline 4023 \end{array}$$
 4023

7 $81 \div 3 = 27$

8
$$\begin{array}{r} 031 \\ 11 \overline{)3^3 4^4 5^5} \end{array}$$
 remainder 4 $31 \frac{4}{11}$

9 a
$$\begin{array}{r} 0.375 \\ 8 \overline{)3.0000} \end{array}$$
 0.375

b $0.7 \times 100 = 70\%$

10 a $70\% = \frac{70}{100} = \frac{7}{10}$

b $0.8 = \frac{8}{10} = \frac{4}{5}$

11 $\frac{1}{2} = \frac{2}{4}$ $\frac{1}{2}$ is larger

$\frac{2}{7} = \frac{8}{28}$ $\frac{1}{4} = \frac{7}{28}$ $\frac{2}{7}$ is larger

$\frac{3}{11} = \frac{12}{44}$ $\frac{1}{4} = \frac{11}{44}$ $\frac{3}{11}$ is larger

$\frac{2}{5} = \frac{8}{20}$ $\frac{1}{4} = \frac{5}{20}$ $\frac{2}{5}$ is larger

All of them.

12 a $\frac{3}{5} + \frac{1}{7} = \frac{21}{35} + \frac{5}{35} = \frac{26}{35}$

b $2 \frac{1}{5} - \frac{7}{10} = \frac{11}{5} - \frac{7}{10} = \frac{22}{10} - \frac{7}{10} = \frac{15}{10} = 1 \frac{1}{2}$

c $\frac{2}{3} \div \frac{4}{9} = \frac{2}{3} \times \frac{9}{4} = \frac{3}{2} = 1 \frac{1}{2}$

13 $0.25 - 0.07 = 0.18 = \frac{18}{100} = \frac{108}{600}$

$\frac{2}{3} - \frac{1}{2} = \frac{4}{6} - \frac{3}{6} = \frac{1}{6} = \frac{100}{600}$

0.25 - 0.07 is larger

14 $\frac{3}{5} \times \frac{5}{4} = \frac{3}{4}$

15 $\frac{45}{1000} = \frac{9}{200}$

16 $8.6 \div 100 = 0.086$

$0.086 \times 25 = \text{£}2.15$

17 a 9 b 5

18 $25 < 30 < 36$, so:

$\sqrt{25} < \sqrt{30} < \sqrt{36}$

$\sqrt{25} = 5$ (or -5) and $\sqrt{36} = 6$ (or -6)

So $5 < \sqrt{30} < 6$

It lies between 5 and 6.

19 a 3.4×10^9 b 3.04×10^{-7}

20 $37.55 \leq x < 37.65$

21 a 51 b 12, 15, 21, 51, 25, 52

- 22 a** $200 \times 9 \times 10 = 18000 = \text{£}180.00$
b Underestimate since all numbers were rounded down.
- 23** 40% of 600 = 240
 $\frac{1}{5}$ of 600 = 120
 $600 - (240 + 120) = 240$
- 24** More than 33%, less than 50%, multiple of 5. 35%
- 25** No, since 2 is even and a prime number, and odd + odd + even = even.
- 26 a** $4 \times 10^8 \times 2.7 \times 10^2$
 $= 4 \times 2.7 \times 10^8 \times 10^2$
 $= 10.8 \times 10^{8+2}$
 $= 10.8 \times 10^{10}$
 $= 1.08 \times 10^{11}$
b $1.8 \times 10^5 \div (6 \times 10^2)$
 $= (1.8 \div 6) \times (10^5 \div 10^2)$
 $= 0.3 \times 10^{5-2}$
 $= 0.3 \times 10^3$
 $= 3 \times 10^2$
- 27** $0.8 \times 349 = \text{£}279.20$
- 28 a** 3.1 **b** 3.05
- 29 a** 325 000 **b** 320 000
- 30** $3 \times 3 \times 3 \times 3 \times 3 \times 3 = 729$
- 31 a** $26.25 + 18.23 + (4 \times 5.5) = \text{£}66.48$
 $\text{£}66.48 \div 4 = \text{£}16.62$
- 32** $0.19 \times 18\ 000 = 3420$
- 33 a** 2010 and 2011
b $1.1 \times 102.3 = 112.53$

Algebra

Understanding expressions, equations, formulae and identities

- 1 a** $3a + 6 = 10$ (It can be solved to find the value of a .)
b $C = \pi D$ (The value of C can be worked out if the value of D is known.)
c $3(a + 2)$ (It does not have an equals sign.)
d $3ab + 2ab = 5ab$ (Collecting the like terms on the left-hand side gives $5ab$ which is equal to the right-hand side.)
- 2** James is correct.
 $4x - 2 = 2x$ can be solved to find the value of x so it is an equation.
 Or, the two sides of $4x - 2 = 2x$ are not equal for all values of x so it cannot be an identity. For example, when $x = 2$:
 (Left-hand side) $4x - 2 = 4 \times 2 - 2 = 6$
 (Right-hand side) $2x = 2 \times 2 = 4$
 $6 \neq 4$

Simplifying expressions

Stretch it!

The expressions must all contain algebra, so each part must include t .

There are four possible combinations that make $12t^3$:
 $12t \times t \times t$, $2t \times 6t \times t$, $2t \times 3t \times 2t$, $3t \times 4t \times t$.

- 1 a** p^3
b $4 \times b \times c \times 7 = 4 \times 7 \times b \times c = 28bc$
c $4a \times 3b = 4 \times 3 \times a \times b = 12ab$
d $5x \times 4x = 5 \times 4 \times x \times x = 20x^2$
e $2g \times (-4g) = 2 \times (-4) \times g \times g = -8g^2$
f $2p \times 3q \times r = 2 \times 3 \times p \times q \times r = 6pqr$
- 2 a** $10x \div 2 = \frac{10x}{2} = 5x$
b $\frac{14w}{-2} = -7w$
c $6p \div p = \frac{6p}{p} = 6$
d $8mn \div 2m = \frac{8mn}{2m} = 4n$
e $\frac{12xy}{3y} = 4x$
f $9abc \div bc = \frac{9abc}{bc} = 9a$

Collecting like terms

- 1 a** $5f$
b $7b$
c $5mn$
d $4a + 6 - a - 5 = 4a - a + 6 - 5 = 3a + 1$
e $3d + 4e + d - 6e = 3d + d + 4e - 6e = 4d - 2e$
f $2x + 5y + 3x - 2y - 2 = 2x + 3x + 5y - 2y - 2$
 $= 5x + 3y - 2$
g $3a - 2b + 4a + 7b = 3a + 4a - 2b + 7b = 7a + 5b$
h $2a - b - 5a - 3 = 2a - 5a - b - 3 = -3a - b - 3$
i $x^2 + x^2 = 2 \times x^2 = 2x^2$
j $2t^3 + 4 - t^3 - 4 = 2t^3 - t^3 + 4 - 4 = t^3$
k $2a + b^2$
l $(4 + 3)\sqrt{x} = 7\sqrt{x}$
m $(7 - 4)\sqrt{x} = 3\sqrt{x}$
n $(12 - 1 - 4)\sqrt{x} = 7\sqrt{x}$

Using indices

- 1 a** $x^5 \times x^4 = x^{5+4} = x^9$
b $p \times p^4 = p^{1+4} = p^5$
c $2m^4 \times 3m^4 = 2 \times 3 \times m^4 \times m^4 = 6 \times m^{4+4} = 6m^8$
d $3m^4n \times 5m^2n^3$
 $= 3 \times 5 \times m^4 \times m^2 \times n \times n^3$
 $= 15 \times m^{4+2} \times n^{1+3} = 15m^6n^4$
e $u^{-2} \times u^5 = u^{-2+5} = u^3$
f $t^7 \times t^{-6} = t^{7+(-6)} = t$
- 2 a** $x^4 \div x^2 = x^{4-2} = x^2$
b $\frac{y^7}{y^3} = y^{7-3} = y^4$
c $\frac{p^9}{p^8} = p^{9-8} = p$
d $8x^6 \div 4x^3 = \frac{8x^6}{4x^3}$
 $(8 \div 4) \times (x^6 \div x^3) = 2 \times x^{6-3} = 2x^3$
e $m^3 \div m^5 = m^{3-5} = m^{-2} = \frac{1}{m^2}$
f $\frac{5x^8}{15x^4} = \frac{5}{15} \times \frac{x^8}{x^4} = \frac{1}{3} \times x^{8-4} = \frac{x^4}{3}$
g $3x^2 \div 9x = \frac{3x^2}{9x} = \frac{3}{9} \times \frac{x^2}{x} = \frac{1}{3} \times x^{2-1} = \frac{x}{3}$

- 3 a $(x^2)^3 = x^{2 \times 3} = x^6$
 b $(y^4)^4 = y^{4 \times 4} = y^{16}$
 c $(p^5)^2 = p^{5 \times 2} = p^{10}$
 d $(4m^5)^2 = 4^2 \times (m^5)^2 \times 16 \times m^{5 \times 2} = 16m^{10}$
 e $(x^2)^{-3} = x^{2 \times (-3)} = x^{-6} = \frac{1}{x^6}$
 f $(n^{-4})^{-2} = n^{-4 \times (-2)} = n^8$
- 4 $\frac{x^3 \times x^5}{x^4} = \frac{x^{3+5}}{x^4} = \frac{x^8}{x^4} = x^{8-4} = x^4$

Expanding brackets

Stretch it!

- a $a\sqrt{3} + a^2$ or $\sqrt{3}a + a^2$
 b $b\sqrt{5} - b^2$ or $\sqrt{5}b - b^2$
 c $c + d$

Stretch it!

- 1 $2 + 4 = 6$ and $2 \times 4 = 8$, so the numbers are 4 and 8.
 $(x+2)(x+4) = x^2 + 6x + 8$
- 2 a $(x + 3)(2x + 2)$
 $= 2x^2 + 6x + 2x + 6$
 $= 2x^2 + 8x + 6$
- b $(3x - 2)(x + 4)$
 $= 3x^2 - 2x + 12x - 8$
 $= 3x^2 + 10x - 8$
- c $(2x + 3)(3x - 1)$
 $= 6x^2 + 9x - 2x - 3$
 $= 6x^2 + 7x - 3$
- d $(2x - y)(3x + y)$
 $= 6x^2 + 2xy - 3xy - y^2$
 $= 6x^2 - xy - y^2$
- 1 a $3a + 6$ b $4b - 16$ c $10c + 25$
 d $6 - 2e$ e $4x + 4y + 8$ f $-2y - 4$
 g $x^2 - 2x$ h $2a^2 + 10a$
- 2 a $6a - (3a + 5)$
 $= 6a - 3a - 5$
 $= 3a - 5$
- b $4x - 6 + 2(x + 5)$
 $= 4x - 6 + 2x + 10$
 $= 4x + 2x - 6 + 10$
 $= 6x + 4$
- 3 a $2(2x + 3) + 4(x + 5) = 4x + 6 + 4x + 20 = 8x + 26$
 b $3(3y + 1) + 2(4y - 3) = 9y + 3 + 8y - 6 = 17y - 3$
 c $4(2m + 4) - 3(2m - 5) = 8m + 16 - 6m + 15 = 2m + 31$
- 4 a $(x + 2)(x + 3) = x^2 + 3x + 2x + 6 = x^2 + 5x + 6$
 b $(y - 3)(y + 4) = y^2 + 4y - 3y - 12 = y^2 + y - 12$
 c $(a + 3)(a - 7) = a^2 - 7a + 3a - 21 = a^2 - 4a - 21$
 d $(m - 1)(m - 6) = m^2 - 6m - m + 6 = m^2 - 7m + 6$
- 5 a $(x + 1)^2 = (x + 1)(x + 1)$
 $= x^2 + x + x + 1 = x^2 + 2x + 1$
- b $(x - 1)^2 = (x - 1)(x - 1)$
 $= x^2 - x - x + 1 = x^2 - 2x + 1$
- c $(m - 2)^2 = (m - 2)(m - 2)$
 $= m^2 - 2m - 2m + 4 = m^2 - 4m + 4$
- d $(y + 3)^2 = (y + 3)(y + 3)$
 $= y^2 + 3y + 3y + 9 = y^2 + 6y + 9$

Factorising

Stretch it!

The width of the rectangle = $x + 1$, since $x^2 + 3x + 2 = (x + 2)(x + 1)$

- 1 a $3(a + 3)$ b $5(b - 2)$
 c $7(1 + 2c)$ d $d(d - 2)$
- 2 a $4(2a + 5)$ b $4(b - 3)$
 c $9(2 + c)$ d $d(2d - 3)$
- 3 a $2(2x - 3y)$ b $m(a + b)$
 c $4a(3a + 2)$ d $x(4x + 3y)$
 e $n(2 - 9n)$ f $5x(1 + 2y)$
 g $4p(q - 3)$ h $4y(x^2 - 2)$
- 4 $4(x - 3) + 3(2x + 6)$
 $= 4x - 12 + 6x + 18$
 $= 4x + 6x - 12 + 18$
 $= 10x + 6$
 $= 2(5x + 3)$
 Compare $2(5x + 3)$ with $a(5x + b)$
 $a = 2, b = 3$
- 5 a $(x + 1)(x + 7)$ b $(x - 1)(x + 5)$
 c $(x + 2)(x - 4)$ d $(x - 2)(x - 3)$
 e $(x - 3)(x - 3)$ f $(x + 3)(x + 4)$
 g $(x - 2)(x + 5)$ h $(x + 4)(x - 5)$
- 6 a $x^2 - 16 = x^2 - 4^2 = (x + 4)(x - 4)$
 b $x^2 - 36 = x^2 - 6^2 = (x + 6)(x - 6)$
 c $x^2 - 81 = x^2 - 9^2 = (x + 9)(x - 9)$
 d $y^2 - 100 = y^2 - 10^2 = (y + 10)(y - 10)$

Substituting into expressions

- 1 When $a = 3$ and $b = -2$,
 $5a + 2b = 5 \times 3 + 2 \times (-2) = 15 + (-4) = 11$
- 2 a $2 - 2 \times (-4) = 2 - (-8) = 10$
 b $3 \times 2 \times (-4) = -24$
 c $4 \times (-4) - 3 \times 2 = -16 - 6 = -22$
 d $2^2 + (-4)^2 = 4 + 16 = 20$
 e $2 \times 2 + 4(2 - (-4)) = 2 \times 2 + 4 \times 6 = 4 + 24 = 28$
 f $\frac{1}{2}(2 + (-4)) = \frac{1}{2} \times -2 = -1$
- 3 false
 When $a = 3$: $3a^2 = 3 \times 3^2 = 3 \times 9 = 27$
- 4 When $p = \frac{1}{2}$ and $q = -4$,
 a $10pq = 10 \times \frac{1}{2} \times (-4) = -20$
 b $8p^2 = 8 \times \left(\frac{1}{2}\right)^2 = 8 \times \frac{1}{4} = 2$
 c $\frac{q}{p} = \frac{-4}{\frac{1}{2}} = -4 \times 2 = -8$
 d $2q^2 - 12p = 2 \times (-4)^2 - 12 \times \frac{1}{2}$
 $= 2 \times 16 - 12 \times \frac{1}{2}$
 $= 32 - 6$
 $= 26$

- c** $4x + 3 = 2x + 8$
 $2x + 3 = 8$
 $2x = 5$
 $x = \frac{5}{2}$
 (Or $x = 2.5$ or $x = 2\frac{1}{2}$)
- d** $3 - 2p = 6 - 3p$
 $3 + p = 6$
 $p = 3$
- e** $3y - 8 = 5y + 4$
 $-8 = 2y + 4$
 $-12 = 2y$
 $-6 = y$
- 7 a** $2(x + 5) = x + 6$
 $2x + 10 = x + 6$
 $x + 10 = 6$
 $x = -4$
- b** $7b - 2 = 2(b + 4)$
 $7b - 2 = 2b + 8$
 $5b - 2 = 8$
 $5b = 10$
 $b = 2$
- c** $4(2y + 1) = 3(5y - 1)$
 $8y + 4 = 15y - 3$
 $4 = 7y - 3$
 $7 = 7y$
 $1 = y$
- d** $2x - 1 = 8 - 4x$
 $6x - 1 = 8$
 $6x = 9$
 $x = \frac{9}{6} = \frac{3}{2}$
 (Or $x = 1.5$ or $x = 1\frac{1}{2}$)

Writing linear equations





- 1 a** Perimeter = $4 \times (2s + 3) = 8s + 12$
 (Or, Perimeter = $2s + 3 + 2s + 3 + 2s + 3 + 2s + 3 = 8s + 12$)
- b** $8s + 12 = 84$
 $8s = 72$
 $s = 9$ cm
- 2 a** Angles in a quadrilateral add up to 360° so:
 $x + 20 + 2x - 15 + x + 65 + 2x - 10 = 360$
 $6x + 60 = 360$
 $6x = 300$
 $x = 50$
- b** Largest angle: $x + 65 = 50 + 65 = 115^\circ$
 (Other angles: $x + 20 = 50 + 20 = 70^\circ$;
 $2x - 15 = 2 \times 50 - 15 = 85^\circ$;
 $2x - 10 = 2 \times 50 - 10 = 90^\circ$)
- 3** Let $a =$ Karen's age
 Monica is 4 years younger: $a - 4$
 $a + a - 4 = 64$
 $2a - 4 = 64$
 $2a = 68$
 $a = 34$

Karen is 34 years old.
 $a - 4 = 34 - 4 = 30$
 Monica is 30 years old.


- 4** Let $n =$ number.
 $2n + 4 = 16 - n$
 $3n + 4 = 16$
 $3n = 12$
 $n = 4$
 The number is 4.
- 5** Let $l =$ length of rectangle.
 Width is 2 cm smaller: $l - 2$
 Perimeter = $2l + 2(l - 2)$
 $= 2l + 2l - 4 = 4l - 4$
 $4l - 4 = 36$
 $4l = 40$
 $l = 10$
 Length is 10 cm.
 $l - 2 = 10 - 2 = 8$
 Width is 8 cm.
- 6** Base angles of an isosceles triangle are equal so:
 $4a - 20 = 2a + 16$
 $2a - 20 = 16$
 $2a = 36$
 $a = 18$
 When $a = 18$: $4a - 20 = 4 \times 18 - 20 = 52$
 So $2a + 16 = 52$
 Angles in a triangle add up to 180° so:
 $4b - 2a + 52 + 52 = 180$
 $4b - 2a + 104 = 180$
 $4b - 2a = 76$ (Substitute $a = 18$)
 $4b - 2 \times 18 = 76$
 $4b - 36 = 76$
 $4b = 112$
 $b = 28$

Linear inequalities

- 1 a** $x = 3, 4, 5$
b $x = 2, 3, 4, 5$
c $x = 0, 1, 2, 3$
d $x = -3, -2, -1, 0, 1$
- 2 a** $x < 3$ **b** $x \geq -2$
c $-1 \leq x \leq 5$




- 4 a** $2x - 2 > 4$
 $2x > 6$
 $x > 3$



b $2x - 7 \leq 3$

$2x \leq 10$

$x \leq 5$




c $4x + 3 \leq 13$

$4x \leq 10$

$x \leq \frac{10}{4} = \frac{5}{2}$


(Or $x \leq 2.5$ or $x \leq 2\frac{1}{2}$)



d $4x < 2x - 10$

$2x < -10$

$x < -5$



e $4x - 8 < 6x$

$4x < 6x + 8$

$-2x < 8$


$x > -4$

Alternative method:

$4x - 8 < 6x$

$-8 < 2x$

$-4 < x$



f $7x + 2 \geq 3x - 2$

$4x + 2 \geq -2$

$4x \geq -4$

$x \geq -1$



5 a $-12 < 4x \leq 8$

$-3 < x \leq 2$

$x = -2, -1, 0, 1, 2$

b $-8 \leq 2x < 14$

$-4 \leq x < 7$

$x = -4, -3, -2, -1, 0, 1, 2, 3, 4, 5, 6$

c $-6 < 6x \leq 18$

$-1 < x \leq 3$

$x = 0, 1, 2, 3$

d $9 \leq 3n \leq 15$

$3 \leq n \leq 5$

$n = 3, 4, 5$

6 a $4 - x \leq 1$

$4 \leq x + 1$

$3 \leq x$ (or $x \geq 3$)

Alternative method:

$4 - x \leq 1$

$-x \leq -3$

$x \geq 3$

b $6 - 3x > 9$

$6 > 3x + 9$

$-3 > 3x$

$-1 > x$ (or $x < -1$)

Alternative method:

$6 - 3x > 9$

$-3x > 3$

$x < -1$

c $8 - 2x \geq 7$

$8 \geq 2x + 7$

$1 \geq 2x$

$\frac{1}{2} \geq x$ (or $x \leq \frac{1}{2}$)

Alternative method:

$8 - 2x \geq 7$

$-2x \geq -1$

$x \leq \frac{1}{2}$

d $-2 < -x \leq 3$

$2 > x \geq -3$

(or $-3 \leq x < 2$)

Formulae

1 wage earned = $8 \times 35 + 25 = 280 + 25 = 305$
 £305

2 $F = \frac{9}{5} \times 45 + 32$
 $= 81 + 32 = 113$
 113°F

3 $v = 10 + (-20) \times 5 = 10 + (-100) = -90$

4 $C = 25d + 50$

5 a Distance in kilometres = $\frac{8}{5} \times$ distance in miles

$k = \frac{8}{5}m$

b $k = \frac{8}{5} \times 200$
 $= 320$ km

6 a $P = 2a + 2(a + 3) = 2a + 2a + 6 = 4a + 6$

(or $P = a + a + a + 3 + a + 3 = 4a + 6$)

b $P = 4 \times 6 + 6 = 24 + 6 = 30$

$P = 30$ cm

7 $-10 = \frac{D}{6.5}$
 $-65 = D$

8 a $v = u + at$

$v - u = at$

$\frac{v - u}{t} = a$

b $V = \frac{1}{3}Ah$

$3V = Ah$

$\frac{3V}{A} = h$

c $y = 3(x - 3)$

$y = 3x - 9$

$y + 9 = 3x$

$\frac{y + 9}{3} = x$

(or $x = \frac{y}{3} + 3$)

d $v^2 = u^2 + 2as$

$v^2 - u^2 = 2as$

$\frac{v^2 - u^2}{2a} = s$

e $T = \sqrt{\frac{2s}{g}}$

$T^2 = \frac{2s}{g}$

$gT^2 = 2s$

$g = \frac{2s}{T^2}$

Linear sequences

- 1 a The term-to-term rule is add 6, so the next two terms are:

$$21 + 6 = 27$$

$$27 + 6 = 33$$

b $9 = 3 + 6$

$$15 = 9 + 6$$

$$21 = 15 + 6$$

The term-to-term rule is 'add 6'.

2 a 1st term = $1 \times 4 - 2 = 2$

2nd term = $2 \times 4 - 2 = 6$

3rd term = $3 \times 4 - 2 = 10$

4th term = $4 \times 4 - 2 = 14$

b 20th term = $20 \times 4 - 2 = 78$

3 a

Pattern number	1	2	3	4	5
Number of sticks	7	10	13	16	19

- b Three new sticks are added each time, so the term-to-term rule is add 3.

Pattern 5 has 19 sticks, so:

$$\begin{aligned} \text{pattern 7 has } & 19 + (7 - 5) \times 3 = 19 + 2 \times 3 \\ & = 25 \text{ sticks} \end{aligned}$$

4 a 1st term = $2 \times 1 + 3 = 5$

2nd term = $2 \times 2 + 3 = 7$

3rd term = $2 \times 3 + 3 = 9$

5, 7, 9

b 100th term = $2 \times 100 + 3 = 203$

c $2n + 3 = 73$

$$2n = 70$$

$$n = 35$$

35th term has a value of 73.

- 5 a Common difference is 7. Hence the term-to-term rule is add 7, so:

$$-11 + 7 = -4$$

$$-4 + 7 = 3$$

$$3 + 7 = 10$$

3, 10

b $15 - 2n < 0$

$$-2n < -15$$

$$-n < -7.5$$

$$n > 7.5$$

n is an integer.

8th term:

$$15 - 2 \times 8 = -1$$

Compare 7th term:

$$15 - 2 \times 7 = 1$$

So 8th term is the first term with a negative value.

6 a 3, 7, 11, 15

Common difference = +4

$4 \times$ term number = 4, 8, 12, 16

- 1 to get each term in the original sequence

So, n th term is $4n - 1$.

- b If $4n - 1 = 100$, then:

$$4n = 101$$

$$n = 25.25$$

But 25.25 is not an integer, so 100 cannot be a term in the sequence.

Non-linear sequences

- 1 In a geometric progression there is a multiplier - each term is multiplied by the same amount to get the next one.

In b:

$$2 = 1 \times 2$$

$$4 = 2 \times 2$$

$$8 = 4 \times 2$$

$$16 = 8 \times 2$$

There is a multiplier (2), so b is a geometric progression.

None of the other sequences has a multiplier (a is arithmetic, c is a Fibonacci-type sequence, and d is quadratic).

2

$$3 \quad 6 \quad 11 \quad 18 \quad 27 \quad 38$$

$$27, 38$$

3 2nd term = $0.5 = 5 \div 10$

3rd term = $0.05 = 0.5 \div 10$

4th term = $0.005 = 0.05 \div 10$

So the term-to-term rule is divide by 10.

5th term = $0.005 \div 10 = 0.0005$

4 4th term = 6

5th term = 10

6th term = $6 + 10 = 16$

7th term = $10 + 16 = 26$

8th term = $16 + 26 = 42$

5 1st term = $1^2 + 5 = 6$

2nd term = $2^2 + 5 = 9$

3rd term = $3^2 + 5 = 14$

4th term = $4^2 + 5 = 21$

6, 9, 14, 21

Show that...

Stretch it! If n is even, $n - 1$ is odd and $n + 1$ is odd. If you multiply two odd numbers the answer will always be odd.

If m is odd, $m - 1$ is even and $m + 1$ is even. If you multiply two even numbers the answer will always be even.

or: $(n + 1)(n - 1) = n^2 - 1$ and $(m + 1)(m - 1) = m^2 - 1$

n^2 will be even \times even = even, so $n^2 - 1$ will be odd.

m^2 will be odd \times odd = odd, so $m^2 - 1$ will be even.

1 a LHS = $4(x - 3) + 2(x + 5) = 4x - 12 + 2x + 10 = 6x - 2$

RHS = $3(2x - 1) + 1 = 6x - 3 + 1 = 6x - 2$

LHS = RHS

So $4(x - 3) + 2(x + 5) \equiv 3(2x - 1) + 1$

b LHS = $(x + 2)(x - 2) = x^2 - 2x + 2x - 4 = x^2 - 4$

LHS = RHS

So $(x + 2)(x - 2) = x^2 - 4$

Number machines

- 1 a $y = 3x - 3$
 b $10 \times 3 - 3 = 30 - 3 = 27$
 c $y = 3x - 3$
 $y + 3 = 3x$
 $x = \frac{y+3}{3}$
 d If $x = y$,
 $3x - 3 = x$
 $2x - 3 = 0$
 $2x = 3$
 $x = \frac{3}{2}$
 (or $x = 1\frac{1}{2}$)

Coordinates and midpoints

Stretch it!

Difference in x coordinates of A and B is equal to difference in x coordinates of B and C .

Difference = $2 - (-1) = 3$

So x coordinate of C is:

$-1 - 3 = -4$

Difference in y coordinates of A and B is equal to difference in y coordinates of B and C .


Difference = $4 - 3 = 1$

So y coordinate of C is:

$3 - 1 = 2$

C has coordinates $(-4, 2)$.

- 1 a $(2, 2)$
 b, c and d




- d B is $(-4, 2)$, D is $(2, -3)$
 x coordinate of midpoint: $\frac{2 + (-4)}{2} = -1$
 y coordinate of midpoint: $\frac{-3 + 2}{2} = -0.5$
 Midpoint is $(-1, -0.5)$

Straight-line graphs

Stretch it!


To solve the equation, you need to find where the graph of $y = 3x - 2$ intersects the graph of $y = -x$.



So the solution to $3x - 2 = -x$ is $x = 0.5$.



1

x	-2	-1	0	1	2
y	4	3	2	1	0



2

x	-2	0	2
y	-7	-3	1



$$\text{Gradient} = \frac{\text{difference in } y \text{ coordinate}}{\text{difference in } x \text{ coordinate}} = \frac{5}{5} = 1$$

Gradient, $m = 1$

y -intercept, $c = 1$

Using the general form of the equation of a line
 $y = mx + c$: the equation of the line is $y = x + 1$

4 Gradient, $m = 2$

$y = mx + c$ (General form of the equation of a line)

$$y = 2x + c$$

For point $(1, -2)$, $x = 1$, $y = -2$:

$$-2 = 2 \times 1 + c$$

$$-2 = 2 + c$$

$$-4 = c$$

So the equation of the line is $y = 2x - 4$

5 Gradient = $\frac{\text{difference in } y \text{ coordinate}}{\text{difference in } x \text{ coordinate}} = \frac{4 - 2}{0 - 4} = \frac{-2}{-4} = \frac{1}{2}$

Gradient, $m = \frac{1}{2}$

Given the point $(0, 4)$:


$$4 = \frac{1}{2}(0) + c$$

$$4 = c \text{ (} y\text{-intercept)}$$

So the equation of the line is $y = \frac{1}{2}x + 4$.


6 a

x	0	6
y	6	0



b

x	0	4
y	2	0



7 First line:

$$4x + 4y = 8$$

$$4y = 8 - 4x$$

$$y = 2 - x$$

$$y = -x + 2$$

Second line:

$$2x = 6 - 2y$$

$$2y + 2x = 6$$

$$2y = 6 - 2x$$

$$y = 3 - x$$

$$y = -x + 3$$

With both lines in the form, $y = mx + c$, you can see that they both have gradient $m = -1$, so they are parallel.

8 $y + 2x = 4$

$$y = 4 - 2x$$


$$y = -2x + 4$$

Compare with $y = mx + c$:

gradient, $m = -2$

y -intercept $c = 4$

Coordinates of the y -intercept are $(0, 4)$.



- a $y = 3.4$ b $x = 0.5$
 c $x \approx 2.5$ d $x \approx -0.5$

Solving simultaneous equations

1 $x + y = 16$ (1)

$x - y = 5$ (2)

(1) + (2):

$$2x = 21$$

$$x = 10.5$$

Substitute $x = 10.5$ into 2:

$$10.5 + y = 5$$

$$y = 5 - 10.5$$

$$y = -5.5$$

Solution: $x = 10.5, y = -5.5$

2 a $2x + y = 4$ (1)

$3x - y = 1$ (2)

(1) + (2): $5x = 5$

$$x = 1$$

Substitute $x = 1$ in (1)

$$2 \times 1 + y = 4$$

$$2 + y = 4$$

$$y = 2$$

Solution: $x = 1, y = 2$

b $x - y = 5$ (1)

$2x + y = 4$ (2)

(1) + (2): $3x = 9$

$$x = 3$$

Substitute $x = 3$ into (1)

$$3 - y = 5$$

$$3 = y + 5$$

$$y = -2$$

Solution: $x = 3, y = -2$

c $2x + y = 8$ (1)
 $x + y = 2$ (2)
 (1) - (2): $x = 6$
 Substitute $x = 6$ into (2)
 $6 + y = 2$
 $y = -4$
 Solution: $x = 6, y = -4$

d $4x - y = 10$ (1)
 $x + 2y = 7$ (2)
 (1) $\times 2$: $8x - 2y = 20$ (3)
 (2) + (3): $9x = 27$
 $x = 3$
 Substitute $x = 3$ into (2):
 $3 + 2y = 7$
 $2y = 4$
 $y = 2$
 Solution: $x = 3, y = 2$

e $2x + y = 7$ (1)
 $x - 4y = 8$ (2)
 (1) $\times 4$: $8x + 4y = 28$ (3)
 (2) + (3): $9x = 36$
 $x = 4$
 Substitute $x = 4$ into (1):
 $2 \times 4 + y = 7$
 $8 + y = 7$
 $y = -1$
 Solution: $x = 4, y = -1$

f $2x + 3y = 7$ (1)
 $3x - 2y = 4$ (2)
 (1) $\times 2$: $4x + 6y = 14$ (3)
 (2) $\times 3$: $9x - 6y = 12$ (4)
 (3) + (4): $13x = 26$
 $x = 2$
 Substitute $x = 2$ into (1)
 $2 \times 2 + 3y = 7$
 $4 + 3y = 7$
 $3y = 3$
 $y = 1$
 Solution: $x = 2, y = 1$


3 $x + y = 21$ (1)
 $x - y = 7$ (2)
 (1) + (2): $2x = 28$
 $x = 14$
 Substitute $x = 14$ into (1)
 $14 + y = 21$
 $y = 7$

The two numbers are 7 and 14.

4 Let b = burger and c = cola.
 $3b + 2c = 505$ (1)
 $3b + 4c = 725$ (2)
 (2) - (1): $2c = 220$
 $c = 110$
 Substitute $c = 110$ into (1)


$3b + 2 \times 110 = 505$
 $3b + 220 = 505$
 $3b = 285$
 $b = 95$
 A burger costs 95p.
 A cola costs £1.10.

5 a



$x = 2, y = 8$

b




$x = 8, y = 1$

Quadratic graphs

Stretch it!

Rearrange $x^2 - 3x = 3$, to give $x^2 - 3x - 2 = 1$


You can solve this graphically by finding where the lines $y = x^2 - 3x - 2$ and $y = 1$ intersect.



So the solutions to the equation $x^2 - 3x = 3$ are $x = 3.8$ and $x = -0.8$. Acceptable readings from the graph would be in the range 3.6 to 3.9 and -0.6 to -0.9 .

1 a


x	-2	-1	0	1	2	3	4
y	6	1	-2	-3	-2	1	6



- b $x = 1$
 c $(1, -3)$

2


x	-2	-1	0	1	2	3	4
y	8	3	0	-1	0	3	8



- b The roots are where the curve cuts the x axis (the line $y = 0$). These are where $x = 0$ and $x = 2$.

3 a

x	-3	-2	-1	0	1	2	3
y	2	-1	-2	-1	2	7	14



Read off the values of x where the graph cuts the line $y = 0$ (the x -axis).

$x \approx -2.4$ and $x \approx 0.4$

Solving quadratic equations

Stretch it!

$$\frac{x^2}{2} = 8$$

$$x^2 = 16$$

$$x = \sqrt{16}$$

$$\text{So } x = 4 \text{ or } x = -4$$

$$2x^2 = 50$$

$$x^2 = 25$$

$$x = \sqrt{25}$$

$$\text{So } x = 5 \text{ or } x = -5$$

1 a $x^2 - 4x = 0$

$$x(x - 4) = 0$$

$$\text{Either } x = 0 \text{ or } x - 4 = 0$$

$$x = 4$$

$$\text{So } x = 0 \text{ or } x = 4$$

b $x^2 + 7x = 0$

$$x(x + 7) = 0$$

$$\text{Either } x = 0 \text{ or } x + 7 = 0$$

$$x = -7$$

$$\text{So } x = 0 \text{ or } x = -7$$

c $x^2 - 16 = 0$ ($x^2 - 16 = x^2 - 4^2$, Factorise)

$$(x + 4)(x - 4) = 0$$

$$\text{Either } x + 4 = 0 \text{ or } x - 4 = 0$$

$$x = -4 \quad x = 4$$

$$\text{So } x = -4 \text{ or } x = 4$$

d $x^2 + 10x + 9 = 0$

$$(x + 1)(x + 9) = 0$$

$$\text{Either } x + 1 = 0 \text{ or } x + 9 = 0$$

$$x = -1 \quad x = -9$$

$$\text{So } x = -1 \text{ or } x = -9$$

e $x^2 + x - 12 = 0$

$$(x - 3)(x + 4) = 0$$

$$\text{Either } x - 3 = 0 \text{ or } x + 4 = 0$$

$$x = 3 \quad x = -4$$

$$\text{So } x = 3 \text{ or } x = -4$$

f $x^2 - 6x - 16 = 0$

$$(x + 2)(x - 8) = 0$$

$$\text{Either } x + 2 = 0 \text{ or } x - 8 = 0$$

$$x = -2 \quad x = 8$$

$$\text{So } x = -2 \text{ or } x = 8$$

2 a $y = x^2 - 49$ (Set $y = 0$)

$$x^2 - 49 = 0$$
 ($x^2 - 49 = x^2 - 7^2$, Factorise)

$$(x + 7)(x - 7) = 0$$

$$\text{Either } x + 7 = 0 \text{ or } x - 7 = 0$$

$$x = -7 \quad x = 7$$

$$\text{So } x = -7 \text{ or } x = 7$$

b $y = x^2 - 3x$ (Set $y = 0$)

$$x^2 - 3x = 0$$

$$x(x - 3) = 0$$

$$\text{Either } x = 0 \text{ or } x - 3 = 0$$

$$x = 3$$


$$\text{So } x = 0 \text{ or } x = 3$$

- c $y = x^2 + 7x + 6$ (Set $y = 0$)
 $x^2 + 7x + 6 = 0$
 $(x + 1)(x + 6) = 0$
 Either $x + 1 = 0$ or $x + 6 = 0$
 $x = -1$ $x = -6$
 So $x = -1$ or $x = -6$

Cubic and reciprocal graphs


1 a

x	-3	-2	-1	0	1	2	3
y	-28	-9	-2	-1	0	7	26




2 a

x	-4	-3	-2	-1	$-\frac{1}{2}$	$-\frac{1}{4}$	$\frac{1}{4}$	$\frac{1}{2}$	1	2	3	4
y	$\frac{1}{4}$	$\frac{1}{3}$	$\frac{1}{2}$	1	2	4	-4	-2	-1	$-\frac{1}{2}$	$-\frac{1}{3}$	$-\frac{1}{4}$



Drawing and interpreting real-life graphs




- b See lines drawn on graph.
 i €36 ii £75

- c From the graph: £30 = €36
 So £90 = €36 × 3 = €108
 The ring is cheaper in France.
- 2 a Monthly charge = £10 (cost of 0 minutes from the graph)
 b Gradient = $\frac{30}{240} = 0.125$
 Charge per minute of calls is 13p.


3 a and c

d	0	1	2	3	4
C	20	32	44	56	68




- b This is the flat rate that you pay just for hiring the sander, before you pay for the number of days. It is the intercept with the vertical axis:
 days (d) = 0
 cost (C) = £20
- c Using a graphical method: plot the second equation, $C = 10d + 30$, on the same axes. The line for the second shop has a lower gradient, and after the lines cross over (at $d = 5$), the second shop is cheaper. So you would use the second shop.
 Alternatively, using an algebraic method:
 Let $d = 6$ days (more than 5 days)
 First shop: $C = 12 \times 6 + 20 = 92$
 Second shop: $C = 10 \times 6 + 30 = 90$
 To hire the sander for more than 5 days use the second shop as it is cheaper.

- 4 a 30 minutes (Horizontal line on graph)
 b 55 km
 c Speed before break = $\frac{\text{distance (km)}}{\text{time (hours)}} = \frac{30}{1} = 30$ km/hr
 Speed after break = $\frac{\text{distance (km)}}{\text{time (hours)}} = \frac{25}{1.5} = 16.7$ km/hr




- 5 a Reading off maximum height value from graph: **12.8 m**.
 b Reading from the graph, the ball is thrown at time = 0 seconds and returns to the ground at time = **3 seconds**.
 c Draw a horizontal line on the graph at height = 8 m.



0.4 seconds and 2.4* seconds

- d The ball is thrown from a height of 3 m above the ground.
- 6 a




b Acceleration = $\frac{\text{change in speed (m/s)}}{\text{time (s)}} = \frac{20-10}{30} = \frac{10}{30} = 0.3 \text{ m/s}^2$

- 7 a The maximum depth of water in the bath before the person got in was 35 cm.
 b Between C and D, the person was taking their bath.
 c Between D and E, the person got out of the bath.
 d Running water into the bath was quicker. The slope of the line between O and A (filling the bath) is steeper than the slope of the line between E and F (emptying the bath).

Review it!

- 1 a P(3, 5)
 b, c and e




- d Q(3, -2), S(-4, 5)
 x-coordinate: $-4 + 3 = -1$
 $-1 \div 2 = -0.5$
 y-coordinate: $5 + (-2) = 3$
 $3 \div 2 = 1.5$
 Midpoint is (-0.5, 1.5)

- 2 a $2x + 8 = 4$
 $2x = -4$
 $x = -2$
 b When $x = 2$ and $y = -4$
 A: $\frac{y}{x} = \frac{-4}{2} = -2$
 B: $x - y = 2 - (-4) = 6$
 C: $xy = 2 \times -4 = -8$
 Expression C has the smallest value.
 c Millie is correct.
 When $x = 4$, $3x^2 = 3 \times 4^2 = 3 \times 16 = 48$
 (George has worked out $(3x)^2$ instead.)
 3 a $7a - (3a + 4) = 7a - 3a - 4 = 4a - 4$
 b $4(2x + 3)$
 c $m^4 \times m = m^{4+1} = m^5$
 d $\frac{x^8}{x^3} = x^{8-3} = x^5$

4 a

x	-1	0	1	2	3
y	-1	1	3	5	7



- c Compare $y = 2x + 1$ with $y = mx + c$ (general form of the equation of a line):
 Gradient, $m = 2$
 5 $4x + 4 = x + 13$
 $3x + 4 = 13$
 $3x = 9$
 $x = 3$
 6 a 2 is included, and so are all values lower than 2.
 $x \leq 2$
 b
 c 1 can be included, but 4 cannot.
 $x = 1, 2, 3$
 d $4x + 2 \leq 2x + 5$
 $2x + 2 \leq 5$
 $2x \leq 3$
 $x \leq \frac{3}{2}$
 (or $x \leq 1\frac{1}{2}$)

7 a $6x$ or $x + 65$

b $6x = x + 65$

$5x = 65$

$x = 13$

Luke is 13 years old.

8 a The term-to-term rule is 'add 6'.

$21 + 6 = 27$

$27 + 6 = 33$

b No. The n th term is $6n - 3$.

$6n = 2 \times 3n =$ always even

Because 3 (odd) is always taken away from $6n$, every term in the sequence will be odd. As 44 is even it is not in the sequence.

c When $n = 5$:

$2n^2 - 3 = 2 \times 5^2 - 3 = 2 \times 25 - 3 = 47$

9 $(x + 3)(x + 4) = x^2 + 4x + 3x + 12 = x^2 + 7x + 12$

10 Smallest value of $a - b$ is where a is as small as possible and b is as large as possible.

$a > 30$ so its smallest value is 31

$b < 20$ so its largest value is 19

Using $a = 31$ and $b = 19$:

$a - b = 31 - 19 = 12$

11 The opposite sides of a rectangle are equal in length so:

$5x - 8 = 2x + 4$

$3x - 8 = 4$

$3x = 12$

$x = 4$

$14 - 2y = 4y + 2$

$14 = 6y + 2$

$12 = 6y$

$2 = y$

12 $3(ax - 4) + 2(4x + b) \equiv 14x - 6$

$3ax - 12 + 8x + 2b \equiv 14x - 6$

$3ax + 8x - 12 + 2b \equiv 14x - 6$

$3ax + 8x \equiv 14x$

$3a + 8 = 14$

$3a = 6$

$a = 2$

$-12 + 2b \equiv -6$

$2b = 6$

$b = 3$

13 a $12m$

b $3p \times 4p = 3 \times 4 \times p \times p = 12p^2$

c $12x \div 2 = \frac{12x}{2} = 6x$

14 a $5(w - 4) = 35$

$5w - 20 = 35$

$5w = 55$

$w = 11$

b When $a = 7$ and $b = -2$,


$5a + 7b = 5 \times 7 + 7 \times (-2)$

$= 35 + (-14)$

$= 21$

c $5a + 8b$

15 a



P and Q are vertically above each other, because they share an x coordinate (3).

R and Q are horizontally aligned, because they share a y coordinate (-2).

As the fourth vertex, S must share an x coordinate with R (-5) and a y coordinate with P (4).

S is the point (-5, 4).

b Length is x coordinate of P or Q minus x coordinate of R or S :


$= 3 - -5 = 8$

Width is y coordinate of S or P minus y coordinate of R or Q :

$= 4 - -2 = 6$

Length is 8 units and width is 6 units.

16 a*



b i From the graph: 10 inches = 25 cm

ii From the graph: 50 cm = 20 inches

So 50 cm = $10 \times 2 = 20$ inches

c From the graph: 10 inches = 25 cm

So 60 inches = $25 \times 6 = 150$ cm

Cost of beading = $150 \times 2 = 300$ p

Cost = £3.00

17 a $T = 12.50x + 10$

b $72.50 = 12.50x + 10$

$62.50 = 12.50x$

$5 = x$

Suzanne hired the costume for 5 days.

*This answer differs from the one in the Revision Guide due to an error in our first edition. This answer has now been re-checked and corrected.

- 18 $P = \frac{Q}{4} + R$
 $P - R = \frac{Q}{4}$
 $4(P - R) = Q$
- 19 a $m(m + 8)$
 b $(x + 3)(x + 4)$
- 20 a 2, 5, 8, 11, 14
 Common difference = +3
 3 × term number: 3, 6, 9, 12, 15
 - 1 to get each term in the original sequence
 So n th term = $3n - 1$
- b 50th term = $3 \times 50 - 1 = 149$
- c $2n - 3 = 112$
 $2n = 115$
 $n = 57.5$
 No, Kadena is incorrect.
 112 cannot be term in the sequence because 57.5 is not an integer.
- 21 a $4(x + 5) - 3(2x - 1) = 4x + 20 - 6x + 3 = -2x + 23$
 b $4a^3b^2 \times 5a^2b = 4 \times 5 \times a^3 \times a^2 \times b^2 \times b$
 $= 20 \times a^{3+2} \times b^{2+1} = 20a^5b^3$
- 22 Perimeter = $3x - 2 + 2x + 1 + 3x + 5 + 2x = 10x + 4$
 $10x + 4 = 49$
 $10x = 45$
 $x = 4.5$
- 23 A: output = $6x - 4$
 B: output = $3x + 2$
 $6x - 4 = 4(3x + 2)$
 $6x - 4 = 12x + 8$
 $-4 = 6x + 8$
 $-12 = 6x$
 $-2 = x$
 Input = -2
- 24 1st term: $4 + 2a$
 2nd term: $4 + 4a$
 3rd term: $4 + 6a$
 4th term: $4 + 8a$
 5th term: $4 + 10a$
 $4 + 10a = 64$
 $10a = 60$
 $a = 6$
- 25 Roots 2 and -4 are x -intercepts where the curve cuts the x -axis.
 These give factors:
 $(x - 2)$ and $(x - (-4))$
 Equation is $(x - 2)(x + 4) = 0$
 Equation C.

Ratio, proportion and rates of change

Units of measure

- 1 a 3000 m
 b 75 mins
 c 13 000 cm²
 d 3.52 litres
 e 7200 seconds
 f 14 kg
- 2 $4.5 - 0.325 = 4.175$ kg or $4500 - 325 = 4175$ g
- 3 $5 \div 2.2 = 2.\dot{2}\dot{7}$ kg

Ratio

Stretch it! $31 + 25 = 56$, fraction male = $\frac{31}{56}$

- 1 a 1:4 b 1:3:4 c 4:5
- 2 $35:5 = 7:1$
- 3 $375 \div 250 = 1.5$.
 Allow 1 part cement for 1.5 parts sand.
- 4 a number in evening = 7 parts
 number in afternoon = 1 part
 7:1
 b total parts = $7 + 1 = 8$
 1 part = $800 \div 8 = 100$
 1 part sold in afternoon
 So 100 tickets were sold in the afternoon.
- 5 a There are 5 parts to the ratio. Ratio is:
 $3 : (5 - 3)$
 $= 3 : 2$
 b 1 part = $200 \div 5 = 40$
 $40 \times 3 = 120$
 120 cats
- 6 There are 5 parts to the ratio.
 1 part = $1.5 \div 5 = 0.3$ kg
 2 parts of flour needed:
 $2 \times 0.3 = 0.6$
 0.6 kg or 600 g
- 7 $9 = 3 \times 3$ so multiply the other lengths by 3.
 $4 \times 3 = 12$
 $5 \times 3 = 15$
 12 cm and 15 cm
- 8 To work out the number of students per teacher (s), you multiply the number of teachers (t) by 20, so: $s = 20t$.
- 9 $5 - 2 = 3$ parts = 60 g more flour
 $60 \div 3 = 20$
 $2 \times 20 = 40$
 40 g of sugar

Scale diagrams and maps

Stretch it! 50 miles on ground = $50 \div x$ or $\frac{50}{x}$ miles on map

1 mile = 1610 m = 161 000 cm

50 miles on ground = $\frac{50}{x} \times 161\,000$ cm* on map

- 1 A, B, F
- 2 a $3 \times 12 = 36$ km
 b $15 \div 12 = 1.25$ cm

- 3 $12 \times 1000 = 12\,000\text{ cm} = 120\text{ m}$
- 4 a $2\text{ cm}: 2 \times 50\,000 = 100\,000\text{ cm} = 1\text{ km}$
 (Any answer within the range of $1\text{ km} - 1.1\text{ km}$ is acceptable.)
- b 250°

Fractions, percentages and proportion

- 1 $\frac{20}{3500} = \frac{1}{175}$
- 2 $2 + 3 + 8 = 13\text{ hours}$
 $24 - 13 = 11\text{ hours}$
 $\frac{11}{24}$ of the day remaining
- 3 a $\frac{15}{20} = \frac{3}{4}$
- b $1 - \frac{3}{4} = \frac{1}{4} = 25\%$
- 4 $1 + 2 + 7 = 10, \frac{1}{10} = 10\%$
- 5 School A: $125:145 = 25:29$
 School B: $100:120 = 5:6$
 No since the ratios are not equivalent.
- 6 $150 \div 100 = 1.5$
 $1.5 \times 22\text{ g} = 33\text{ g}$
 $33\text{ g} \div 8 = 4.125\text{ g}$

Direct proportion

Stretch it!

For two values to be in direct proportion, when one is 0 the other must be 0. Here, when distance is 0 miles, the fee is £2.

- 1 A and E
- 2 a i 20 meringues = 2 eggs, divide both by 2 to give:
 10 meringues = 1 egg
 3 eggs: $3 \times 10 = 30$ meringues
- ii 20 meringues = 120 g of sugar, divide both by 2 to give:
 10 meringues = 60 g of sugar. Multiply both by 10 to give 100 meringues
- b 20 meringues = 2 eggs, divide both by 2 to give
 10 meringues = 1 egg, multiply both by 7 to give
 70 meringues = 7 eggs
- 3 $675 \div 4.5 = 150\text{ minutes} = 2\text{ hours } 30\text{ minutes}$
- 4 A, D

Inverse proportion

- 1 D
- 2 At 60 miles it takes 15 minutes.
 $60 \times \frac{2}{3} = 40$
 $15 \div \frac{2}{3} = 22.5\text{ mins}$
- 3 $2 \times 3 = 6$ decorators
 $5 \div 3 = 1\frac{2}{3}$ of a day
- 4 a 2
- b The age of the chicken and the number of eggs it lays are in inverse proportion, this means that as the age of the chicken increases, the number of eggs it lays decreases.

Working with percentages

Stretch it! £128

Stretch it! Let percentage rate = x

$$(1 + \frac{x}{100})^5 \times \text{£}100 = \text{£}110$$

$$(1 + \frac{x}{100})^5 = \frac{110}{100}$$

$$1 + \frac{x}{100} = \sqrt[5]{\frac{110}{100}}$$

$$1 + \frac{x}{100} = 1.02$$

$$\frac{x}{100} = 0.02$$

$$x = 2$$

Percentage interest is 2%

- 1 a $1.03 \times 50 = \text{£}51.50$
- b $2.48 \times 400 = 992$
- c $0.195 \times 64 = 12.48$
- 2 $45 - 40 = 5, \frac{5}{40} \times 100 = 12.5\%$
- 3 $24 \div 115 = 0.209, 0.209 \times 100 = 20.9^\circ\text{C}$
- 4 $15\,000 \times 1.20^3 = 25920$
- 5 20% is $\frac{1}{5}$ of the price.
 $30 \times 5 = \text{£}150$
- 6 $(200 \div 225) \times 100 = 88.9\%$ (to 1 d.p.)
 The number of employees in Year 2 is 88.9% of the number in Year 1.

Compound units

Stretch it! $\frac{100}{x}$ mph

- 1 $29.50 \div 0.18 = 164$ or $2950 \div 18 = 164$ units
- 2 Time = $\frac{80}{120} = \frac{2}{3}$ hour = 40 minutes
- 3 Density = $\frac{0.72}{3} = 0.24\text{ g/cm}^3$
- 4 Pressure = $\frac{12}{2} = 6\text{ N/m}^2$
- 5 $3\text{ m/s} = 3 \times 60\text{ m/minute} = 3 \times 60 \times 60\text{ m/hour}$
 $= 10800\text{ m/hour} = 10.8\text{ km/hour}$
- 6 0.6 litres per second = 0.6×60 litres per minute
 $= 0.6 \times 60 \times 60$ litres per hour
 $= 2160$ litres per hour.
 $2160 \div 4.55 = 475$ gallons
 475 gallons per hour (to the nearest whole number)
- 7 Bolt: $100\text{ m in } 9.58\text{ seconds} = 10.4\text{ m/s}$
 Cheetah: $120\text{ km/h} = 120000\text{ m/hour}$
 $= 120000 \div 60\text{ m/min}$
 $= 2000\text{ m/min}$
 $= 2000 \div 60\text{ m/sec} = 33.3\text{ m/s}$

The cheetah is faster.

Review it!





- 1 a $3.2 \times 1000 = 3200\text{ m}$
- b $9 \times 60 = 540$ seconds
- c $0.4 \times 1000 = 400\text{ ml}$
- 2 $4600 \div 1000 = 4.6\text{ km}$
- 3 $2.5 \times 60 = 150$ minutes
- 4 $1.1 \times 0.32 = 0.352\text{ m}^2$ or $110 \times 32 = 3520\text{ cm}^2$
- 5 $3 \times 10000 = 30000\text{ cm}^2$
- 6 $\frac{5}{12}$

- 7 $26:18 = 13:9$
- 8 $100 - 85 = 15$, $15 \div 3 = 5$ minutes
- 9 density = $\frac{345}{0.15} = 2300 \text{ kg/m}^3$
- 10 $10 - 8 = 2 \text{ km}$, $\frac{2}{8} \times 100 = 25\%$
- 11 $25 - 13 = 12$, $\frac{12}{25}$ OR 48%
- 12 $15 + 5 + 3 = 23$ mins
 $\frac{23}{90}$
- 13 $20 \div (\frac{4}{5}) = 25$ hours = 1 day and 1 hour
- 14 a $50 \div 5 = 10$, Josie: $1 \times 10 = 10$ marbles,
Charlie: $4 \times 10 = 40$ marbles, Charlie has 30 more.
b $C = 4J$
- 15 $\frac{100}{360} \times 100 = 28\%$
- 16 $0.8 \times 1200 = \text{£}960$
 $0.9 \times 960 = \text{£}864$
- 17 1 cm: 50 000 cm
50 000 cm = 0.5 km
 $3 \text{ km} \div 0.5 = 6$
6 cm
- 18 $1.02^3 \times 1500 = \text{£}1591.81$
- 19 $32\ 000 \div 4 = 8000$ people
- 20 $393 \div 125 = 3.144$ hours = 3 hours 9 minutes
- 21 $2.50 + 1.90 + (2 \times 5.30) = \text{£}15$
 $1.05 \times \text{£}15 = \text{£}15.75$
- 22 $37 + 15 + 4 + 19 = 75$
 $\frac{15}{75} \times 100 = 20\%$
- 23 $0.045 \times 3000 = \text{£}135$
 $3000 + (5 \times 135) = \text{£}3675$
- 24 $30 \div 3 = 10$
boys = $2 \times 10 = 20$
Girls = $1 \times 10 = 10$
Boys = $20 - 2 = 18$
Girls = $10 + 3 = 13$
18:13
- 25 Men to women is $7:6 = 35:30$
Ratio of women to children is $15:2 = 30:4$
Ratio of men to women to children is $35:30:4$
 $35 + 30 + 4 = 69$
 $3450 \div 69 = 50$
 $35 \times 50 = 1750$ men
- 26 No – for two things to be in direct proportion when one is zero the other must be zero; the graph does not go through the origin so this is not the case.
- 27 Neither, since the time taken to cook increases as the weight increases it is not in indirect proportion. It is not in direct proportion since a graph to illustrate the relationship would not go through the origin.
- 28 speed = $\frac{\text{distance}}{\text{time}} = \frac{0.05}{17} = \frac{1}{340}$ hours = $\frac{3}{17}$ mins
= 11 seconds
- 29 She is incorrect since the ratio of females to males must be the same for them to have equivalent proportions: $35:60$ is not equivalent to $12:37$.
- 30 $90 \times 2.5 = 225 \text{ g}$

Geometry and measures

Measuring and drawing angles

- 1 a 43° b Acute



- b 18°

Using the properties of angles

- 1 Angles around a point add up to 360° so:
 $a + 112 + 88 + 106 = 360$
 $a + 306 = 360$
 $a = 54^\circ$
- 2 a $a = (180 - 40) \div 2 = 70^\circ$
Base angles of an isosceles triangle are equal.
- b Exterior angle of a triangle is equal to the sum of the interior angles at the other two vertices so:
 $b = 70 + 40$
 $b = 110^\circ$
Or, angles on a straight line add up to 180° so:
 $b = 180 - 70 = 110^\circ$
- 3 Angles around a point add up to 360° so:
 $5x + 9x + 108 = 360$
 $14x + 108 = 360$
 $14x = 252$
 $x = 18^\circ$
- 4 a $x = 180 - 126 = 54^\circ$
Angles on a straight line add up to 180° .

b Angles in a quadrilateral add up to 360° so:

$$y + 135 + 54 + 88 = 360$$

$$y + 277 = 360$$

$$y = 83^\circ$$

5 a Angles on a straight line add up to 180° so:

$$x = 180 - 84$$

$$x = 96^\circ$$

b $y = 96^\circ$

Use the fact that corresponding angles are equal, then the fact that vertically opposite angles are equal.

Or, use the fact that alternate angles are equal, then use angles on a straight line add up to 180° .

6 a Base angles of an isosceles triangle are equal so $a = 58^\circ$.

b Angles in a triangle add up to 180° so:

$$b = 180 - 58 - 58$$

$$b = 64^\circ$$

c Alternate angles are equal so $c = 58^\circ$ (since angle $a =$ angle c).

Or, since opposite angles of a parallelogram are equal:

$$b + c = 122$$

$$64 + c = 122$$

$$c = 58^\circ$$

7 Angle $BAD = 62^\circ$ (Opposite angles of a parallelogram are equal)

Angle $ADE = 62^\circ$ (Alternate angles are equal)

$x = 180 - 62 - 62$ (Base angles of an isosceles triangle are equal)

$$x = 56^\circ$$

8 Angle $ACB = 36^\circ$ (Base angles of an isosceles triangle are equal)

Angle $ABC = 180 - 36 - 36$ (Angles in a triangle add up to 180°)

Angle $ABC = 108^\circ$

$x = 108^\circ$ (Alternate angles are equal)

Using the properties of polygons

Stretch it!

1 The angle sum of a triangle is 180° .

Sum of interior angles of a hexagon = $4 \times 180^\circ = 720^\circ$.

2

Polygon	Number of sides (n)	Number of triangles formed	Sum of interior angles
Triangle	3	1	180°
Quadrilateral	4	2	360°
Pentagon	5	3	540°
Hexagon	6	4	720°
Heptagon	7	5	900°
Octagon	8	6	1080°
Decagon	10	8	1440°

3 $n - 2$

4 $180 \times (n - 2)$

Stretch it! Exterior angle of a regular hexagon = $360 \div 6 = 60^\circ$

Interior angle = $180 - 60 = 120^\circ$

Three hexagons meet at a point, so $120 + 120 + 120 = 360^\circ$

Similarly, interior angle of an octagon = $180 - (360 \div 8) = 135^\circ$

Interior angle of a square = 90° , so $135 + 135 + 90 = 360^\circ$.

Regular pentagons have an interior angle of 108° . This does not divide equally into 360° , so these shapes will not fit together at a point in this way.

1 Regular decagon has 10 equal sides.

Exterior angle = $360^\circ \div 10 = 36^\circ$

2 a Number of sides = $360^\circ \div 15^\circ = 24$

b Angles on a straight line add up to 180° so:

Interior angle + exterior angle = 180

Interior angle + $15 = 180$

Interior angle = 165°

Sum of interior angles = $24 \times 165 = 3960^\circ$

3 Sum of interior angles of regular pentagon = $180^\circ \times (5 - 2)$

= $180^\circ \times 3 = 540^\circ$

One interior angle of regular pentagon = $540^\circ \div 5 = 108^\circ$

If this is a regular pentagon, $AB = AE$ and triangle ABE is isosceles.

In triangle ABE :


angle $ABE =$ angle $AEB = (180^\circ - 108^\circ) \div 2 = 36^\circ$

so angle $CBE = 108^\circ - 36^\circ = 72^\circ$

Using bearings

1 a $360 - 45$ (acute angle) = 315°

b




2 Bearing of P from Q = $180^\circ + 164^\circ = 344^\circ$


3 Kirsty is correct.

The bearing is 314° ($360^\circ - 46^\circ$) as it must be measured clockwise from North.

Properties of 2D shapes


Stretch it!





b kite

2 a 8 possible lines of symmetry:



b 8

3 a A rectangle has rotational symmetry of order 2.

b A **rhombus** has all sides equal and rotational symmetry of order 2.

c A kite has **1** line of symmetry and **no** rotational symmetry.

d The diagonals of a **square** and a **rhombus** bisect each other at 90° .

Congruent shapes

1 Any accurate copy of shape A, in any orientation.

2 a Corresponding angles are equal so $x = 120^\circ$

b Corresponding sides are the same length so $y = 12$ cm


3 a SSS (each triangle has equal sides: 3 cm, 3 cm, 2.5 cm)

b ASA (two angles, 70° and 60° , and the included side, 8 cm, are equal)



Constructions

Stretch it!




A triangle with sides of 5 cm with constructions lines indicating the use of compasses



Angle size 60°



Not drawn accurately



Drawing circles and parts of circles



1 a A **chord** is a straight line that does not pass through the centre of a circle but touches the circumference at each end.

b A **tangent** is a straight line that touches the outside of a circle at one point only.



c A **diameter** is a straight line through the centre of a circle that touches the circumference at each end.

d An **arc** is part of the circumference of a circle.


- e A **radius** is a straight line from the centre of a circle that is half the length of the diameter.
 f The part of a circle that has a chord and an arc as its boundary is called a **segment**.



Loci



- 3 a and b



Perimeter

- $4 \times 7.2 = 28.8 \text{ cm}$
- $7 + 9 + 9 + 5 + 5 + 7 = 42 \text{ cm}$
- Curved edge = $2\pi r \div 2 = (2 \times \pi \times 4) \div 2 = 4\pi$
 Perimeter = $4\pi + 2 \times r = 4\pi + 8 \text{ cm}$
 So $k = 4$ and $b = 8$
- Perimeter = $(\pi \times 30) + 100 + 100 = 200 + 30\pi \text{ m}$
- Perimeter = $(\frac{1}{2} \times \pi \times 32) + 32 + 32 = 16\pi + 64 \text{ cm}$
 Ribbon = $16\pi + 64 + 5 = 16\pi + 69 \text{ cm} = 119.3 \text{ cm}$
 120 cm must be bought
 $12 \times \text{£}0.15 = \text{£}1.80$

Area

Stretch it! Area of a semicircle = $\frac{\pi r^2}{2}$,
 area of a quarter circle = $\frac{\pi r^2}{4}$

- $4.5 \times 2 = 9.0 \text{ cm}^2$
 - $3 \times 1.5 = 4.5 \text{ cm}^2$
 - $\frac{(5+9)}{2} \times 4 = 28.0 \text{ cm}^2$
 - $\frac{1}{2} \times 2 \times 5 = 5.0 \text{ cm}^2$
 - $\pi \times 4.5^2 = 63.6 \text{ cm}^2$
- Length of side = $12 \div 4 = 3 \text{ cm}$
 Area = $3^2 = 9 \text{ cm}^2$
- Shaded triangles would fit together to form one triangle with base $10 - 6 = 4$.
 So area of shaded triangles = $\frac{1}{2} \times 4 \times 7 = 14 \text{ cm}^2$
 Area of trapezium = $\frac{(6+10)}{2} \times 7 = 56 \text{ cm}^2$
 Fraction of the shape that is shaded = $\frac{14}{56} = \frac{1}{4}$
- Area of whole shape = $6 \times 8 = 48 \text{ cm}^2$
 Fraction shaded = $\frac{6}{16} = \frac{3}{8}$
 Area shaded = $(\frac{3}{8}) \times 48 = 18 \text{ cm}^2$
- Area of square = $46 \times 46 = 2116 \text{ cm}^2$
 Each circle has radius = 11.5 cm
 Area of four circles = $4 \times \pi \times 11.5^2 = 1661.9 \text{ cm}^2$
 Shaded area = $2116 - 1661.9 = 454.1 \text{ cm}^2$

Sectors


- Area = $\frac{1}{2} \times \pi \times 5^2 = 39.3 \text{ cm}^2$
 Perimeter = $\frac{1}{2} \times \pi \times 10 + 10 = 25.7 \text{ cm}$
- Area = $\frac{3}{4} \times \pi \times 4^2 = 12\pi \text{ cm}^2$
- Area = $\frac{1}{2} \times \pi \times 3^2 = 14.1 \text{ m}^2$
 $14.1 \div 2 = 7.05$, so 8 bags needed.
 $8 \times 14.99 = \text{£}119.92$

3D shapes


Stretch it!

3D shape	Faces	Edges	Vertices
Cube	6	12	8
Cuboid	6	12	8
Square-based pyramid	5	8	5
Tetrahedron	4	6	4
Triangular prism	5	9	6
Hexagonal prism	8	18	12



Stretch it!






- 2 Draw a square in the middle with sides of 4 units (1 unit represents 1 cm). Set your compasses to 6 units and draw pairs of intersecting arcs from the corners of the square. These are the apices (top points) of the triangular sides. Draw lines for the sides of the triangles.



- 1 a 6 possible rectangular faces:



- b Kelli has not counted the hidden edges.



Volume

- 1 $\frac{4}{3} \times \pi \times 4.5^3 = 381.7 = 382 \text{ cm}^3$ (to 3 s.f.)
- 2 $\pi r^2 h + \frac{1}{3} \pi r^2 h = \pi \times 0.5^2 \times 2 + \frac{1}{3} \times \pi \times 0.5^2 \times 1.5$
 $= 0.625 \pi = 1.96 \text{ m}^3$
- 3 $\frac{1}{3} \times \pi \times 6^2 \times 22 = \frac{1}{3} \times 792 \times \pi = 264\pi \text{ cm}^3$
 $k = 264$
- 4 Volume of water = $18 \times 7 \times 7 = 882 \text{ cm}^3$
 $882 = 7 \times 20 \times h$
 $882 = 140 \times h$
 $h = 6.3 \text{ cm}$

Surface area

- 1 $6 \times (5 \times 5) = 150 \text{ cm}^2$
- 2 $4\pi r^2 = 4 \times \pi \times 3^2 = 36\pi \text{ cm}^2$
- 3 $18 - 4 = 14 \text{ cm}^2$
- 4 Sloping surface = $\pi \times 14 \times 45 = 630\pi \text{ cm}^2$
 Base = $\pi \times 14^2 = 196\pi \text{ cm}^2$
 Total surface area = $196\pi + 630\pi = 826\pi$
 Percentage yellow = $\frac{630}{826} \times 100 = 76.3\%$

Using Pythagoras' theorem

- 1 Using Pythagoras' theorem $c^2 = a^2 + b^2$:

$$\begin{aligned} AC^2 &= AB^2 + BC^2 \\ 15^2 &= 11^2 + BC^2 \\ BC^2 &= 15^2 - 11^2 = 104 \\ BC &= \sqrt{104} \\ BC &= 10.2 \text{ cm (to 3 s.f.)} \end{aligned}$$

- 2 $c^2 = a^2 + b^2$

$$\begin{aligned} c^2 &= 3.6^2 + 4.8^2 \\ c^2 &= 36 \\ c &= \sqrt{36} \\ c &= 6 \end{aligned}$$

The ladder is 6 m long.

- 3 Using Pythagoras' theorem $c^2 = a^2 + b^2$:

$$\begin{aligned} XZ^2 &= XY^2 + YZ^2 \\ 15^2 &= XY^2 + 9^2 \\ XY^2 &= 15^2 - 9^2 = 144 \\ XY &= \sqrt{144} \\ XY &= 12 \text{ cm} \end{aligned}$$

$$\begin{aligned} \text{Area} &= \frac{1}{2}bh = \frac{1}{2} \times 9 \times 12 \\ \text{Area} &= 54 \text{ cm}^2 \end{aligned}$$


- 4 If the triangle is right-angled, $PQ^2 = PR^2 + RQ^2$

$$\begin{aligned} PQ^2 &= 13^2 = 169 \\ PR^2 + RQ^2 &= 8^2 + 5^2 = 64 + 25 = 89^* \\ PQ^2 &\neq PR^2 + RQ^2 \end{aligned}$$

Claudia is not correct.


Notice that $PR + RQ = 8 + 5 = 13 \text{ cm} = \text{length of } PQ$, so PQR isn't a triangle at all, it is just a straight line!

- 5 $P(-2, 5)$, $Q(5, -3)$




$$\begin{aligned} PQ^2 &= 8^2 + 7^2 = 113 \\ PQ &= \sqrt{113} \\ PQ &= 10.63 \text{ (to 2 d.p.)} \end{aligned}$$

- 6



$$\begin{aligned} A(2, 0), B(5, 6) \\ AB^2 &= (6 - 0)^2 + (5 - 2)^2 \\ &= 6^2 + 3^2 \\ &= 45 \\ AB &= \sqrt{45} = 6.7 \text{ (1 d.p.)} \end{aligned}$$

- 7



Using Pythagoras' theorem $c^2 = a^2 + b^2$:

$$\begin{aligned} AB^2 &= AX^2 + BX^2 \\ AB^2 &= 7^2 + 20^2 = 449 \\ AB &= \sqrt{449} \\ AB &= 21.2 \text{ (to 3 s.f.)} \end{aligned}$$

$$\begin{aligned} \text{Perimeter of field } ABCD &= 15 + 20 + 8 + 21.2 \\ &= 64.2 \approx 65 \text{ m} \end{aligned}$$

$$\text{Cost of fencing} = 65 \times \text{£}14 = \text{£}910$$

Trigonometry

Stretch it!

Opposite could have been 1 m, hypotenuse could have been 2 m. They could be any lengths that keep opposite and hypotenuse in the ratio 1:2.

- 1 a 0.4 b 0.6 c 1.0
d 26.6 e 48.6 f 54.7
- 2 $\cos 72^\circ = \frac{MN}{15}$ $MN = 15 \cos 72^\circ = 4.6 \text{ cm}$

- 3 $\tan ABC = \frac{6}{7}$
 $ABC = \tan^{-1}\left(\frac{6}{7}\right)$
 $ABC = 40.6^\circ$

- 4 Let x be the depth of water.
 $\sin 15^\circ = \frac{x}{10}$
 $x = 10 \sin 15^\circ$
 $x = 2.6 \text{ m}$

Exact trigonometric values

- 1 a 0.5 b 0 c 0
d $\frac{1}{\sqrt{2}}$ e $\sqrt{3}$
- 2 $\tan 45^\circ = 1 = \frac{\text{opposite}}{\text{adjacent}} = \frac{4}{AC}$
Therefore $AC = 4 \text{ cm}$
 $\cos 45^\circ = \frac{1}{\sqrt{2}} = \frac{4}{BC}$
 $BC = 4\sqrt{2}$
Therefore $BC = 4\sqrt{2} \text{ cm}$


*This answer differs from the one in the Revision Guide due to an error in our first edition. This answer has now been re-checked and corrected.

- 3 Since: $\tan 30^\circ = \frac{1}{\sqrt{3}}$ one angle must be 30° and therefore the other is 60°
- 4 $\sin 30^\circ = \frac{1}{2}$ therefore $ABC = 30^\circ$
- 5 $\cos 30^\circ = \frac{\sqrt{3}}{2} = 0.866$ (3 d.p.) $\tan 45^\circ = 1$
Smallest to largest = $0.5, \frac{3}{4}, \cos 30^\circ, \tan 45^\circ$


Transformations

Stretch it!


Yes. Reflection in the x -axis followed by reflection in the y -axis (or vice versa) will always produce a rotation of 180° .



- 2 Translation by vector $\begin{pmatrix} -4 \\ -2 \end{pmatrix}$
- 3 a and b



- 4 Reflection in the y -axis
- 5 Enlargement by scale factor $\frac{1}{2}$, centre $(3, 3)$
- 6 a and b



- c Rotation of 90° clockwise about $(0, 0)$

Similar shapes

Stretch it!

Perimeter of $ABC = 3 + 6 + 5 = 14$ cm






Perimeter of $DEF = 6 + 12 + 10 = 28$ cm


The perimeter of a shape enlarged by scale factor 2 will also be enlarged by scale factor 2.

In general, all lengths on an enlarged shape, including the perimeter, are enlarged by the same scale factor.

- 1 a Angle $DFE = 30^\circ$ (Corresponding angles are the same)
- b Scale factor of enlargement = $\frac{\text{enlarged length}}{\text{original length}} = \frac{12}{3} = 4$
Length of $EF = 4$ cm $\times 4 = 16$ cm
- c Length of $AB = 8$ cm $\div 4 = 2$ cm
- 2 a Angle $MLO = 80^\circ$ (Corresponding angles are the same: angle $MLO =$ angle QPS)
- b Scale factor of enlargement = $\frac{\text{enlarged length}}{\text{original length}} = \frac{9}{3} = 3$
Length of $QR = 4.4$ cm $\times 3 = 13.2$ cm
- c Length of $LO = 12$ cm $\div 3 = 4$ cm

Vectors





c $\mathbf{a} = \begin{pmatrix} 4 \\ 3 \end{pmatrix}$

So $-2\mathbf{a} = -2 \times \begin{pmatrix} 4 \\ 3 \end{pmatrix} = \begin{pmatrix} -8 \\ -6 \end{pmatrix}$

3 a $\mathbf{a} + \mathbf{b} = \begin{pmatrix} 2 \\ 6 \end{pmatrix} + \begin{pmatrix} -5 \\ 3 \end{pmatrix} = \begin{pmatrix} -3 \\ 9 \end{pmatrix}$

b $\mathbf{a} + 2\mathbf{b} = \begin{pmatrix} 2 \\ 6 \end{pmatrix} + 2 \times \begin{pmatrix} -5 \\ 3 \end{pmatrix} = \begin{pmatrix} 2 \\ 6 \end{pmatrix} + \begin{pmatrix} -10 \\ 6 \end{pmatrix} = \begin{pmatrix} -8 \\ 12 \end{pmatrix}$

c $\mathbf{a} - \mathbf{b} = \begin{pmatrix} 2 \\ 6 \end{pmatrix} - \begin{pmatrix} -5 \\ 3 \end{pmatrix} = \begin{pmatrix} 7 \\ 3 \end{pmatrix}$

d $\mathbf{b} - 2\mathbf{a} = \begin{pmatrix} -5 \\ 3 \end{pmatrix} - 2 \times \begin{pmatrix} 2 \\ 6 \end{pmatrix} = \begin{pmatrix} -5 \\ 3 \end{pmatrix} - \begin{pmatrix} 4 \\ 12 \end{pmatrix} = \begin{pmatrix} -9 \\ -9 \end{pmatrix}$


4 a $\overrightarrow{PQ} = 4\mathbf{a}$ (\overrightarrow{PQ} and \overrightarrow{SR} are parallel and the same length)

b $\overrightarrow{QR} = -3\mathbf{b}$ (\overrightarrow{QR} and \overrightarrow{PS} are parallel and the same length; \overrightarrow{PS} has opposite direction to \overrightarrow{SP})

c $\overrightarrow{PR} = \overrightarrow{PQ} + \overrightarrow{QR}$
 $= 4\mathbf{a} - 3\mathbf{b}$

d $\overrightarrow{QS} = \overrightarrow{QR} + \overrightarrow{RS}$
 $= -3\mathbf{b} - 4\mathbf{a}$

Review it




b $BC = 3.8\text{cm}$


c $x = 50^\circ$

2 a 5 faces


b 6 vertices



4 A shape that matches the shape given in the question. It could be on its side or upside down.



5 Area $A = 7 \times (10 - 2 - 2) = 7 \times 6 = 42\text{ cm}^2$




Area $B = 10 \times 3 = 30\text{ cm}^2$

Total area $= 42 + 30 = 72\text{ cm}^2$

6 Area of parallelogram $= 3 \times 12 = 36\text{ cm}^2$

Length of side of square $= \sqrt{36} = 6\text{ cm}$

Perimeter of square $= 4 \times 6 = 24\text{ cm}$



8 Rotation of 180° about $(1, 0)$

9 Angle $ACB = 50^\circ$ (alternate angles are equal)

Angle $ABC = \text{angle } ACB = 50^\circ$ (base angles of an isosceles triangle are equal)

$x = 180^\circ - 50^\circ = 130^\circ$ (angles on a straight line add up to 180°)

10 Shaded area $= (10 \times 12) - ((\frac{1}{2} \times 12 \times 3) + (\frac{1}{2} \times 8 \times 7) + (\frac{1}{2} \times 10 \times 4))$
 $= 120 - (18 + 28 + 20)$
 $= 54\text{ cm}^2$

Proportion $= \frac{54}{120} = \frac{9}{20} = 45\%$


11 If triangle ABC is right-angled, $c^2 = a^2 + b^2$

$c^2 = 8^2 = 64$

$a^2 + b^2 = 4^2 + 6^2 = 16 + 36 = 52$

$c^2 \neq a^2 + b^2$ so triangle ABC is not right-angled.

12



13 a $\cos 45^\circ = \frac{1}{\sqrt{2}}$


b Ratio of adjacent to hypotenuse is 1:2

Therefore $AB = 3\text{ cm}$

14 a $a + 2b = \begin{pmatrix} 4 \\ -5 \end{pmatrix} + 2 \times \begin{pmatrix} 2 \\ 3 \end{pmatrix}$
 $= \begin{pmatrix} 4 \\ -5 \end{pmatrix} + \begin{pmatrix} 4 \\ 6 \end{pmatrix}$
 $= \begin{pmatrix} 8 \\ 1 \end{pmatrix}$

b $b - 2a = \begin{pmatrix} 2 \\ 3 \end{pmatrix} - 2 \times \begin{pmatrix} 4 \\ -5 \end{pmatrix}$
 $= \begin{pmatrix} 2 \\ 3 \end{pmatrix} - \begin{pmatrix} 8 \\ -10 \end{pmatrix}$
 $= \begin{pmatrix} -6 \\ 13 \end{pmatrix}$

15 Any correct answer will have two pairs of equal adjacent sides, two equal angles, and one line of symmetry.



16 Three lines of symmetry and all sides the same length mean it must be an **equilateral triangle**.

17 a 35°

Triangle WYZ is isosceles, and base angles of an isosceles triangle are equal.

b Angles in a triangle add up to 180° so:

$$b = 180 - 35 - 35$$

$$= 110^\circ$$

c Triangle XYZ is isosceles, and base angles of an isosceles triangle are equal so:

$$c = (180 - 70) \div 2 = 55^\circ$$

18 Using Pythagoras' theorem $c^2 = a^2 + b^2$:

$$AC^2 = AB^2 + BC^2$$

$$14^2 = 6^2 + BC^2$$

$$BC^2 = 14^2 - 6^2 = 160$$

$$BC = \sqrt{160}$$

$$BC = 12.6 \text{ cm (to 1 d.p.)}$$

19 Interior angle of a square = 90°

Sum of interior angles of an octagon (with $n = 8$)
 $= 180 \times (n - 2) = 180 \times (8 - 2) = 1080^\circ$

Interior angle of a regular octagon = $1080^\circ \div 8 = 135^\circ$


(Or, exterior angle of a regular octagon = $360^\circ \div 8 = 45^\circ$.
 Then interior angle = $180^\circ - 45^\circ = 135^\circ$)

Angles around a point add up to 360° so:

$$x = 360 - 90 - 135$$

$$x = 135^\circ$$

20



21 Divide the trapezium into a rectangle and a triangle. Draw a line DX parallel to AB , with X on the line BC . $BX = 5 \text{ cm}$, $CX = 7 \text{ cm}$.

Using Pythagoras' theorem $c^2 = a^2 + b^2$:

$$DC^2 = CX^2 + DX^2$$

$$DC^2 = 7^2 + 4^2 = 65$$

$$DC = \sqrt{65}$$

$$DC = 8.06 \text{ (to 2 d.p.)}$$

$$\text{Perimeter of } ABCD = 4 + 5 + 8.06 + 12 = 29.06 \text{ cm}$$

22 Length of arc = $\frac{1}{4} \times 2 \times \pi \times 4 = 2\pi$

$$\text{Perimeter} = 4 + (2 \times 9) + 4 + 2\pi = 32.3 \text{ cm}$$

23 Area of square = $6 \times 6 = 36 \text{ cm}^2$

$$\text{Area of circle} = \pi \times 3^2 = 9\pi \text{ cm}^2$$

$$\text{Shaded area} = 36 - 9\pi = 7.7 \text{ cm}^2$$

24 Volume of cylinder = $\pi \times 3^2 \times 15 = 135\pi \text{ cm}^3$

$$2 \text{ litres} = 2000 \text{ ml} = 2000 \text{ cm}^3$$


$$2000 \div 135\pi = 4.7$$

Glass can be completely filled 4 times.

25 Scale factor of enlargement = $\frac{\text{enlarged length}}{\text{original length}} = \frac{11}{5} = 2.2$

$$\text{Length } x = 6 \text{ cm} \times 2.2 = 13.2 \text{ cm}$$

26



$$DE^2 = 6^2 + 4^2$$

$$= 36 + 16$$

$$= 52$$

$$DE = 7.2 \text{ cm (1 d.p.)}$$

27 a Curved surface area = $\pi \times 6 \times 10 = 60\pi \text{ cm}^2$

$$\text{Base area} = \pi \times 6^2 = 36\pi \text{ cm}^2$$

$$\text{Total surface area} = 60\pi + 36\pi = 96\pi = 300 \text{ cm}^2 \text{ to 2 s.f.}$$

b Volume = $\frac{1}{3} \times \pi \times 6^2 \times 8 = 96\pi = 300 \text{ cm}^3$


- 28 $\tan x = \frac{8}{6}$
 $x = \tan^{-1}\left(\frac{8}{6}\right)$
 $x = 53.1^\circ$
- 29 Translation by vector $\begin{pmatrix} -7 \\ -6 \end{pmatrix}$
- 30 $\vec{AC} = \vec{AB} + \vec{BC}$
 $= 2\mathbf{a} + 3\mathbf{b} + 3\mathbf{a} - \mathbf{b}$
 $= 5\mathbf{a} + 2\mathbf{b}$

Probability

Basic probability

Stretch it! No – each time the probability of getting an even number is $\frac{1}{2}$. You would expect to get even numbers approximately 50 times but cannot guarantee it.

1 $\frac{4}{10}$



2 a Total number of sweets = $12 + 3 + 10 = 25$

$\frac{3}{25}$
 b $\frac{(3+10)}{25} = \frac{13}{25}$

- 3 Pair **a**, because when you flip a coin, you can't get both a head and a tail at the same time. (Prime numbers on a dice are 2, 3, 5 and odd numbers are 1, 3, 5, so events **b** are **not** mutually exclusive because 3 is in both groups.)
- 4 Pair **b**, because the first sweet chosen is replaced, so the possible outcomes of the second choice remain the same. (If the first sweet chosen is eaten, the possible outcomes of the second choice are altered, and so events **a** are **not** independent.)
- 5 $P(6) = 1 - (0.1 + 0.15 + 0.1 + 0.02 + 0.2)$
 $= 1 - 0.57 = 0.43$
- 6 $P(\text{green or red}) = 1 - 0.4 = 0.6$
 $P(\text{green}) = 2 \times P(\text{red})$
 $P(\text{red}) = \frac{0.6}{3} = 0.2$
 $P(\text{green}) = 2 \times 0.2 = 0.4$

Two-way tables and sample space diagrams

1

	Chicken	Beef	Vegetarian
Fruit	12	6	4
Cake	5	3	8
Total	17	9	12

- a 12 (this is worked out by using the numbers in the 'Total' row, which must add up to 38)
- b As shown in the table.
- 2 a


		Dice 1					
		1	2	3	4	5	6
Dice 2	1	2	3	4	5	6	7
	2	3	4	5	6	7	8
	3	4	5	6	7	8	9
	4	5	6	7	8	9	10
	5	6	7	8	9	10	11
	6	7	8	9	10	11	12

- b i $\frac{2}{36} = \frac{1}{18}$
 ii $\frac{3}{36} = \frac{1}{12}$
 iii 0


- 3 To score 6, the player must pick two cards showing 3. To score 2, the player must pick two cards showing 1. Since the probability of getting 3 and 3 is more than 0, and the probability of getting 1 and 1 is more than 0, there must be at least 2 of each of those numbers. So the cards must be 1, 1, 3, 3.

Sets and Venn diagrams

Stretch it! None




- b $A \cap B = \{\text{multiples of 6 less than 20}\}$ because these numbers are multiples of both 2 and 3.
- 2 a $C \cap T$ is the set of students who travel by car AND train
 $C' \cap B$ is the set of students who do NOT travel by car AND travel by bus.*
- b i $P(C) = \frac{(14 + 11 + 11 + 2)}{(14 + 11 + 11 + 2 + 17 + 19 + 26)} = \frac{38}{100} = \frac{19}{50}$
 ii $P(B \cup T) = \frac{(19 + 11 + 2 + 0 + 11 + 17)}{100} = \frac{60}{100} = \frac{3}{5}$
 iii $P(B' \cap T) = \frac{(11 + 17)}{100} = \frac{28}{100} = \frac{7}{25}$
- 3 $P(A \cap B) = \frac{3}{20}$ so there must be 3 elements in the intersection.
 $P(A) = \frac{3}{10} = \frac{6}{20}$ so there must be a total of 6 elements in A. The total number of elements must sum to 20.




Frequency trees and tree diagrams

- 1 a Apple = 123
 Pear = $347 - 123 = 224$
 Apple fruiting = 102
 Apple not fruiting = $123 - 102 = 21$
 Pear not fruiting = 34
 Pear fruiting = $224 - 34 = 190$

*This answer differs from the one in the Revision Guide due to an error in our first edition. This answer has now been re-checked and corrected.



b $\frac{190}{347}$




b $\frac{3}{5} \times \frac{3}{5} = \frac{9}{25}$

c $\frac{3}{5} \times \frac{3}{5} = \frac{9}{25}$

$\frac{2}{5} \times \frac{3}{5} = \frac{6}{25}$

$\frac{3}{5} \times \frac{2}{5} = \frac{6}{25}$

$P(\text{at least one prime}) = 1 - P(\text{no primes})$
 $= 1 - \frac{2}{5} \times \frac{2}{5} = 1 - \frac{4}{25} = \frac{21}{25}$



b $P(\text{two glass marbles}) = \frac{3}{7} \times \frac{2}{6} = \frac{6}{42}$

$P(\text{glass then plastic}) = \frac{3}{7} \times \frac{4}{6} = \frac{12}{42}$

$P(\text{plastic then glass}) = \frac{4}{7} \times \frac{3}{6} = \frac{12}{42}$

$P(\text{at least one glass}) = 1 - P(\text{both plastic})$
 $= 1 - \frac{4}{7} \times \frac{3}{6} = 1 - \frac{12}{42}$
 $= 1 - \frac{2}{7} = \frac{5}{7}$

Expected outcomes and experimental probability

Stretch it! The dice has not been rolled enough times to decide if it is biased. More tests need to be carried out.

1 $0.45 \times 300 = 135$

2 Red = $\frac{2}{10} = \frac{1}{5}$
 $\frac{1}{5} \times 100 = 20$ red sweets


3 $\frac{1}{2} \times 100 = 50$ primes

4 a Charlie – he has carried out the most tests.

b $\frac{(112 + 10 + 28)}{(112 + 10 + 28 + 74 + 7 + 19)} \times 10 = 6$

Review it!

1 $0.12 \times 250 = 30$



3 $1 - 0.3 = 0.7$

4 B, C

5 a $\frac{3}{5}$

b $(\frac{1}{5}) \times 25 = 5$

6

	Pizza	Pasta	Risotto	Total
Cake	12	6	1	19
Ice Cream	10	11	10	31
Total	22	17	11	50

7 $0.2 + 5x + 0.2 + x = 1$

$6x + 0.4 = 1$

$x = 0.1$

$P(\text{white}) = 5x + 0.2 = 0.7$

8 a No, he has not tested his dice enough times.

b $P(2) = \frac{9}{(12 + 9 + 16 + 7 + 6 + 0)} = \frac{9}{50}$
 $\frac{9}{50} \times 100 = 18$

9 $P(R, R) = 0.1 \times 0.5 = 0.05$

$P(R, G) = 0.1 \times 0.5 = 0.05$

$P(G, R) = 0.9 \times 0.5 = 0.45$

$0.05 + 0.05 + 0.45 = 0.55$

Or $P(\text{at least one red}) = 1 - P(\text{green, green})$
 $= 1 - (0.9 \times 0.5)$
 $= 1 - 0.45$
 $= 0.55$

10 a

		Dice					
		1	2	3	4	5	6
Coin	Heads	2	4	6	8	10	12
	Tails	3	4	5	6	7	8


b i $\frac{2}{12} = \frac{1}{6}$

ii $\frac{2}{12} = \frac{1}{6}$

11 a i 6 ii 1 iii 5

b $\frac{4}{8} = \frac{1}{2}$

12 a



b $\frac{7}{9} \times \frac{6}{8} = \frac{42}{72} = \frac{7}{12}$

13 a Possible fractions: $\frac{1}{2}, \frac{1}{3}, \frac{1}{4}, \frac{2}{3}, \frac{2}{4}, \frac{3}{4}$

Less than $\frac{1}{2}$ are $\frac{1}{3}$ and $\frac{1}{4}$.

$P(\text{less than } \frac{1}{2}) = \frac{2}{6} = \frac{1}{3}$

$\frac{1}{3} \times 30 = 10$

b $\frac{1}{3}$ is only a theoretical probability and therefore will not necessarily be accurate in real life.

14 45% of 300 = 135

135 boys and 165 girls.

$\frac{2}{3}$ of 135 = 90

$\frac{4}{5}$ of 165 = 132

Total playing sport = 222

Probability = $\frac{222}{300} = \frac{37}{50} = 0.74$

15 $P(\text{hooking a winning duck}) = \frac{5}{20} = 0.25$

If 100 people play, expected number of winners = $0.25 \times 100 = 25$ people.

The game makes $\text{£}1 \times 100 \text{ people} = \text{£}100$.

The money paid out in prizes = $25 \text{ winners} \times \text{£}2 = \text{£}50$

Profit = $\text{£}100 - \text{£}50 = \text{£}50$

16 a Milo will have the better estimate as he has surveyed a greater number of people.

b Number of left-handed students = $5 + 4 + 7 + 7 = 23$

Number of right-handed students = $23 + 18 + 51 + 60 = 152$

$P(\text{left-handed}) = \frac{23}{23 + 152} = \frac{23}{175}$

$\frac{23}{175} \times 2000 = 262.8$

You would expect to find 263 left-handed students in a school with 2000 students.

Statistics

Data and sampling

Stretch it! A random sample could be taken; you could allocate a number to each pupil and randomly generate the numbers to survey. Any method is acceptable as long as each person in the school has an equally likely chance of being chosen. Alternatively a stratified sample could be taken.

1 Primary source: Recording the data by measuring it yourself.

Secondary source: Any sensible source, e.g. the Meteorological Office, local paper etc.

2 Qualitative data.

*This answer differs from the one in the Revision Guide due to an error in our first edition. This answer has now been re-checked and corrected.

3 It is cheaper and quicker than surveying the whole population.

4 a The people working for an animal charity are more likely to be opposed to wearing real fur; every member of the population does not have an equal chance of being chosen.

b Surveying people in the street, a random telephone survey, any sensible method that ensures that any member of the population has an equal chance of being chosen.

5 a $\frac{3}{200} \times 800\,000 = 12\,000$

b The sample is relatively small. The sample is not a random sample as it is taken on one day in a year.

Frequency tables

1

Number of people on the bus	Frequency
0*–9	4
10–19	12
20–29	3
30–39	1

2 a

Number of courgettes	Frequency
0	1
1	0
2	1
3	1
4	9
5	3
6	0

b $(0 \times 1) + (1 \times 0) + (2 \times 1) + (3 \times 1) + (4 \times 9) + (5 \times 3) = 56$

3 a There are gaps between his groups – times that fall between groups cannot be recorded, e.g. 15.5 hours.

His groups do not have the same width.

b Although one or more of the data values may fall in the $30 \leq h < 40$ group, this doesn't mean that those people trained for 40 hours. They could have trained for any length of time between 30 and 40 hours.

Bar charts and pictograms

1 a $15 + 4 + 1 = 20$

b $4 + 1 = 5$

$\frac{5}{20} \times 100 = 25\%$

2 a $11 - 7 = 4$

b Total number of people surveyed = $18 + 18 + 12 + 3 = 51$

Total number of boys = $11 + 6 + 3 = 20$


$\frac{20}{51} \times 100 = 39.2\%$

c Proportion of boys who played two sports = $\frac{6}{18} = \frac{1}{3}$

Proportion of boys who played three sports = $\frac{3}{12} = \frac{1}{4}$

$\frac{1}{3} > \frac{1}{4}$ so the proportion who played two sports is larger.

- 3 $50 - (11 + 15) = 24$
 $24 \div 3 = 8$
 Therefore: $2 \times 8 = 16$ size 7 shoes
 $1 \times 8 = 8$ size 8 shoes




- 4 a $90 - 20 = 70$
 b Total number of bikes = $50 + 50 + 20 + 90 = 210$
 $\frac{50}{210} = \frac{5}{21}$

Pie charts

Stretch it!

Round appropriately – but check the angles sum to 360°

- 1 $27 + 42 + 21 = 90$
 $360^\circ \div 90 = 4^\circ$
 French = $27 \times 4 = 108^\circ$
 Spanish = $42 \times 4 = 168^\circ$
 German = $21 \times 4 = 84^\circ$



- 2 a $\frac{1.5}{360} \times 240 = 1$ student earned more than £40 000.
 b $\frac{288 + 63}{360} \times 100 = 97.5\%$ of students earned less than £30 000.
 3 a $18 + 10 = 28$
 b The bar chart, since the frequency is easy to read from the bar chart.

Measures of central tendency: mode

- 1 The other three must be 12.2.
 2 $1 < t \leq 2$
 3 Max is correct, the modal number of pets is the group with the highest frequency, therefore 2 pets is the mode.

Measures of central tendency: median

- 1 Ordering the data gives; 2.9, 3.1, 4.3, 6.5, 8.7, 9.2
 Median = $\frac{4.3 + 6.5}{2} = 5.4$

- 2 $29 + 28 + 30 + 3 + 10 = 100$
 $\frac{(100 + 1)}{2} = 50.5$ – median term is between the 50th and 51st terms.
 Both these lie in the $2 \leq b < 4$ class*.
 3 If there are 5 integers in the list, then the middle value is the 3rd integer.

Measures of central tendency: mean

Stretch it! a mode b mean/median c mean/median

- 1 a Total frequency = $12 + 3 + 5 = 20$
 Mean = $\frac{(2 \times 12) + (6 \times 3) + (10 \times 5)}{20} = 4.6$
 b You are using the midpoint of the groups as an estimate of the actual value for each group.
 2 $\frac{(5 \times 9) + 6}{5 + 1} = 8.5$
 3 No – they could be any pair of numbers which sum to 10.


Range

- 1 $9.5 - 0.7 = 8.8$
 2 a Girls = $18 - 15 = 3$
 b Boys = $18 - 16 = 2$
 3 Range for Athlete A = $15.2 - 13.0 = 2.2$
 Range for Athlete B = $15.2 - 14.3 = 0.9$
 Athlete A has the greatest range.
 4 $45\% - 10 = 35\%$ or $45\% + 30 = 75\%$

Comparing data using measures of central tendency and range

- 1 a i Mean = $\frac{(32 + 29 + 18 + 41 + 362 + 19)}{6} = \frac{501}{6} = 83.5$ minutes
 ii Ordered data: 18, 19, 29, 32, 41, 362
 Median = $\frac{(29+32)}{2} = 30.5$ minutes
 b The extreme value (362 mins) affects the mean but not the median.
 2 All the data is used to find the mean.
 3 Either as long as suitably justified:
 Car A – although the mean time is higher, it is more consistent in performance since the range is smaller.
 Car B – the acceleration is quicker on average.
 4 a and b The mode or median since the mean will not be a whole number and therefore not meaningful.

Time series graphs




*This answer differs from the one in the Revision Guide due to an error in our first edition. This answer has now been re-checked and corrected.

- 2 a 67°C
 b Approx. 27°C
 c No, since it is extrapolation (beyond the limits of the data).
- 3 a 17 000 b i April ii August
 c The number of tourists peaks in April and again in December. The low seasons are February/March and July/August/September/October*.


Scatter graphs

1 a and b*



- c Positive
 d This will vary according to the line of best fit: approximately 4.7 kg. A range of 4.6kg to 4.8kg would be acceptable.
 e This is beyond the limits of the data and therefore extrapolation.


2 a, b and c



- c The seeds failed to germinate or the seedling died.
 d The further the seedling is from the light source the shorter its height.
- 3 No, although the two things correlate one does not cause another. There may be many reasons why the crime rate is high in the area, perhaps there is poverty and inequality causing social tension.


Review it!

- 1 The sample is too small and he only asked his friends. His data is therefore not representative of the population of TV viewers.
- 2 a Margherita
 b Total frequency = 11 + 2 + 6 + 1 = 20
 $\frac{1}{20} = \frac{5}{100} = 5\%$
 c $360^\circ \div 20 = 18^\circ$
 Pepperoni = $1 \times 18^\circ = 18^\circ$
 (or 5% of $360^\circ = 18^\circ$)
- 3 a $\frac{90}{360} = \frac{1}{4}$
 b $45^\circ = \frac{1}{8}$ of 360°
 Therefore $\frac{1}{8}$ of the pie chart represents 60 cars.
 The whole pie chart = $8 \times 60 = 480$ cars
 c $(\frac{105}{360}) \times 480 = 140$ cars
- 4 a The number of people doing their grocery shopping online is increasing.
 b Any sensible answer, approximately 75%
 c No – it is outside the limits of the data therefore extrapolation.
- 5 If the mean is 6, then the total of the numbers is $6 \times 4 = 24$.
 If the mode is 7, this number must occur at least twice.
 The median is the average of the middle two numbers, as there are 4 numbers. It is 7, so the middle two numbers must both be 7.
 If all the numbers were 7, the mean would be $7 \times 4 = 28$, which is not true, so they are not all 7.
 If three of the numbers were 7, the fourth number would be 3, because $7 \times 3 + 3 = 24$ (the mean). However, the range is 6, and $7 - 3 = 4$. So 7 must appear only twice.
 If the biggest value is 8, the smallest must be $8 - 6 = 2$.
 Test: $2 + 7 + 7 + 8 = 24 =$ the correct total.
 So the numbers are 2, 7, 7 and 8, and you can now answer all the questions.
 The four numbers **must** be: 2, 7, 7, 8.
- a true
 b true
 c true
 d false
- 6 a Comparative bar chart or compound bar chart:



*This answer differs from the one in the Revision Guide due to an error in our first edition. This answer has now been re-checked and corrected.

Or:



b Total number of students = $(3 + 5 + 1 + 0 + 8 + 2 + 2 + 4) = 25$

Number of cats = $3 + 8 = 11$
 $\frac{11}{25}$

7 a Total frequency = $17 + 2 + 32 + 23 + 9 = 83$
 Median value = $\frac{(83 + 1)}{2} = 42\text{nd term}$

42nd term is in group 40–59
 Median class = 40–59

b The youngest person is between 0 and 19, the youngest may be any age in this range and the oldest is between 80 and 99 therefore any age in this range.

8 a 7

b Size 5

c Mean = $\frac{(3 \times 2) + (4 \times 1) + (5 \times 7) + (6 \times 5) + (7 \times 3)}{2 + 1 + 7 + 5 + 3} = 5.\bar{3}$


d Mode – the mean is not an actual shoe size.

9 a $\frac{50}{150} = \frac{1}{3}$

b $60 - 40 = 20$

c Biology

10 a* and **c**



b Negative

d Approximately 40 minutes: it depends on line of best fit.


e This is outside the limits of the data and therefore extrapolation.

f As the age of the customer increases the time spent on the phone decreases.

11 a $\frac{(65 \times 3) + (75 \times 5) + (85 \times 2)}{3 + 5 + 2} = 74 \text{ kg}$

b The midpoint of the class is used as the age of each of the patients rather than the actual age.

12 Annual income for surveyed population



13 Mean = $\frac{(10.3 \times 10) + 9.5}{11} = 10.2$ (1 d.p.)

14 Mean is 3.8 so the sum of the scores is $3.8 \times 5 = 19$

Mode is 3 so she must roll at least two 3s.

Range is 4.

If the range is 4 then the lowest and highest must be either 1 and 5 or 2 and 6.

The numbers are: 2, 3, 3, 5 and 6

*This answer differs from the one in the Revision Guide due to an error in our first edition. This answer has now been re-checked and corrected.

AQA Foundation Mathematics Exam Practice Book

Full worked solutions

Number

Factors, multiples and primes

1 $18 = \begin{matrix} 2 \\ \times \end{matrix} \times \begin{matrix} 3 \\ \times \end{matrix} \times 3$
 $24 = \begin{matrix} 2 \\ \times \end{matrix} \times 2 \times 2 \times \begin{matrix} 3 \\ \times \end{matrix}$

Find the factors that they both share (2 and 3) and multiply together:

$2 \times 3 = 6$, so HCF is 6.

2 17, 19, 23 are the only numbers in this range with only 2 factors (1 and the number itself).

3 $60 = 20 \times 3$
 $= 2 \times 2 \times 5 \times 3$
 $= 2^2 \times 3 \times 5$

4 Drummer 1 hits her drum at: 6 12 18 24 30 36 42 48 54 60 seconds

Drummer 2 hits his drum at: 8 16 24 32 40 48 56 seconds

They hit their drums at the same time twice (two times), after 24 seconds and after 48 seconds.

Ordering integers and decimals

1 Negative numbers are smaller than zero

-12 is further left on the number line than -8, -1 is larger than -8 (and -12) so it appears next.

Then comes 0, then 2.

So the order is:

-12, -8, -1, 0, 2

2 First look at the place value for 10ths: 0.32 and 0.3 have the higher number of 10ths.

Now compare their 100ths. 0.32 has 2 100ths but 0.3 doesn't have any, so it's smaller.

Similarly, 0.23 and 0.203 both have 2 10ths, but 0.23 is bigger than 0.203 because it has 3 100ths while 0.203 only has 3 1000ths.

So the order is:

0.32, 0.3, 0.23, 0.203

3 a $-4 < 0.4$ (the negative number is smaller)

b $4.200 < 4.3$ (the larger number has more 10ths)

c $-0.404 > -0.44$ (because they are both negative, the one with more 100ths is smaller)

d $0.33 < 0.4$ (the larger number has more 10ths)

Calculating with negative numbers

1 a $-7 + -3 = -7 - 3 = -10$

b $-7 - -3 = -7 + 3 = -4$

c $8 + -5 - -2 = 8 - 5 + 2 = 5$

d $-4 - -6 + -1 = -4 + 6 - 1 = 1$

2 a -18

b $-12 \div -3 = 12 \div 3 = 4$

c $-4 \times -2 \times 5 = 4 \times 2 \times 5 = 40$

d $(-24 \div 3) \times 2 = -8 \times 2 = -16$

3 $-7 - 4 = -11$, $-11 \times -2 = 22$

4 Let a = number of correct answers, b = number of incorrect answers

$3a - 2b = -5$ (1)

There are five questions, so $a + b = 5$ and $b = 5 - a$ (2)

Substituting this for b in (1): $3a - 2(5 - a) = -5$

$3a - 10 + 2a = -5$

$5a = 5$

$a = 1$

Substituting this in (2):

$b = 5 - 1$

$b = 4$

Sally got 1 correct answer and 4 incorrect answers in the test.

Multiplication and division

1 a
$$\begin{array}{r} 357 \\ \times 6 \\ \hline 2142 \end{array}$$

2142

b
$$\begin{array}{r} 261 \\ \times 43 \\ \hline 783 \quad (=261 \times 3) \\ 10440 \quad (=261 \times 40) \\ \hline 11223 \end{array}$$

11223

c
$$\begin{array}{r} 092 \\ 6 \overline{)55^12} \end{array}$$

92

d
$$\begin{array}{r} 052 \\ 13 \overline{)676} \\ \underline{-65} \\ 026 \end{array}$$

52

2 a
$$\begin{array}{r} 012 \text{ remainder } 12 \\ 24 \overline{)306} \end{array}$$

So 12 boxes are filled.

b $24 \times 12 = 20 \times 12 + 4 \times 12 = 240 + 48 = 288$

$300 - 288 = 12$

There are 12 books left over.

3 $12500 - 440 = 12060$

$$\begin{array}{r} 00335 \\ 36 \overline{)121060} \\ \underline{-108} \\ 1216 \\ \underline{-108} \\ 0180 \end{array}$$

Each repayment is £335.

4 $52 - 6 = 46$ weeks $46 \times 26 = 1196$ hours

Calculating with decimals

1 First note the combined number of decimal places in both numbers (2).

Remove the decimal points to do the calculation:

$$\begin{array}{r} 92 \\ \times 83 \\ \hline 276 \quad (=92 \times 3) \\ 7360 \quad (=92 \times 80) \\ \hline 7636 \\ \hline 1 \end{array}$$

Now you've got the digits right put the decimal point back, counting in from the right 2 places, to give a number with 2 decimal places:

76.36

$$\begin{array}{r} 2 \quad 22.50 \\ + 19.99 \\ \hline 42.49 \\ \\ 50.00 \\ - 42.49 \\ \hline 07.51 \end{array}$$

She should get £7.51 change.

$$3 \quad \begin{array}{r} 038.29 \\ 6 \overline{)229.754} \end{array}$$

38.29

$$4 \quad \text{Kirsty raises } \frac{172.50}{5+1}$$

Kirsty raises $\text{£}28.75 \times 5 = 28.75 \times 10 \div 2 = 287.5 \div 2 = 143.75$

$$\begin{array}{r} 028.75 \\ 6 \overline{)172.50} \end{array}$$

Kirsty raises

£143.75

$$\begin{array}{r} 6 \quad 11 \quad 14 \quad 1 \\ 172.50 \\ - 143.75 \\ \hline 028.75 \end{array}$$

Flo raises £28.75

Rounding and estimation

- The first non-zero digit is 7, so round the digit after the 8. This is below 5, so the 8 doesn't change.
0.798
 - Look at the digit in the third decimal place. It is 5 or above (8), so round the 9 up to 10 and the 7 up to 8.
0.80
- $\frac{9.74 \times 4.02}{7.88} \approx \frac{10 \times 4}{8} = 5$
- $40 \times 500 = 20000$
 $20000 - 12500 = \text{£}7500$
 - Overestimate, because the concert ticket price and number of tickets sold were rounded up, and so the amount of income was estimated more than it really is.

Converting between fractions, decimals and percentages

- There are $0 \times 10^{\text{ths}}$, $7 \times 100^{\text{ths}}$ and $1 \times 1000^{\text{ths}}$
 $= 071 \times 1000^{\text{ths}}$ so: $\frac{71}{1000}$
 - $63 \div 100 = 0.63$
 - $0.4 \times 100 = 40\%$
 - $32\% = \frac{32}{100} = \frac{8}{25}$
- $5 \div 16 = 0.3125$
 - To convert a number to a percentage, multiply its decimal value by 100.
 $0.3125 \times 100 = 31.25\%$
- $\frac{5}{8} = 0.625$, $60\% = 0.6$, so 0.65 is the largest.

Ordering fractions, decimals and percentages

- $\frac{1}{2} = \frac{5}{10} = 0.5$, so
 $\frac{1}{2} < 0.6$
 - $\frac{3}{4} = 3 \div 4 = 0.75$, so
 $\frac{3}{4} > 0.7$

$$c \quad \frac{-3}{10} = -0.3, \text{ so}$$

$$\frac{-3}{10} < 0.2$$

- LCM of 12, 15 and 20 is 60

$$\frac{5}{12} = \frac{25}{60}$$

$$\frac{7}{15} = \frac{28}{60}$$

$$\frac{9}{20} = \frac{27}{60}$$

So order from lowest to highest is $\frac{5}{12}$, $\frac{9}{20}$, $\frac{7}{15}$

$$b \quad 45\% = \frac{45}{100} = 0.45$$

$$\frac{1}{25} = \frac{4}{100} = 0.04$$

$$0.04 < 0.4 < 0.45$$

So order is:

$$\frac{1}{25}, 0.4, 45\%$$

- Shop C is cheapest ($\frac{2}{5} = 40\%$), then Shop A ($\frac{1}{3} = 33.3\% \dots$), and Shop B offers the least discount at 30%.

$$4 \quad \frac{5}{9} = 0.\dot{5}$$

$$38.5\% = 0.385$$

$$\frac{3}{10} = 0.3$$

So the order is $\frac{5}{9}$, 38.5%, 0.38, $\frac{3}{10}$

Calculating with fractions

- $\frac{1}{5} + \frac{4}{9} = \frac{9}{45} + \frac{20}{45} = \frac{9+20}{45} = \frac{29}{45}$
- $2\frac{3}{4} - 2\frac{2}{3} = \frac{11}{4} - \frac{8}{3} = \frac{33}{12} - \frac{32}{12} = \frac{1}{12}$
- $1\frac{5}{6} \times \frac{2}{7} = \frac{11}{6} \times \frac{2}{7} = \frac{22}{42} = \frac{11}{21}$
- $6 \div \frac{3}{5} = 6 \times \frac{5}{3} = \frac{30}{3} = 10$
Jo can make 10 necklaces.

Percentages

- $\frac{40}{100} \times 25 = 10$
- $16 \times 0.85 = \text{£}13.60$
- $12450 \times 1.14 = 14193$
- $40 \times 7 \times 3 = \text{£}840$
 $840 \times 1.2 = \text{£}1008$

Order of operations

- $3^2 - 6 \div (2 + 1) = 9 - \frac{6}{3} = 9 - 2 = 7$
- $2^3 + 3x\sqrt{25} = 8 + (3 \times 5) = 8 + 15 = 23$
- $(1.7 - 0.12)^2 + \sqrt[3]{4.096} = 4.0964$

Exact solutions

- Area of triangle = $\frac{1}{2} \times \text{base} \times \text{vertical height} = 0.5 \times 0.76 \times 0.35 = 0.133 \text{ cm}^2$
- $(1\frac{1}{3})^2 = (\frac{4}{3})^2 = \frac{16}{9} = 1\frac{7}{9} \text{ m}^2 \text{ cm}^2$
- $\sqrt{2} \times \sqrt{6} = \sqrt{12} = 2\sqrt{3} \text{ cm}^2$
- Area of a circle = πr^2
The fraction of the circle shown = $\frac{3}{4}$
The area of the circle shown = $\frac{3}{4} \times \pi r^2$
The radius = 2 cm
So area of shape shown = $\frac{3}{4} \times \pi \times 2^2 = 3\pi \text{ cm}^2$

Indices and roots

- $7 \times 7 \times 7 \times 7 = 7^4$
 - $\frac{1}{5 \times 5 \times 5} = \frac{1}{5^3} = 5^{-3}$
- $2^4 = 2 \times 2 \times 2 \times 2 = 16$
 - $10^{-2} = \frac{1}{10^2} = \frac{1}{100}$

3 $2^3 = 2 \times 2 \times 2 = 8$
 $3^{-2} = \frac{1}{9}$
 $\sqrt[3]{27} = 3$
 $\sqrt{25} = 5$ or -5

Assuming the square root of 25 is positive, the answer is:

$3^{-2}, \sqrt[3]{27}, \sqrt{25}, 2^3$

If it were negative, the answer would be:

$\sqrt{25}, 3^{-2}, \sqrt[3]{27}, 2^3$

4 $\frac{9^5}{9^3 \times 9^2} = \frac{9^5}{9^5} = 1$

Standard form

- 1 2750
- 2 1.5×10^8
- 3 Move the decimal point three places to the right to give 6.42×10^{-3}
- 4 $(1.4 \times 10^{-5}) \times 20 = (2.8 \times 10^{-5}) \times 2 \times 10 = 2.8 \times 10^{-4}$ km

Listing strategies

- 1 259, 295, 529, 592, 925, 952

2 a

		4-sided spinner			
		0	1	2	3
3-sided spinner	1	1	2	3	4
	2	2	3	4	5
	3	3	4	5	6

b 3

3

		Gift tag				
		1	2	3	4	5
Wrapping paper	Red	Red, 1	Red, 2	Red, 3	Red, 4	Red, 5
	Blue	Blue, 1	Blue, 2	Blue, 3	Blue, 4	Blue, 5
	Green	Green, 1	Green, 2	Green, 3	Green, 4	Green, 5

Terrence can use 15 different combinations of wrapping paper and gift tags.

- 4 spj; spi; sfj; sfi ; bpj; bpi; bfj; bfi

Algebra

Understanding expressions, equations, formulae and identities

- 1 a identity b equation c expression
- 2 a Equation, because it has an equals sign and can be solved.
 b Formula, because it has letter terms, an equals sign and the values of the letters can vary.
 c Expression, because it has letter terms and no equals sign.
 d Formula, because it has letter terms, an equals sign and the values of the letters can vary.
- 3 a Any of: $2x + 10$ or $10x + 2$ or $x + 210$ or $x + 102$
 b Any of: $2x = 10$ or $10x = 2$

Simplifying expressions

- 1 $8x$
- 2 a $6a \times 8a = (6 \times 8) \times (a \times a) = 48 \times a^2 = 48a^2$
 b $2p \times 3p \times 5p = (2 \times 3 \times 5) \times (p \times p \times p) = 30 \times p^3 = 30p^3$
- 3 $35yz \div 7z = (35 \div 7) \times (yz \div z) = 5 \times y = 5y$
- 4 $\frac{32uv}{4v} = \frac{32}{4} \times \frac{uv}{v} = 8 \times u = 8u$

Collecting like terms

- 1 a $7m + 6n - 4m - 2n = (7 - 4)m + (6 - 2)n = 3m + 4n$
 b $9q - 5r - 12q + 3r = (9 - 12)q + (3 - 5)r = -3q - 2r$
- 2 a $11a + 5b - 10a + 8b = (11 - 10)a + (5 + 8)b = a + 13b$
 b $6c - 4d - 7c + 5d = (6 - 7)c + (5 - 4)d = -c + d$
 c $m + m + m + n \times n = 3m + n^2$

- 3 a $9p^3 + p - 4p^3 = (9 - 4)p^3 + p = 5p^3 + p$
 b $12 - 5x^2 + 3x - 2x^2 = 12 - (5 + 2)x^2 + 3x = -7x^2 + 3x + 12$
 c $m + m + m + n \times n = 3m + n^2$
- 4 $3\sqrt{5} - f - 8\sqrt{5} + 2f = (3 - 8)\sqrt{5} + (2 - 1)f = -5\sqrt{5} + f$

Using indices

- 1 a $p^3 \times p = p^{(3+1)} = p^4$
 b $4y^2 \times 3y^3 = (4 \times 3) \times y^{(2+3)} = 12 \times y^5 = 12y^5$
 c $2a^4b \times 5ab^2 = (2 \times 5) \times a^{(4+1)} \times b^{(1+2)} = 10 \times a^5 \times b^3 = 10a^5b^3$
- 2 a $q^{-2} \times q^{-4} = q^{(-2-4)} = q^{-6}$
 b $(u^{-3})^2 = u^{(-3) \times 2} = u^{-6}$
 c $x^{-1} \times x = x^{-1} \times x^1 = x^{(-1+1)} = x^0 = 1$
- 3 a $b^4 \div b^3 = b^{(4-3)} = b^1 = b$
 b $\frac{f^5}{f^2} = f^{(5-2)} = f^3$
 c $\frac{xy^3}{x^2y} = x^{(1-2)} \times y^{(3-1)} = x^{-1} \times y^2 = \frac{1}{x} \times y^2 = \frac{y^2}{x}$
- 4 Let the first box = x and the second box = y
 $(xm^3)^y = x^y m^{3y} = 8m^9$
 comparing terms, $3y = 9$
 $y = 3$
 Substitute in the y value: $(xm^3)^3 = 8m^9$
 $x^3 = 8$
 $x = \sqrt[3]{8} = 2$
 Therefore, the completed expression is $(2m^3)^3$

Expanding brackets

- 1 a $4(m + 3) = (4 \times m) + (4 \times 3) = 4m + 12$
 b $2(p - 1) = (2 \times p) + (2 \times -1) = 2p - 2$
 c $10(3x - 5) = (10 \times 3)x + (10 \times -5) = 30x - 50$
- 2 a $3(m + 2) + 5(m + 1) = 3m + 6 + 5m + 5 = 8m + 11$
 b $6(x - 1) - 2(x - 4) = 6x - 6 - 2x + 8 = 4x + 2$
- 3 a $(y + 3)(y + 7) = y^2 + 7y + 3y + 21 = y^2 + 10y + 21$
 b $(b + 2)(b - 4) = b^2 - 4b + 2b - 8 = b^2 - 2b - 8$
 c $(x - 4)(x - 6) = x^2 - 6x - 4x + 24 = x^2 - 10x + 24$
- 4 a $(q + 1)^2 = (q + 1)(q + 1) = q^2 + q + q + 1 = q^2 + 2q + 1$
 b $(c - 3)^2 = (c - 3)(c - 3) = c^2 - 3c - 3c + 9 = c^2 - 6c + 9$
 c $(4m - n)(3m + n) = 12m^2 + 4mn - 3mn - n^2 = 12m^2 + mn - n^2$

Factorising

- 1 Divide the expression by the highest common factor (HCF) of both terms to find the bracket, and then place the HCF outside of the bracket to give the full factorisation.
 a $(4x + 8) \div 4 = x + 2$
 factorisation: $4(x + 2)$
 b $(3d - 15) \div 3 = d - 5$
 factorisation: $3(d - 5)$
 c $(8y - 12) \div 4 = 2y - 3$
 factorisation: $4(2y - 3)$
- 2 Divide the expression by the common term to find the bracket, and then place the common term outside of the bracket to give the full factorisation.
 a $(q^2 + q) \div q = q + 1$
 factorisation: $q(q + 1)$
 b $(a^2 + 6a) \div a = a + 6$
 factorisation: $a(a + 6)$
 c $(10z^2 + 15z) \div 5z = (2z + 3)$
 factorisation: $5z(2z + 3)$
- 3 Find which factors of the number term add together to give the coefficient of the x term.
 a $12 = 3 \times 4$
 $7 = 3 + 4$
 factorisation: $(x + 3)(x + 4)$

b $-16 = (-2) \times 8$
 $6 = -2 + 8$
 factorisation: $(x - 2)(x + 8)$

c $24 = (-6) \times (-4)$
 $-10 = (-6) + (-4)$
 factorisation: $(a - 6)(a - 4)$

4 a Write $y^2 - 4$ in the form of $a^2 - b^2$:
 $y^2 - 2^2$
 Using the formula for the difference of two squares, the factorisation is
 $(y + 2)(y - 2)$

b Write $x^2 - 9$ in the form of $a^2 - b^2$:
 $x^2 - 3^2$
 Using the formula for the difference of two squares, the factorisation is
 $(x + 3)(x - 3)$

c Write $p^2 - 100$ in the form of $a^2 - b^2$:
 $p^2 - 10^2$
 Using the formula for the difference of two squares, the factorisation is
 $(p + 10)(p - 10)$

Substituting into expressions

1 $4x + 5y = 4 \times 3 + 5 \times (-2) = 12 - 10 = 2$

2 $s = ut + \frac{1}{2}at^2$
 $= 12 \times 2 + \frac{1}{2} \times 10 \times 2^2$
 $= 12 \times 2 + \frac{1}{2} \times 40$
 $= 24 + 20$
 $s = 44$

3 a $f = 3c - 2(c - d)$
 $= 3 \times 7 - 2 \times (7 - (-5))$
 $= 21 - 2 \times (12)$
 $= 21 - 24$
 $f = -3$

b $f = -c(d^2 - 3c)$
 $= -7 \times ((-5)^2 - 3 \times 7)$
 $= -7 \times (25 - 21)$
 $= -7 \times 4$
 $f = -28$

c $f^2 = 7c - 3d$
 $= 7 \times 7 - 3 \times (-5)$
 $= 49 + 15$
 $= 64$
 $f = \sqrt{64}$
 $f = \pm 8$

4 $\frac{p(q-3)}{r} = \frac{6(7-3)}{-8}$
 $= \frac{6 \times 4}{-8}$
 $= \frac{24}{-8}$
 $= -3$

Writing expressions

1 a $n + 3$ **b** $(n \times 2) - 9 = 2n - 9$

2 a $x + y$ **b** $5 \times x = 5x$

c $(12 \times x) + (11 \times y) = 12x + 11y$

3 $2 \times 9p + 2(5p + 2) = 18p + 10p + 4 = 28p + 4$

4 The area of the rectangle is given by height \times length, which is $s \times (5s + 1) = s(5s + 1)$.

Solving linear equations

1 a $x = 12 - 5$
 $x = 7$

b $x = 10 + 3$
 $x = 13$

c $x = \frac{20}{4}$
 $x = 5$

d $x = 6 \times 3$
 $x = 18$

2 a $2x + 3 = 15$
 $2x = 12$
 $x = 6$

b $3x - 5 = 16$
 $3x = 21$
 $x = 7$

c $\frac{x}{5} + 3 = 8$
 $\frac{x}{5} = 5$
 $x = 25$

d $7 - 2x = 1$
 $7 = 2x + 1$
 $6 = 2x$
 $x = 3$

3 a $3(x + 9) = 30$
 $3x + 27 = 30$
 $3x = 3$
 $x = 1$

b $5(p - 2) = 10$
 $5p - 10 = 10$
 $5p = 20$
 $p = 4$

c $2(10 - 3m) = 8$
 $20 - 6m = 8$
 $12 = 6m$
 $m = 2$

d $4(6 - 2q) + 10 = 2$
 $24 - 8q + 10 = 2$
 $32 = 8q$
 $q = 4$

4 a $4x - 6 = x + 9$
 $3x - 6 = 9$
 $3x = 15$
 $x = 5$

b $2y + 5 = 4y - 3$
 $5 = 2y - 3$
 $8 = 2y$
 $y = 4$

c $4(2x + 3) = 11x + 3$
 $8x + 12 = 11x + 3$
 $9 = 3x$
 $x = 3$

d $3(n + 4) = 2(2n + 3)$
 $3n + 12 = 4n + 6$
 $12 = n + 6$
 $n = 6$

Writing linear equations

1 Sum of the angles in a triangle are 180°
 $(2x + 3) + 81 + (3x - 4) = 180$
 $5x + 80 = 180$
 $5x = 100$
 $x = 20$

2 Let Jamie's age = x years. Sophie's age = $\frac{x}{2}$
 $x + \frac{x}{2} = 18$
 $\frac{3x}{2} = 18$
 $3x = 36$
 $x = 12$

Jamie is 12 years old

3 Let width = x so length = $x + 3$
 Perimeter = $2x + 2(x + 3)$
 $46 = 4x + 6$
 $x = 10$


Length = 10cm and width = 13cm

Area = $10 \times 13 = 130\text{cm}^2$


4 Opposite angles are equal so $3x + 10 = 5x - 20$
 $30 = 2x$ giving $x = 15$
 Also $3x + 10 + 7x + 5y = 180$
 $10x + 10 + 5y = 180$
 Now $x = 15$ so $150 + 10 + 5y = 180$
 Solving this gives $y = 4$

Linear inequalities


1 a $-1, 0, 1, 2, 3, 4, 5$




2 a $4x > 20$
 $x > 5$



b $3x - 8 \leq 13$
 $3x \leq 21$
 $x \leq 7$



c $2(x - 3) < 10$
 $2x - 6 < 10$
 $2x < 16$
 $x < 8$



3 a $2 \leq 3x + 5$
 $-3 \leq 3x$
 $-1 \leq x$
 $3x + 5 < 11$
 $3x < 6$
 $x < 2$
 Hence $-1 \leq x < 2$

b $-4 > 5x + 6$
 $-10 < 5x$
 $-2 < x$
 $5x + 6 \leq 6$
 $5x \leq 0$
 $x \leq 0$
 Hence $-2 < x \leq 0$

4 $n + n + 3 < 15$
 $2n + 3 < 15$
 $2n < 12$
 $n < 6$
 Possible integer values of $n = 1, 2, 3, 4, 5$

Formulae

1 a $t = (40 \times 2) + 20 = 100$ minutes = 1 hour 40 minutes
 b $t = (40 \times 1.5) + 20 = 80$ minutes = 1 hour 20 minutes
 The chicken should be put in the oven 1 hour and 20 minutes earlier than 1.30 pm, which is a time of 12.10 pm.

2 a $C = l + kn$
 b $C = 90 + 6.5 \times 3$
 $C = \pounds 109.50$

3 a $p = \frac{qs}{3}$
 $3p = qs$
 $q = \frac{3p}{s}$
 b $p = \frac{q}{r} + t$
 $p - t = \frac{q}{r}$
 $q = rp - rt$ or $r(p - t)$

c $p = 3(q + r)$
 $\frac{p}{3} = q + r$
 $q = \frac{p}{3} - r = \frac{p - 3r}{3}$

d $p = \sqrt{2q}$
 $p^2 = 2q$
 $q = \frac{p^2}{2}$

Linear sequences

- 1 a The term in position 1 is $1 \times 5 + 1 = 6$
 The term in position 2 is $2 \times 5 + 1 = 11$
 The term in position 3 is $3 \times 5 + 1 = 16$
 The term in position 4 is $4 \times 5 + 1 = 21$
 b The term in position 50 is $50 \times 5 + 1 = 251$
- 2 a Each pattern has 2 more dots than the last, so pattern 7 will have 8 more dots than pattern 3. Pattern 7 will have 19 dots.
 b No, Rachel is not correct, because the number of triangles is not the pattern number multiplied by 2. Instead, it is the pattern number plus 2, so there will be 6 triangles in pattern 4.
- 3 a Common difference = 11, so 11n is in the sequence.
 When $n = 1$:
 $11n = 11$, but the 1st term is 3.
 $3 = 11n - 8$
 So the expression for the sequence is $11n - 8$
 b Assume 100 is in the sequence. Then:
 $11n - 8 = 100$
 $11n = 108$
 $n = 108 \div 11 = 9$ remainder 9
 But n must be a whole number, and it is not; so 100 is not in this sequence.

Non-linear sequences

- 1 a Rule is multiply by 2.
 $8 \times 2 = 16$
 $16 \times 2 = 32$
 So terms are 16, 32.
 b Rule is divide by 10.
 $1 \div 10 = 0.1$
 $0.1 \div 10 = 0.01$
 So terms are 0.1, 0.01
 c Rule is multiply by -2 .
 $-12 \times -2 = 24$
 $24 \times -2 = -48$
 So terms are 24, -48 .
 d They involve multiplying and dividing, not adding and subtracting, so they are geometric.
- 2 a Next term = $6 + 9 = 15$
 b 5th term = $6 + 9 = 15$
 6th term = $9 + 15 = 24$
 7th term = $15 + 24 = 39$
 8th term = $24 + 39 = 63$
 9th term = $39 + 63 = 102$
 The 9th term is the first term in the sequence over 100

3 a

Day	Mon	Tue	Wed	Thu	Fri
Number of ladybirds	2	$8 (= 2 \times 4)$	$32 (= 8 \times 4)$	$32 \times 4 = 128$	$128 \times 4 = 512$

The gardener is correct. There will be more than 500 ladybirds.

b Saturday, because $512 \times 4 = 2048$.

4 a First term: $\frac{1}{2} \times 1^2 = \frac{1}{2}$

Second term: $\frac{1}{2} \times 2^2 = 2$

Third term: $\frac{1}{2} \times 3^2 = \frac{9}{2} = 4\frac{1}{2}$

b If 32 is in the sequence, then:

$$\frac{1}{2}n^2 = 32$$

$$n^2 = 64$$

$$n = 8$$

This gives n as a whole number, 8, so 32 is the 8th term in the sequence.

Show that...

1 LHS = $2x + 1$; RHS = $2x + 1$; LHS = RHS. Therefore,

$$2\left(x + \frac{1}{2}\right) \equiv x + x + 1$$

2 LHS = $x^2 - 25 + 9 = x^2 - 16$; RHS = $x^2 - 16$

3 Let the three consecutive numbers be $n, n + 1$ and $n + 2$.

$n + n + 1 + n + 2 = 3n + 3 = 3(n+1)$. Therefore, the sum of three consecutive numbers is a multiple of 3.

4 a Width of pond = $x - y + x + x + x - y = 4x - 2y$

Length of pond = $4x$

$$\text{Perimeter} = 4x - 2y + 4x - 2y + 4x + 4x = 16x - 4y$$

b Yes Sanjit is correct, because $16x - 4y = 4(4x - y)$, showing that when x and y are whole numbers, the perimeter is always a multiple of 4.

Number machines

1 a when $x = 3, y = 3 \times 4 - 1 = 11$

b when $y = 23, 4x - 1 = 23$, therefore $x = (23 + 1) \div 4 = 6$

c To get y you multiply x by 4 and subtract 1, so $y = 4x - 1$

2


x	Operations	y
-2	$(-2) \times 2 + 3$	-1
0	$0 \times 2 + 3$	3
3	$(9 - 3) \div 2$	9

3

x	Operations	y
-2	$(-2) \div 2 + 1$	0
1	$(1) \div 2 + 1$	$1\frac{1}{2}$
8	$(5 - 1) \times 2$	5


Coordinates and midpoints

1 a 1 along the x -axis and 2 up the y -axis: (1, 2)



c $4\frac{1}{2}$ along the x -axis and 1 up the y -axis: $(4\frac{1}{2}, 1)$

2 a 1 along the x -axis and -4 'up' the y -axis: (1, -4)



S = (6, -2) to make a parallelogram

3 a x coordinate = $\frac{4 + (-2)}{2} = 1$

y coordinate = $\frac{5 + 1}{2} = 3$

Midpoint of XY is (1, 3).


b Midpoint of $XZ = \left(\frac{4+4}{2}, \frac{5+(-4)}{2}\right) = (4, \frac{1}{2})$

c Midpoint of $YZ = \left(\frac{(-2)+4}{2}, \frac{1+(-4)}{2}\right) = (1, -1\frac{1}{2})$


Straight line graphs

1 a For $y = 2x + 3$

x	-1	0	1	2
Operations	$2 \times (-1) + 3$	$2 \times (0) + 3$	$2 \times (1) + 3$	$2 \times (2) + 3$
y	1	3	5	7



2 a and b



a For $y = 3x$

x	-2	-1	0	1	2
Operations	$3 \times (-2)$	$3 \times (-1)$	$3 \times (0)$	$3 \times (1)$	$3 \times (2)$
y	-6	-3	0	3	6

b Rearrange the equation to give $y = -2x - \frac{1}{2}$

x	Operations	y
-3	$-2 \times (-3) - \frac{1}{2}$	$5\frac{1}{2}$
-2	$-2 \times (-2) - \frac{1}{2}$	$3\frac{1}{2}$
-1	$-2 \times (-1) - \frac{1}{2}$	$1\frac{1}{2}$
0	$-2 \times (0) - \frac{1}{2}$	$-\frac{1}{2}$
1	$-2 \times (1) - \frac{1}{2}$	$-2\frac{1}{2}$
2	$-2 \times (2) - \frac{1}{2}$	$-4\frac{1}{2}$

c Line C goes through points (0, -3), (1, 1) and (2, 5)
The y intercept is -3.

The gradient is $\frac{\text{difference in } y \text{ coordinates}}{\text{difference in } x \text{ coordinates}} = \frac{5 - 1}{2 - 1} = 4$

The equation of line C is $y = 4x - 3$.

3 a B and C, because they have the same gradient of 2.

b A and B, because they both have a y-intercept at (0, 1).

4 The gradient is $\frac{2 - (-6)}{3 - (-1)} = \frac{8}{4} = 2$

Using point (3, 2) and gradient $m = 2$:


$$y = 2x + c$$

$$2 = 2 \times 3 + c$$

$$c = -4$$


Equation of the line is $y = 2x - 4$

5 a




Draw a vertical line up from $x = 3$ to the graph, and then a horizontal line to the y-axis to read off the result: $y = 2$.

b




Draw a horizontal line across from $y = -2$ to the graph, and then a vertical line up to the x-axis to read off the result: $x = 1$.

6 a



Draw a horizontal line across from $y = -2$ to the graph, and then a vertical line up to the x-axis to read off the result: $x = 2.5$.


b and c



b Draw a vertical line up from $x = -0.5$ to the graph, and then a vertical line across to the y-axis to read off the result: $y = 4$.


c Reading up from $x = 1.2$ and then across to y axis gives $y = 0.6$. Any value from 0.6 to 0.75 is acceptable.

7 a and b



b Where the graphs cross, draw a vertical line up to the x-axis. It meets the axis two thirds of the way between $x = -1$ and $x = 0$, so the solution is approximately $x = -0.67$. Any value from -0.6 to -0.7 is acceptable.

8



Compare the equations $y = 4x - 3$ and $4x - 3 = 2$.
 y has been replaced with 2, so add line $y = 2$ to the graph.
 The intersection point of the two graphs gives the solution to the equation $4x - 3 = 2$
 $x = 1.25$. Any answer between 1.2 and 1.3 is acceptable.

Solving simultaneous equations

- 1 a Substituting $y = 2x$ into the first equation gives
 $3x + 2x = 15$
 $5x = 15$
 $x = 3$

When $x = 3$, $y = 2 \times 3 = 6$

- b $3x = 12$
 $x = 4$

Substituting $x = 4$ into the first equation gives
 $8 + y = 9$
 $y = 1$

- c $5x = 10$
 $x = 2$

Substituting $x = 2$ into the first equation gives
 $6 + y = 4$
 $y = -2$

- 2 a $2x + 2y = 14$ (1)
 $3x + y = 11$ (2)
 $(2) \times 2 \quad 6x + 2y = 22$
 $(3) - (1) \quad 4x = 8$
 $x = 2$

Substitute into (1) $4 + 2y = 14$
 $2y = 10$
 $y = 5$

Solution is $x = 2, y = 5$

- b $4x - 2y = 2$ (1)
 $2x - 3y = 7$ (2)
 $(2) \times 2 \quad 4x - 6y = 14$ (3)
 $(3) - (1) \quad -4y = 12$
 $y = -3$

Substitute into (1) $4x + 6 = 2$
 $4x = -4$
 $x = -1$

Solution is $x = -1, y = -3$

- c $2x + 3y = 20$ (1)
 $3x + 2y = 15$ (2)
 $(1) \times 2 \quad 4x + 6y = 40$ (3)
 $(2) \times 3 \quad 9x + 6y = 45$ (4)
 $(4) - (3) \quad 5x = 5$
 $x = 1$

Substituting $x = 1$ into equation (1) $2 + 3y = 20$
 $3y = 18$
 $y = 6$

Solution is $x = 1, y = 6$


3 a $x + y = 2$

x	0	2
y	2	0

$2x - y = 1$


x	0	$\frac{1}{2}$
y	-1	0

b



From the intersection of the two lines, $x = 1, y = 1$.

c



From the intersection of the two lines, $x = 2, y = 3$.


Quadratic graphs

- 1 a C and D – they are straight lines so they are linear.
 b E – it is quadratic, with a positive multiplier for x^2 , and is symmetrical about the origin.
 c A – it is quadratic, with a negative multiplier for x^2 , and is symmetrical about the origin.
 d D – the x coordinates are all different, but all the y coordinates on this line are 1.
 e B – it is the same as E except that it has been moved 1 unit up the y -axis.

2 a

x	-3	-2	-1	0	1	2	3
y	6	1	-2	-3	-2	1	6


b




c $x = 0$

d $(0, -3)$

3 a



b $x = -1$ c $(-1, -2)$




b $x = -2$ or $x = 1$


5 $x = -3$ or $x = 1$

6 a

x	-3	-2	-1	0
$x^2 + 3x + 1$	$(-3)^2 + 3 \times (-3) + 1$	$(-2)^2 + 3 \times (-2) + 1$	$(-1)^2 + 3 \times (-1) + 1$	$(0)^2 + 3 \times (0) + 1$
y	1	-1	-1	1



b Compare the equations $y = x^2 + 3x + 1$ and $x^2 + 3x + 1 = -1$. y has been replaced with -1 , so the solutions to the equation $x^2 + 3x + 1 = -1$ are where $y = -1$.



$x = -2$ and $x = -1$

c Compare the equations $y = x^2 + 3x + 1$ and $x^2 + 3x + 1 = 0$. y has been replaced with 0 , so the solutions to the equation $x^2 + 3x + 1 = 0$ are where the graph crosses the x -axis.

$x = -2.6$ and $x = -0.38$ (any answer close to -0.4 is acceptable)

Solving quadratic equations


- | | |
|---|--|
| 1 a $x(x + 6) = 0$
$x = 0$ or -6 | 3 a $(x + 3)(x + 2) = 0$
$x = -2$ or -3 |
| b $y(y - 11) = 0$
$y = 0$ or 11 | b $(x + 5)(x - 2) = 0$
$x = -5$ or 2 |
| c $3d(d - 3) = 0$
$d = 0$ or 3 | c $(x - 7)(x - 2) = 0$
$x = 2$ or 7 |
| 2 a $(x + 4)(x - 4) = 0$
$x = 4$ or -4 | 4 a $0 = x(x - 3)$
$x = 0$ or 3 |
| b $(a + 9)(a - 9) = 0$
$a = 9$ or -9 | b $0 = (x - 5)(x + 5)$
$x = 5$ or -5 |
| c $(z - 10)(z + 10) = 0$
$z = 10$ or -10 | c $0 = (x + 6)(x - 3)$
$x = -6$ or 3 |

Cubic and reciprocal graphs

- 1 a A, C and D – they are not continuous curves with two turning points (s-shaped curves).
 b B – it has two turning points and has rotational symmetry about the origin.
 c E – it has two turning points and is a reflection of B, raised up one unit on the y -axis.
 d D – this is the form for a reciprocal graph.

2 a


x	-2	-1	0	1	2
y	-4	3	4	5	12



3 a cubic b $(0, -8)$ c $(2, 0)$


4 a

x	-3	-2	-1	$-\frac{1}{2}$	$\frac{1}{2}$	1	2	3
y	$\frac{2}{3}$	$\frac{1}{2}$	0	-1	3	2	$1\frac{1}{2}$	$1\frac{1}{3}$



Drawing and interpreting real-life graphs


- 1 a The initial charge is the value at 0 miles, where the graph cuts the y -axis (the y -intercept): \$3.
 b The charge per mile is given by the gradient of the graph.
 Gradient = $\frac{\text{difference in } y \text{ coordinates}}{\text{difference in } x \text{ coordinates}} = \frac{23 - 3}{8 - 0} = \frac{20}{8} = 2.5$
 Charge per mile is \$2.50



- d From the conversion graph, $\text{£}15 = \text{\$}18$.
 From the graph of the cost of a taxi in New York, \$18 allows you to travel 6 miles.
 2 a You can read this from the highest point of the graph: 16 m.
 b Read from the highest point down to the value on the horizontal axis: 4 seconds.
 c This is when the ball reaches 0 height for the second time: 8 seconds.
 d Read from 12 on the vertical axis across to the curve: 2 seconds and 6 seconds.

- 3 **a** This is the highest value on the vertical axis: 10 m/s.
b The cyclist is travelling at a constant speed of 10 m/s.
c The cyclist is decelerating, so it will be a negative value.
 Acceleration = $\frac{-10}{20} = -\frac{1}{2}$ m/s²

- d** The speed returns to 0: the cyclist stops.
- 4 **a** Assuming that water pours into the container at a constant rate,
 A: depth goes up increasingly slowly as the container widens, so 2.
 B: depth rises steadily and fairly slowly in a broad container of consistent diameter, so 4.
 C: depth rises quickly at first, then more slowly as the container widens, then more quickly as it gets narrow towards the top, so 3.
 D: depth rises steadily and quickly in a narrow container of consistent diameter, so 1.



Ratio, proportion and rates of change

Units of measure

- 1 **a** $4 \text{ (m)} \times 100 = 400 \text{ cm}$
b $500 \text{ (g)} \div 1000 = 5 \text{ kg}$
c $1.5 \text{ (l)} \times 1000 = 1500 \text{ ml}$
d $8250 \text{ (m)} \div 1000 = 8.25 \text{ km}$
- 2 $6 \text{ (litres)} \times 1000 = 6000 \text{ ml}$
 $6000 - 3500 = 2500$
 Sally had **2500 ml** of lemonade left.
- 3 **a** Luke: 240 seconds; Adam: $3 \times 60 + 47 = 227$ seconds.
 Adam arrived first.
or Luke: $240 \div 60 = 4$ minutes; Adam: 3 minutes 47 seconds. Adam arrived first.
- b** $4 \text{ minutes} - 3 \text{ minutes } 47 \text{ seconds} = 13 \text{ seconds}$.
 Adam waited **13 seconds** for Luke to arrive at school.
- 4 $\text{Ben} = 1.25 \text{ m} = 3.2 + 0.8 \text{ feet} = 4 \text{ feet}$. Tom is taller.
or $\text{Tom} = 4.8 \text{ feet} = 3.2 + 1.6 \text{ feet} = 1 + 0.5 \text{ metres} = 1.5 \text{ metres}$. Tom is taller.

Ratio

- 1 12 stories and 8 colouring
 Ratio of story: colouring = $12 : 8 = 3 : 2$
- 2 Total parts = $1 + 2 = 3$
 1 part = $\frac{15}{3} = 5$
 Spent = $2 \times 5 = \text{£}10$
- 3 Total parts = $1 + 2 + 3 = 6$
 1 part = $\frac{60}{6} = 10$
 Amount given to charity = $3 \times 10 = \text{£}30$
- 4 Ratio of blue to yellow required is $3 : 7$.
 There are $3 + 7 = 10$ parts. He needs to make 5 litres.
 $10 \text{ parts} = 5000 \text{ ml}$
 1 part = 500 ml
 Phil needs $3 \times 500 \text{ ml} = 1500 \text{ ml} = 1.5 \text{ litres}$ of blue paint.
 He has 2 litres of blue paint.
 Phil needs $7 \times 500 \text{ ml} = 3500 \text{ ml} = 3.5 \text{ litres}$ of yellow paint.

He has 3 litres of yellow paint.

Phil has enough blue paint, but does not have enough yellow paint.

Scale diagrams and maps

- 1 1 cm on the map is **10000 cm** in real life.
 This means 1 cm on the map is **100 m** in real life.
- 2 1 cm on the map represents 50 m in real life.
 $3 \times 50 \text{ m} = 150 \text{ m}$, so the bus stop is **3 cm** from the village shop on the map.
- 3 Measure the distance between the trees on the diagram = 5 cm
 1 cm on the diagram represents 4 m in real life.
 $5 \times 4 = 20$
 The trees are **20 m** apart.
- 4 A scale of $1 : 400$ means 1 cm on the model represents 4 m (= 400 cm) in real life.
 $96 \div 4 = 24$
 The scale model is **24 cm** tall.

Fractions, percentages and proportion

- 1 $1 + 3 = 4$ parts so Bess receives $\frac{3}{4}$
- 2 **a** $1 : 3 : 6$
b $1 + 3 + 6 = 10$ items in the basket
 Fruit = $\frac{3}{10}$
c Tins = $\frac{6}{10} = 60\%$
- 3 $\frac{50}{4000} = \frac{1}{80}$
- 4 Total parts = $3 + 8 + 14 = 25$
 $\frac{8}{25} = \frac{32}{100} = 32\%$

Direct proportion

- 1 **a** One ticket costs $\text{£}80 \div 5 = \text{£}16$
b Nine tickets cost $9 \times \text{£}16 = \text{£}144$
- 2 **a** Read up from 6 packs on the horizontal axis, to the line, then across to the vertical axis to find the cost: $\text{£}1.20$
b There are 10 pencils in a pack, so 1 pencil is 0.1 of a pack. Reading off the graph using this value, the price is 2p.
c It is a straight-line graph; the graph passes through the origin (0, 0).
- 3 **a** Sally needs to make $28 \div 4 = 7$ lots of the recipe.
 She will need $1 \times 7 = 7$ teaspoons of turmeric,
 $2 \times 7 = 14$ teaspoons of chilli powder and
 $2\frac{1}{2} \times 7 = 17\frac{1}{2}$ teaspoons of cumin.
b Sally has 75 g of chilli powder. That is $75 \div 3 = 25$ teaspoons.
 Sally needs 14 teaspoons to make the curry for her class.
 She does have enough.

Inverse proportion

- 1 **a** Start from 5 on the x -axis, read up to the graph, then left to the scale on the y -axis.
 5 winners will each get **£400**.
b Start from 200 on the y -axis, read right to the graph, then down to the scale on the x -axis. 10 winners each get $\text{£}200$, so there are **9** other winners.
c Multiply the number of winners by the amount each one gets to find the total prize money. e.g. $5 \times \text{£}400 = \text{£}2000$.
d Compare 2 points on the graph. (Use your answers to parts **a** and **b**.) If the number of winners goes up, the prize money they each receive goes down. The prize money and the number of winners are in inverse proportion.
- 2 **a** The total time needed to decorate the room is $3 \times 2 = 6$ hours.
b $6 \text{ hours} \div 12 \text{ people} = 0.5 \text{ hours} = \text{30 minutes}$ (or 0.5 hours).

- 3 a y decreases from 1 to $\frac{1}{8}$ as x increases from 1 to 8
(when $x = 1, y = 1$; when $x = 8, y = \frac{1}{8}$)
b x is inversely proportional to y , because as x increases, y decreases.
- 4 The printer can print $240 \div 4 = 60$ pages per minute.
 $600 \div 60 = 10$, so it would take **10 minutes** to print the larger document.

Working with percentages

- 1 $125 - 75 = 50$
 $\frac{50}{125} \times 100 = 40\%$
- 2 a Amount of increase = $24 - 15 = \text{£}9$ million
 $\frac{9}{15} \times 100 = 60\%$ increase in sales for Company X
b $125\% = \text{£}35$ million
 $35 \div 125 \times 100 = \text{£}28$ million sales in 2006.
- 3 a $\frac{2}{100} \times 265 = 5.30$ and $3 \times 5.30 = 15.90$
 $15.90 + 265 = \text{£}280.90$
b $265 \times (1.02)^3 = \text{£}281.22$


Compound units

- 1 $\frac{3 \text{ km}}{\text{minute}} = \frac{3000 \text{ m}}{\text{minute}} = \frac{3000 \text{ m}}{60 \text{ seconds}} = 50 \text{ m/s}$
- 2 $20 \div 5 = 4$ minutes to fill the tank.
- 3 Pressure = $\frac{300}{0.05} = 6000$ Newtons/m² (or 6000 N/m²)
- 4 On Saturday Sami drove $4 \times 50 = 200$ miles; on Sunday Sami drove $356 \div 8 \times 5 = 222.5$ miles. Sami drove further on Sunday.

Geometry and measures

Measuring and drawing angles

- 1 a 123°
b 42°
c 331°



- 3 a $100^\circ, 120^\circ, 140^\circ, 160^\circ$
b First angle + second angle = 87° . This means both angles are less than 87° , and so they both must be acute.

Using the properties of angles


- 1 a Angle $ACB = 52^\circ$ (Angles in a triangle add up to 180°)
 $x = 128^\circ$ (Angles on a straight line add up to 180°)
b Angle $ADC = 86^\circ$ (Angles in a quadrilateral add up to 360°)
 $x = 94^\circ$ (Angles on a straight line add up to 180°)
- 2 a Angle $BED = 39^\circ$ (Alternate angles are equal)
Angle $BDE = 39^\circ$ (Base angles in an isosceles triangle are equal)
 $x = 102^\circ$ (Angles in a triangle add up to 180°)
b Angle $DCF = 98^\circ$ (Vertically opposite angles are equal)
 $x = 98^\circ$ (Corresponding angles are equal)
- 3 Angle $CFG = 62^\circ$ (Co-interior angles add up to 180°)
 $x = 66^\circ$ (Angles on a straight line add up to 180°)
- 4 $x + 40 + 3x + 5x - 40 = 180^\circ$
 $9x = 180^\circ$
 $x = 20^\circ$
Angle $BAC = x + 40 = 20 + 40 = 60^\circ$
Angle $ACB = 3x = 3 \times 20 = 60^\circ$
Angle $ABC = 5x - 40 = 5 \times 20 - 40 = 60^\circ$
Triangle ABC has equal angles of 60° . Therefore, it is an equilateral triangle.

Using the properties of polygons


- 1 a $180^\circ \times (6 - 2) = 720^\circ$
b $720^\circ \div 6 = 120^\circ$
c $180^\circ - 120^\circ = 60^\circ$ or $360^\circ \div 6 = 60^\circ$
- 2 a It is an octagon because it has eight sides.
b All angles are equal; all sides are equal.
c $180^\circ \times (8 - 2) = 1080^\circ$
 $1080^\circ \div 8 = 135^\circ$
or $180^\circ - (360^\circ \div 8) = 135^\circ$
- 3 exterior angle = $180^\circ - 144^\circ = 36^\circ$
number of sides = $360 \div 36 = 10$
Therefore it is a decagon.

Using bearings


- 1 a This is the angle measured clockwise from North at A: 065° .
b $180 - 138 = 42^\circ$ This is the acute angle at C.
Bearing of B from $C = 360 - 42^\circ = 318^\circ$
c $180 - 65 = 115^\circ$ This is the angle between the north line at B and AB , measured anticlockwise.
Bearing of A from $B = 360 - 115 = 245^\circ$
- 2 To find a reciprocal bearing, subtract 180 from the original bearing (or add 180 to it).
The bearing of O from $X = 276 - 180 = 096^\circ$
- 3 a Draw a North line at P , then join P to Q and measure the angle between the North line and this line: 060° (any value 058° to 062° accepted).



Properties of 2D shapes





- b trapezium
c one pair of parallel sides



- b 4
 c square
 d Two from: all sides equal in length; all angles are 90° ; diagonals are equal; diagonals bisect each other at 90° .

- 3 a rectangle, rhombus
 b




Congruent shapes


- D and F are exactly the same as A – they are the same size and shape (it doesn't matter that they are rotated). E is similar to A, not congruent – it is smaller.
- If the triangles are congruent, all three angles must be the same in both. You know that two of the angles are 35° and 82° , so $x = 180 - 35 - 82 = 63^\circ$.
- a Identify what values match: SAS (side, angle, side – two sides and the angle between them).
 b Identify what values match: ASA (angle, side angle – two angles and a corresponding side).
- No, they are not congruent. They have the same angles, but the sides may not be the same size (one triangle could be an enlargement of the other).

Constructions


- 1 a and b



- 2 a and b



- 3 a



- b Distance on the diagram = 2.5 cm
 $2.5 \times 100 = 250$ cm in real life = 2.5 m

Drawing circles and parts of circles

- 1 a-d




Diagram not to scale

- 2 a and b










Diagram not to scale

- 3 No, Donald is not correct. A segment of a circle is the area enclosed by a chord and an arc; a sector of a circle is the area enclosed by two radii and the arc between them.

Loci

- 1 a





Perimeter

- A hexagon has 6 sides so perimeter = $6 \times 9 = 54$ cm
- Missing vertical length = $20 - 5 - 5 - 4 = 6$ mm
Missing horizontal lengths are all equal = $25 - 13 = 12$ mm each
Perimeter = $20 + 25 + 6 + 12 + 4 + 12 + 5 + 12 + 5 + 13 = 114$ mm
 $114 \div 10 = 11.4$ cm
- Perimeter of cushion = $\frac{1}{2} \times 2 \times \pi \times 24 + 30 + 48 + 30 = 183$ cm (to nearest cm)
= 1.83 m. So no, Greta does not have enough lace.

Area

- Area = $12 \times 6 = 72$ cm²
 - Area = $\frac{1}{2} \times (3 + 8) \times 4 = 22$ cm²
 - Area of rectangle = $2 \times 10 = 20$ cm²
Area of trapezium = $\frac{1}{2} (a + b)h = \frac{1}{2} (2.5 + 10)10 = 62.5$ cm²
Area of shape = $20 + 62.5 = 82.5$ cm²
- First draw a diagram. Two sides are equal, and are 6 cm. The two other sides are equal, and are x cm.

$$x + x + 6 + 6 = 16$$


$$2x + 12 = 16$$

$$x = \frac{16 - 12}{2} = 2.$$
 Area = $6 \times 2 = 12$ cm²
- Radius of circle = 3.5 cm
Area of circle = $\pi r^2 = \pi \times 3.5^2 = 38.48$ cm²
Area of square = 49 cm²
Area of shaded part = $\frac{49 - 38.48}{4} = 2.63$ cm²

Sectors

- $\frac{120}{360} = \frac{1}{3}$
 - Area = $\frac{1}{3} \times \pi \times 3^2 = 3\pi$ cm²
- Area = $\frac{1}{4} \times \pi \times 2.8^2 = 6.2$ cm² (1 d.p.)
 - Perimeter = $2.8 + 2.8 + \frac{1}{4} \times 2 \times \pi \times 2.8 = 10.0$ cm (1 d.p.)
- Area = $\frac{40}{360} \times \pi \times 5^2 = 8.73$ cm² (2 d.p.)
 - Arc AB = $\frac{40}{360} \times 2 \times \pi \times 5 = 3.49$ cm (2 d.p.)





3D shapes



b triangular prism

c

Number of faces	Number of edges	Number of vertices
5	9	6



3 b The front elevation shows 5 cubes and the side shows that the shape is 2 cubes deep. $5 \times 2 = 10$, so 10 cubes make up the shape.

Volume

1 The front elevation shows 5 cubes and the side shows that the shape is 4 cubes deep. Volume = $5 \times 4 = 20 \text{ cm}^3$

2 a Volume = area of cross-section \times length = $\pi \times 5^2 \times 12 = 942 \text{ cm}^3$ (to 3 s.f.)

b Volume = $\frac{1}{3} \times$ area of cross-section \times length
 = $\frac{1}{3} \times \pi \times 7^2 \times 15 = 770 \text{ cm}^3$ (to nearest cm)

3 Volume = $\frac{1}{2} \times \frac{4}{3} \pi r^3 = \frac{2}{3} \times \pi \times 8^3 = 1072.33 \text{ cm}^3$ (to 2 d.p.)

4 Volume of tank = $40 \times 40 \times 60 = 96\,000 \text{ cm}^3$

Volume of water in tank, 80% full = $0.8 \times 96\,000 = 76\,800 \text{ cm}^3$

Height of water in pond (1st fill) = $76\,800 \div (80 \times 60) = 16 \text{ cm}$

Height of water in pond (2nd fill) = $16 \times 2 = 32 \text{ cm}$

Height of water in pond (3rd fill) = $16 \times 3 = 48 \text{ cm}$

Three tanks of water are needed to fill the pond.

Alternative method: divide volume of pond by volume of water in tank.

$$\frac{80 \times 60 \times 48}{76\,800} = 3$$

Surface area

1 a 6 faces

b Surface area = $60 + 60 + 5 + 5 + 3 + 3 = 136 \text{ cm}^2$

2 surface area = area of triangular side $\times 4$ + area of square base

$$= \left(\frac{1}{2} \times 6 \times 5\right) \times 4 + 6^2$$

$$= (15 \times 4) + 36$$

Surface area = 96 cm^2

3 a Surface area of a sphere = $4\pi r^2 = 4 \times \pi \times 14^2 = 2463.01 \text{ cm}^2$ (to 2 d.p.)

b Surface area of a cone = $\pi r l + \pi r^2 = \pi \times 6 \times 10 + \pi \times 6^2 = 301.59 \text{ cm}^2$ (to 2 d.p.)

4 Area of cylinder = $2\pi r h = 2\pi \times 6 \times 1.5 = 56.55 \text{ cm}^2$

Area of circular base = $\pi r^2 = \pi \times 6^2 = 113.10 \text{ cm}^2$

Area of curved surface area of cone = $\pi r l = \pi \times 6 \times 11.5 = 216.77 \text{ cm}^2$

Total surface area = $56.55 + 113.10 + 216.77 = 386.42 \text{ cm}^2$ (to 2 d.p.)

Using Pythagoras' theorem

$$\begin{aligned} 1 \quad x^2 &= 3^2 + 4^2 \\ &= 9 + 16 \\ &= 25 \\ x &= \sqrt{25} \\ &= 5 \text{ cm} \end{aligned}$$

$$\begin{aligned} 15^2 &= y^2 + 12^2 \\ 225 &= y^2 + 144 \\ 81 &= y^2 \\ y &= \sqrt{81} \\ &= 9 \text{ cm} \end{aligned}$$

$$\begin{aligned} 2 \quad 6^2 &= 4.5^2 + w^2 \\ 36 &= 20.25 + w^2 \\ w &= \sqrt{15.75} \\ &= 3.97 \text{ cm} \end{aligned}$$

Area = $l \times w = 4.5 \times 3.97 = 17.9 \text{ cm}^2$

$$\begin{aligned} 3 \quad AB^2 &= 2^2 + 4^2 \\ &= 4 + 16 \\ &= 20 \\ AB &= \sqrt{20} \\ &= \sqrt{4 \times 5} \\ &= 2\sqrt{5} \text{ units} \end{aligned}$$

4 Square of diagonal of doorway = $70^2 + 190^2 = 41\,000$
 Diagonal of doorway = $\sqrt{41\,000}$
 = $202.48 \text{ cm} = 2.0248 \text{ m} = 2.02 \text{ m}$ (2 d.p.)

Yes, the artwork will fit through the diagonal of the doorway.

Trigonometry

$$1 \quad \tan x = \frac{8}{13} \\ x = 31.6^\circ$$

$$2 \quad \cos 42 = \frac{17}{AC} \\ AC = \frac{17}{\cos 42} = 22.9 \text{ cm}$$

$$3 \quad \sin 49 = \frac{h}{6} \\ h = 6 \sin 49 \\ = 4.53 \text{ m}$$

Exact trigonometric values

$$1 \quad \text{a} \quad \tan x = \frac{\text{opposite}}{\text{adjacent}} = \frac{1}{1} = 1$$

$$\text{b} \quad x = \tan^{-1}(1) = 45^\circ$$

$$2 \quad \cos 30 = \frac{\sqrt{3}}{PR} \\ \frac{\sqrt{3}}{2} = \frac{\sqrt{3}}{PR} \\ PR = 2 \text{ cm}$$



$$3 \quad \frac{YZ}{20} = \sin 30 \\ YZ = 20 \sin 30 \\ = 20 \times \frac{1}{2} \\ = 10 \text{ cm}$$

$$4 \quad \sin 45^\circ = \cos 45^\circ = \frac{1}{\sqrt{2}}$$


So the second angle in the triangle must be 45° .

The third angle will be $180 - 90 - 45 = 45^\circ$.

Transformations



3 Rotation 90° clockwise about (1, -1); or rotation 270° anticlockwise about (1, -1).



c Reflection in $y = 1$

Similar shapes

- 1 a YZW
 b Scale of enlargement = $\frac{\text{enlarged length}}{\text{original length}} = \frac{6.3}{2.1} = 3$
 c $WZ = 4 \times 3 = 12$ cm
 2 a 37.5°
 b Scale of enlargement = $\frac{\text{enlarged length}}{\text{original length}} = \frac{5}{2.5} = 2$
 Length of $AB = 8 \div 2 = 4$ cm
 c They are isosceles, because they have two equal sides and two equal angles. Note that the diagrams are not drawn to scale, as is common practice in maths questions – you have to go by the numbers.
 d Length of $BC = \text{length of } AC = 2.5$ cm
 3 a Scale of enlargement = $\frac{\text{enlarged length}}{\text{original length}} = \frac{4}{6} = \frac{2}{3}$
 b Length of $RT = 4.5 \times \frac{2}{3} = 3$ cm
 4 a Scale of enlargement = $\frac{\text{enlarged length}}{\text{original length}} = \frac{2.2}{4.4} = \frac{1}{2}$
 b Length of $CE = \text{length of } CD \div \frac{1}{2} = 2.8 \times 2 = 5.6$ cm
 c Angle $ACE = 180 - 70 - 64 = 46^\circ$

Vectors

- 1 a $\mathbf{a} = \begin{pmatrix} 3 \\ 2 \end{pmatrix}$ b $\mathbf{b} = \begin{pmatrix} -3 \\ 3 \end{pmatrix}$ c $\mathbf{c} = \begin{pmatrix} -2 \\ -4 \end{pmatrix}$ d $\mathbf{d} = \begin{pmatrix} -4 \\ -2 \end{pmatrix}$
 2 a
-

- b $-\mathbf{p} = \begin{pmatrix} -6 \\ -1 \end{pmatrix}$
 c $2\mathbf{p} = \begin{pmatrix} -12 \\ 2 \end{pmatrix}$
 d $2\mathbf{p} + \mathbf{p} = \begin{pmatrix} -12 \\ 2 \end{pmatrix} + \begin{pmatrix} -6 \\ 1 \end{pmatrix} = \begin{pmatrix} -18 \\ 3 \end{pmatrix}$
 $3\mathbf{p} = 3 \times \begin{pmatrix} -6 \\ 1 \end{pmatrix} = \begin{pmatrix} -18 \\ 3 \end{pmatrix}$
 Therefore, $2\mathbf{p} + \mathbf{p} = 3\mathbf{p}$

- 3 a $\mathbf{a} + \mathbf{b} = \begin{pmatrix} 3 \\ 4 \end{pmatrix} + \begin{pmatrix} -5 \\ -1 \end{pmatrix} = \begin{pmatrix} 3-5 \\ 4-1 \end{pmatrix} = \begin{pmatrix} -2 \\ 3 \end{pmatrix}$
 b $\mathbf{c} = \mathbf{a} - \mathbf{b} = \begin{pmatrix} 3 \\ 4 \end{pmatrix} - \begin{pmatrix} -5 \\ -1 \end{pmatrix} = \begin{pmatrix} 3-(-5) \\ 4-(-1) \end{pmatrix} = \begin{pmatrix} 8 \\ 3 \end{pmatrix}$
 c
-

Probability

Basic probability

- 1 Probability = $\frac{\text{number of successful outcomes}}{\text{total number of possible outcomes}} = \frac{1}{10}$ (or 0.1 or 10%)
 2 $P(\text{not rain}) = 1 - P(\text{rain}) = 1 - 0.6 = 0.4$
 3 a b c
-
- a $P(\text{a number from 1 to 8}) = \frac{8}{8} = 1$
 b $P(\text{a multiple of 3}) = \frac{2}{8} = \frac{1}{4}$
 c $P(\text{a number greater than 3}) = \frac{5}{8}$
 4 $2p - 0.1 + 2p + 0.1 + p = 1$
 $5p = 1$
 $p = 0.2$

Outcome	Red	Blue	Green
Probability	$2p - 0.1$ $= 2 \times 0.2 - 0.1$ $= 0.3$	$2p + 0.1$ $= 2 \times 0.2 + 0.1$ $= 0.5$	$p = 0.2$

Blue is most likely.

Two-way tables and sample space diagrams

- 1 Work out the missing values one by one, for example in the order shown from first to seventh. (There is more than one order you can do it in.)

	Single	Double	King	Totals
Oak	2	Fourth: $42 - 12 =$ $14 = 16$	Fifth: $30 - 16 =$ $2 = 12$	30
Pine	First: $54 - 14$ $- 17 = 23$	14	17	54
Walnut	1	12	Sixth: $32 - 12 =$ $17 = 3$	Seventh: $1 + 12 +$ $3 = 16$
Totals	Second: $2 + 23 +$ $1 = 26$	Third: $100 - 26$ $- 32 = 42$	32	100

2 a

		Spinner			
		1	2	3	4
Coin	Heads	1, H	2, H	3, H	4, H
	Tails	1, T	2, T	3, T	4, T

b $P(1, T) = \frac{1}{8}$
 c $P(2, H) + P(3, H) + P(4, H) = \frac{1}{8} + \frac{1}{8} + \frac{1}{8} = \frac{3}{8}$

3 a


	Study sciences	Do not study sciences	Totals
Boys	$\frac{1}{5} \times 120 = 24$	$45 - 24 = 21$	$\frac{3}{8} \times 120 = 45$
Girls	$75 - 40 = 35$	$\frac{1}{3} \times 120 = 40$	$120 - 45 = 75$
Totals	$24 + 35 = 59$	$21 + 40 = 61$	120

- b 21 boys do not study science, so probability = $\frac{21}{120}$ or $\frac{7}{40}$
 c There are 75 girls and 35 of them study science, so probability = $\frac{35}{75} = \frac{7}{15}$

Sets and Venn diagrams

- 1 a $\xi = \{21, 22, 23, 24, 25, 26, 27, 28, 29\}$
 b $A = \{21, 24, 27\}$
 c $B = \{24, 28\}$
 d $A \cup B = \{21, 24, 27, 28\}$ – A ‘union’ B means **all** the values in A and **all** the values in B
 e $A \cap B = \{24\}$ – A ‘intersect’ B means only those value that are in **both** A and B.

2 a




- b There are 7 adults who only go to art class, so probability = $\frac{7}{20}$
 c 8 adults only go to yoga class and 7 adults only go to art class.
 Probability = $\frac{8+7}{20} = \frac{3}{4}$

- 3 a Total number of respondents = ζ
 $= 25 + 11 + 2 + 12 + 0 + 7 + 3 = 60$
 $P(\text{supermarket only}) = \frac{25}{60} = \frac{5}{12}$
 b $P(F) = \frac{12+2+7+3}{60} = \frac{24}{60} = \frac{2}{5}$
 c $P(L \cap F) = \frac{2+7}{60} = \frac{9}{60} = \frac{3}{20}$
 d $P(S^c) = \frac{0+7+3}{60} = \frac{10}{60} = \frac{1}{6}$


Frequency trees and diagrams

1 a




- b $P(W, B) = \frac{3}{4} \times \frac{1}{4} = \frac{3}{16}$
 c $P(B, B) = \frac{1}{4} \times \frac{1}{4} = \frac{1}{16}$

2 a




b $P(O, O) + P(A, A) = \left(\frac{2}{6} \times \frac{1}{5}\right) + \left(\frac{4}{6} \times \frac{3}{5}\right) = \frac{2}{30} + \frac{12}{30} = \frac{14}{30} = \frac{7}{15}$

3 a



- b One boy chosen at random:
 $P(\text{does not want to go}) = \frac{3}{45} = \frac{1}{15}$

4 a



- b $P(R, R) = \frac{3}{10} \times \frac{1}{2} = \frac{3}{20}$
 c $P(R, R') + P(R', R) = \left(\frac{3}{10} \times \frac{1}{2}\right) + \left(\frac{7}{10} \times \frac{1}{2}\right) = \frac{3}{20} + \frac{7}{20} = \frac{10}{20} = \frac{1}{2}$

The probability of one day having rain and one day having no rain is 50%.

Expected outcomes and experimental probability

- 1 a The spinner was spun $12 + 13 + 10 + 15 = 50$ times.
 b Estimated probability of blue = $\frac{13}{50}$
 c Estimated probability of yellow = $\frac{15}{50} = \frac{3}{10}$
 d You would expect $\frac{10}{15} \times 100 = 20$ green outcomes from 100 spins.
 2 $0.75 \times 20 = 15$ students would be expected to pass the exam.
 3 a Total number of customers = $26 + 20 + 6 + 5 + 3 = 60$
 Estimated probability that someone will buy stamps = $\frac{20}{60} = \frac{1}{3}$
 b $\frac{1}{3} \times 450 = 150$ customers buy stamps each day
 c $\frac{6}{60} \times 450 = 45$ customers buy foreign currency each day
 d $450 \times 6 = 2700$ customers each week
 $\frac{5}{60} \times 2700 = 225$ customers use the post office for banking each week

Statistics

Data and sampling

- 65, because that is 10% of 650 (the entire population).
- $\frac{50}{25000} \times 100 = 0.2\%$. The sample is not big enough.
People in the town centre may not be the only ones using buses. For example, some people may take buses to the local train station, school or hospital.
- $\frac{9}{45} \times 400 = 80$ people like carrot cake
Sam needs to make 80 cakes.
 - Assumptions
Assumed that these are individual carrot cakes. If instead each cake is a large one divided into 8 slices, then only $80 \div 8 = 10$ carrot cakes would be needed.
Assumed the sample is representative of the population; this could affect the answer because not all the 400 people who have accepted the invitation may turn up.

Frequency tables

- $1 + 11 + 9 + 6 + 1 + 2 = 30$ tables
 - $(1 \times 1) + (2 \times 11) + (3 \times 9) + (4 \times 6) + (5 \times 1) + (6 \times 2) = 91$ people

Number of electronic devices	Tally	Frequency
0-1		3
2-3		10
4-5		5
6-7		5
8-9		1


- Continuous

Mass, m (kg)	Tally	Frequency
$50 \leq m < 60$		3
$60 \leq m < 70$		5
$70 \leq m < 80$		4
$80 \leq m < 90$		5
$90 \leq m < 100$		3

- There were 12 people who spent 10 or more minutes, but fewer than 14 minutes, with the GP. However, it could be that none of them spent exactly 10 minutes with the GP.

Bar charts and pictograms

- $9 - 4 = 5$ more boys than girls prefer squash
 - $15 + 6 + 9 + 7 + 4 = 41$ girls were surveyed




- $3 + 3 + 3 + 3 + 2 = 14$ endangered species

- Blue Mountain has 9 endangered species, Salt Springs has 7 endangered species. Blue Mountain has 2 more endangered species than Salt Springs.

Pie charts

- $\frac{150}{360} \times 300 = 125$ people last saw an action movie
 - Romance = 90° , so $\frac{3}{4} \times 300 = 225$ people did not see a romance movie.

	Children	Adults	Senior citizens
Calculation	$\frac{80}{250} \times 360$ $= 115.2$	$\frac{155}{250} \times 360$ $= 223.2$	$\frac{15}{250} \times 360$ $= 21.6$
Angle	115°	223°	22°



- Angle for salad = 72°
 $\frac{360}{72} \times 12 = 60$ children were surveyed
Or:
 $72^\circ = 12$ children
 $\frac{72}{12} = 6^\circ = 1$ child
 $\frac{360}{6} = 60$ children were surveyed

Measures of central tendency: mode

- There are two modes: 3 minutes and 4 minutes, since each appears twice. An alternative correct answer is to say that there is no mode.
- This is the one with the highest frequency: $12 < a \leq 13$.
- This is the number with the biggest slice of the pie: 3.
- The mode is the interest which represents the highest number of students: Sport (boys); Crafts (girls).

Measures of central tendency: median

- Ages in order: 11 11 12 13 13 13 13 (14 15) 15 16 16 17 17 18 18
Median = $14\frac{1}{2}$ years old
- Total frequency = $5 + 12 + 17 + 10 + 6 = 50$
Median = $\frac{50+1}{2} = 25.5$ th person
Median class = $12 < a \leq 13$

Measures of central tendency: mean

- Mean age = $\frac{6+7+11+13+18}{5} = 11$ years old
- Number of bedrooms = $(1 \times 4) + (2 \times 7) + (3 \times 13) + (4 \times 17) = 125$
Number of houses = $4 + 7 + 13 + 17 = 41$
Mean number of bedrooms = $125 \div 41 = 3.05 \approx 3$ bedrooms
- Number of holidays = $(0 \times 4) + (1 \times 21) + (2 \times 9) + (3 \times 2) = 45$
Number of employees = $4 + 21 + 9 + 2 = 36$
Mean number of holidays = $45 \div 36 = 1.25 \approx 1$ holiday

Age of patients, a	Midpoint	Frequency	Midpoint \times frequency
$0 < a \leq 10$	5	3	15
$10 < a \leq 20$	15	18	270
$20 < a \leq 30$	25	6	150
$30 < a \leq 40$	35	11	385
$40 < a \leq 50$	45	10	450
$50 < a \leq 60$	55	19	1045
$60 < a \leq 70$	65	16	1040
$70 < a \leq 80$	75	17	1275
		Total = 100	Total = 4630

Mean age of patients = $\frac{4630}{100} = 46.3 \approx 46$ years old

Range


- Range of boys' ages = $4 - 2 = 2$ years
 - Range of boys' ages = $4 - 1 = 3$ years
 - Range in temperatures for Resort A = $25 - 16 = 9^\circ\text{C}$
 - Range in temperatures for Resort B = $28 - 13 = 15^\circ\text{C}$
 - Business A: Range = $45\,816 - 23\,561 = \pounds 22\,255$
 Mean profit = $\frac{23\,561 + 30\,485 + 39\,210 + 45\,816}{4} = \pounds 34\,768$
 - Business B: Range = $63\,248 - 17\,894 = \pounds 45\,354$
 Mean profit = $\frac{32\,820 + 40\,328 + 17\,894 + 63\,248}{4} = \pounds 38\,572.50$
- c** **Either** Business A because its range in profit is lower and the profit is increasing each year, and so it shows a more consistent performance.
or Business B because its mean profit is higher, and its most recent profit (in Year 4) is $\pounds 17\,432$ more than Business A.

Comparing data using measures of central tendency and range

- Mean time for bus journey = $\frac{32 + 30 + 39 + 32 + 43 + 31}{6} = 34.5$ minutes
 - Mean time for train journey = $\frac{16 + 24 + 18 + 26 + 70 + 17}{6} = 28.5$ minutes
 - Range for bus journey = $43 - 30 = 13$ minutes
 Range for train journey = $70 - 16 = 54$ minutes
 The train journey has the bigger range.
 - Either:** The bus is better because although it takes longer (on average), the range is lower, and so you can predict the time it takes for the journey.
Or: The train is better because it is quicker than the bus (on average), although the range suggests it may be less reliable.
- Mean = 13. This does not represent the age of the people using the playground. In fact, those using the playground are small children (under 10) and their parents (over 25).
 - There are five modes (3, 4, 5, 7, 8), and so the mode does not represent the age of the people using the playground.
 - Ages in order: 3 3 4 4 5 5 7 7 8 8 26 30 33 39
 Median position = $\frac{14 + 1}{2} = 7.5$ th value
 Median age = 7 years old
- Mode = 0; Median = 0; Mean = 2 days. Mode or median are the best averages to use, because the mean is skewed by the student who is absent due to sickness for 24 days.


Time series graphs

1 a and b




- Start at Day 3, read up to the graph line then left to the weight scale: 190 g
- Start at 220 g, read across to the graph line then down: 5 days old
- Mean weight gained = $\frac{\text{total weight gained}}{\text{number of days}} = \frac{270 - 130}{7} = 20$ g per day

2 a



- There is an increase in campers in May. This may be due to May bank holidays, or May half-term, or perhaps there was some very sunny weather.

3 a*




*This answer differs from the one in the Exam Practice Book due to an error in our first edition. This answer has now been re-checked and corrected.

- b** Branch 1 had a steady increase in sales for the first four months. Then sales levelled off to stay at around £25 000. Branch 2 had a slow start to its sales in the first three months. Then perhaps it had a promotion, because sales increased a lot in month 4. Sales have been increasing ever since.


Scatter graphs

- 1 **a** Positive correlation. This means as the temperature rises, more pairs of flip flops are sold.
b Negative correlation. This means as the temperature rises, fewer wellington boots are sold.



- b** The scatter diagram shows a positive correlation between students' maths and physics test percentages. Therefore, the students who got a low percentage in the maths test got the lower percentages in the physics test; the students who got a high percentage in the maths test got the higher percentages in the physics test.

- c** The outlier is the point marked at (0, 58).
d The student was absent for the maths test.



The black line shows the line of best fit.
 The grey line shows the line where a laptop loses £150 every 6 months.
 The shop owner is not correct. The line of best fit shows on average a laptop loses approximately £159/£160 every 6 months.

- b** The line of best fit cannot make a prediction outside the available data. The data only goes as far as 18 months.

Practice papers

Non-calculator

- Digits move one place value to the left:
3.4
- If you rotate C 90° anticlockwise it is an exact reflection of A. It has the same area, proportions, dimensions and shape (in reflection) so it is congruent.
C
 $\frac{10}{14} = \frac{5}{7}$
 $\frac{12}{21} = \frac{4}{7}$ (original fraction cancelled by 3)
 $\frac{16}{28} = \frac{4}{7}$ (original fraction cancelled by 4)
 $\frac{24}{42} = \frac{4}{7}$ (original fraction cancelled by 6)
 $\frac{10}{14}$
- $4x - (2 - 3x)$
 $= 4x - 2 + 3x$
 $= 7x - 2$
- $7.93 \approx 8$
 $12.1 \approx 12$
 $3.86 \approx 4$
 Estimate:
 $\frac{8 + 12}{4}$
 $= \frac{20}{4}$
 $= 5$
- You can cancel the second fraction as follows:
 $\frac{8a}{10a} = \frac{4a}{5a} = \frac{4}{5}$
 So the first two fractions are the same (and are equal to 80%)
8% is not equivalent to the others.
- Its opposite sides are equal (indicated by the pairs of slashed lines across the opposite sides).
 It is a parallelogram (because it has two pairs of equal opposite sides).
Note that:
 - we have no evidence that its vertices are right angles, so we should **not** assume it is a rectangle
 - its diagonals do not bisect at right angles, because its sides are **not** equal, so it is not a rhombus.
- $\frac{3}{8} \times 2400 =$
 $3 \times \frac{2400}{8}$
 $= 3 \times 300$
 $= 900$
- 5:7
 Total number of parts = 12
 7 parts are boys
 $\frac{7}{12}$ are boys
- a** Factor pairs of 36:
 1×36
 2×18
 3×12
 4×9
 6×6
 Multiples of 6:
 6, 12, 18, 24, 30, 36
 Numbers in both groups are:
 6, 12, 18, 36
 From list:
 12 and 18

b $36 = 2 \times 18$
 $= 2 \times 2 \times 9$
 $= 2^2 \times 3^2$

Therefore the only prime factors of 36 are 2 and 3.

c No. A prime number has exactly two factors: 1 and itself. A multiple of 6 would have at least four factors: 1, 2, 3, 6.

11 Let the coins be:

$$2q + r + s + t$$

(Two are the same, represented by $2q$.)

Working in pence:

$$2q + r + s + t = 170$$

Ways of making 70p with 4 coins, with no more than 2 being the same:

$$50p + 10p + 5p + 5p$$

20p + 20p + ? (no way of making 30p without repeating coins)

Without using a 50p or a 20p you can't make 70 out of 4 coins (even if you repeated the largest one, 10p, more than once).

So $q = 5, r = 50, s = 10$

$$(2 \times 5) + 50 + 10 + t = 170$$

$$t = 170 - 10 - 50 - 10 = 170 - 70 = 100$$

So $t = 100p = \text{£}1$

The coins are:

£1, 50p, 10p, 5p, 5p

12 60 minutes in an hour

As a fraction, 25 minutes is:

$$\frac{25}{60} = \frac{5}{12}$$

13 $a + 2b = 6 + 2 \times 4 = 14$ (even)

$$2(a - b) = 2(6 - 4) = 2 \times 2 = 4$$
 (even)

$$\frac{2a}{b} = \frac{2 \times 6}{4} = \frac{12}{4} = 3$$
 (odd)

Therefore $\frac{2a}{b}$ is the expression which gives an odd number.

Or:

$$\text{even} + 2(\text{even}) = \text{even} + \text{even} = \text{even}$$

$$\text{even}(\text{even} - \text{even}) = \text{even} \times \text{even} = \text{even}$$

$$\frac{2 \times \text{even}}{\text{even}} = \frac{\text{even}}{\frac{1}{2} \times \text{even}} = \frac{\text{even}}{\text{even}} \text{ or } \frac{\text{even}}{\text{odd}}$$

so the answer could be either even or odd.

Therefore $\frac{2a}{b}$ is the expression which gives an odd number.

14 Area of rectangle: $4 \times 1.5 = 6\text{m}^2$

Perimeter of triangle = $3 \times 6 = 18\text{m}$

$$BC = 18 - (5.1 + 3.4)$$

$$= 18 - 8.5$$

$$= 9.5\text{m}$$

Note that you cannot use Pythagoras' theorem because BAC is not a right angle.

15 a Ratio 2 : 3

$$16 = 2 \text{ parts}$$

$$1 \text{ part} = 16 \div 2 = 8$$

$$8 \times 3 = 24$$

There are 24 children.

b $16 + 4 = 20$ adults

ratio of adults to children = $20 : 24$

Divide by 4:

$$5 : 6$$

16 The width of the tin is 4 cm and each crayon is 0.8 cm wide.

$$4 \div 0.8 = 5$$

So you could fit the widths of 5 crayons in.

The height of the tin is 1 cm and the height of each crayon is 1 cm.

So you could fit 1 layer of crayons in.

The length of tin is 20 cm and the length of each crayon is 10 cm.

$$20 \div 10 = 2$$

So you could fit 2 rows of crayons in.

Total crayons that will fit in the tin:

$$5 \times 1 \times 2 = 10 \text{ crayons}$$

17 Animals that are not sheep: $(10 + 12 + 18) = 40\%$

$$240 \text{ animals (sheep)} = 60\%$$

$$10\% = 240 \div 6 = 40$$

$$100\% = 10 \times 40 = 400$$

There are 400 animals on the farm.

18 Let price of coffee = x and price of cake = y .

$$2x + y = 4.4 \quad (1)$$

$$x + y = 3.2 \quad (2)$$

Multiply (2) by 2:


$$2x + 2y = 6.4$$

Subtract (1) from this:

$$y = 6.4 - 4.4 = 2$$

The cake was £2.

19 a and b



Reading from graph:

$$y = 2.5$$

$$x = 0.5$$

(Acceptable alternative readings from graph in book would be: 2.4 and 0.4.)

Check answer using the two equations:

$$3 \times 0.5 + 1 = 1.5 + 1 = 2.5$$

$$0.5 + 2 = 2.5$$

20 Let fare = f and number of kilometres = d

In pounds:

$$f = 2.25d + 3.5$$

$$d = \frac{f - 3.5}{2.25}$$

When $f = 12.5$:

$$d = \frac{12.5 - 3.5}{2.25} = \frac{9}{2.25} = 4$$

Abbie travelled 4 km.

21 Ali has not included labels for 'phone call' or 'letter'. This is inconsistent and does not enable someone to know which is which. A better way to do it would be to include a key for the chart.

The 'phone call' and 'letter' angles should not be equal sizes. The phone call angle should be twice as big as the letter angle (40° , and 20° for letter).

22 a i $-2 \leq a < 3$

a is a whole number that can be as low as -2 but is less than 3:

$$-2, -1, 0, 1, 2$$

ii $-5 < b \leq 2$

b is a whole number that is greater than -5 but can be as high as 2:

$$-4, -3, -2, -1, 0, 1, 2$$

b The largest integer in each list is 2:

$$2 \times 2 = 4$$

However, the smallest integers in the lists are $a = -2$ and $b = -4$, and to get ab we multiply them:

$$-2 \times -4 = 8$$

So the largest possible whole number value for $ab = 8$.

23 $1.5 \times 3 \div 10000$
 $= 4.5 \times 10^{-4}$

24 a From the graph:

$$\text{books that are not fiction} = (100 - 60)$$

$$= 40\%$$

Or:

$$\text{Fiction books in library A} = 60\% \times 1500 = \frac{1500 \times 6}{10} = 150 \times 6 = 900$$

$$\text{Books that are not fiction: } 1500 - 900 = 600$$

$$\text{P(not fiction)} = \frac{600}{1500} = \frac{6}{15} = \frac{2}{5} \text{ or } 40\% \text{ or } 0.4$$

b Local history books in library A = $(88 - 60)\% = 28\%$

$$\text{Total local history books in A} = \frac{1500 \times 28}{100} = 15 \times 28 = 1.5 \times 280 = 420$$

Local history books in library B = $(56 - 40)\% = 16\%$

$$\begin{aligned} \text{Total local history books in B} &= \frac{3200 \times 16}{100} \\ &= 32 \times 16 = 30 \times 16 + 2 \times 16 \\ &= 480 + 32 \\ &= 512 \end{aligned}$$

$$512 > 420$$

Library B has more local history books.

25 $3\frac{3}{4} + 1\frac{7}{8}$
 $= \frac{15}{4} + \frac{15}{8}$
 $= \frac{30}{8} + \frac{15}{8}$
 $= \frac{45}{8}$
 $= 5\frac{5}{8}$

26 a $3(x + 2) = 5x - 4$
 $3x + 6 = 5x - 4$
 $6 + 4 = 2x$
 $10 = 2x$
 $x = 5$

b $3x^2 + 27x$
 $= x(3x + 27)$
 $= 3x(x + 9)$

27 $A = \frac{1}{2}bh$
 Base (b) = 6 cm
 Height (h) = $\sqrt{10^2 - 6^2}$
 $= \sqrt{100 - 36}$
 $= \sqrt{64}$
 $= 8 \text{ cm}$
 Therefore, $A = \frac{1}{2} \times 6 \times 8$
 $= 24 \text{ cm}^2$

28 a $5n + 1 = 36$
 $5n = 35$
 $n = 7$

It is the 7th term.

b The value of n needs to be an integer if 36 is in the sequence. Testing them:

$$3n - 1 = 36$$

$$3n = 37$$

$$n = \frac{37}{3} \text{ (not an integer)}$$

$$2n + 3 = 36$$

$$2n = 33$$

$$n = \frac{33}{2} \text{ (not an integer)}$$

$$5n - 8 = 36$$

$$5n = 44$$

$$n = \frac{44}{5} \text{ (not an integer)}$$

$$7n + 1 = 36$$

$$7n = 35$$

$$n = 5$$

$7n + 1$ is a sequence that also includes 36.

29 a $\tan = \frac{\text{opposite}}{\text{adjacent}}$

$$\text{From second triangle, } \tan 55^\circ = \frac{14.3}{10} = 1.43 \text{ (3 s.f.)}$$

b $\sin = \frac{\text{opposite}}{\text{hypotenuse}}$

$$\text{From first triangle, } \sin 55^\circ = \frac{8.19}{10} = 0.819 \text{ (3 s.f.)}$$

30 Area of trapezium = $\frac{1}{2}(a + b)h$

$$\frac{1}{2}(2 + 8) \times 3$$

$$= \frac{10 \times 3}{2}$$

$$= 15 \text{ cm}^2$$

$$\text{Area of semicircle} = \frac{1}{2} \times \pi r^2$$

$$\frac{1}{2} \times \pi \left(\frac{8}{2}\right)^2$$

$$= \frac{1}{2} \times 4^2 \pi$$

$$= 8\pi$$

$$\text{Total area} = 8\pi + 15 \text{ cm}^2$$

Calculator

1 A prime number has exactly 2 factors: 1 and itself.

1 has only 1 factor (1).

$$8 = 1 \times 8 \text{ and } 8 = 2 \times 4$$

$$11 = 1 \times 11$$

$$15 = 1 \times 15 \text{ and } 15 = 3 \times 5$$

Only 11 has exactly 2 factors. 11 is a prime number.

2 All the triangles are right-angled (they are on a grid).

Those that are similar will have the same ratio between their shorter sides (meaning the angles of both triangles are the same).

Ratios of shorter sides are as follows.

A has ratio 3:4

B has ratio 3:6

C has ratio 6:8 = 3:4

D has ratio 4:6

So A and C have the same ratio and are similar.

3 $-10.2 + 7 = -3.2$

4 $11.5 \times 37 = 425.5$

$$£425.50 < £462.50$$

B is the better rate of pay.

5 2 pm is a useful time to work from. The bus departed at (21 - 10) minutes and (25 - 0) seconds before 2 pm (11 minutes, 25 seconds before 2 pm).

In 24-hour clock times, it departed at:

$$= 14:00 \text{ hours} - 11 \text{ minutes} - 25 \text{ seconds}$$


$$= 13.49 \text{ hours} - 25 \text{ seconds}$$

There are 60 seconds in a minute, so:

$$\text{departure} = 13.48 \text{ and } 35 \text{ seconds,}$$

$$\text{or } 1.48 \text{ pm and } 35 \text{ seconds}$$

6 Plot the points you are given on the grid.



To move between points $(-4, 2)$ and $(-2, -2)$ you go 2 spaces along to the right and -4 spaces up (4 spaces down).

Moving in the same way from $(2, 3)$ to the final point gives coordinates:

$$\begin{aligned} & ((2 + 2), (3 - 4)) \\ & = (4, -1) \end{aligned}$$

- 7 a Add all the scores together and divide by the number of scores.

$$\text{mean} = \frac{18 + 16 + 19 + 2 + 18 + 17}{6} = \frac{90}{6} = 15$$

- b Arrange them in ascending order: 2, 16, 17, 18, 18, 19
median is between middle two values: $\frac{17 + 18}{2} = 17.5$

- c The **median** is better because it represents the typical score – the low score of 2 is an outlier and skews the value of the mean.

- 8 $P(\text{snow}) + P(\text{not snow}) = 100\%$

$$\begin{aligned} P(\text{not snow}) &= 100\% - P(\text{snow}) \\ &= (100 - 20)\% \\ &= 80\% \\ &= 0.8 \end{aligned}$$


- 9 a $5x + 2x + 2x = 180^\circ$

$$9x = 180^\circ$$

$$x = 20^\circ$$

$$2x = 40^\circ$$

$$5x = 100^\circ$$



- b The triangle has two equal angles (and two equal sides) and is therefore **isosceles**.

- 10 String used to tie plants:

$$4 \times \frac{3}{8} = \frac{3}{2} \text{ m}$$

Remaining string:

$$4 - \frac{3}{2} = \frac{8}{2} - \frac{3}{2} = \frac{5}{2} \text{ m}$$

String used to lay out flower bed:

$$\frac{5}{2} \times \frac{4}{5} = \frac{5 \times 2}{1 \times 5} = \frac{1 \times 2}{1 \times 1} = 2 \text{ m}$$

$$\text{Total string used} = \frac{3}{2} + 2 = \frac{3}{2} + \frac{4}{2} = \frac{7}{2} \text{ m}$$

$$\text{String left} = 4 - \frac{7}{2} = \frac{8}{2} - \frac{7}{2} = \frac{1}{2} \text{ m}$$

$$\frac{1}{2} \text{ m} = 50 \text{ cm}$$

- 11 (Note that the 10 mark is missing from the vertical axis of the graph in the book.)

- a Profit = sales – costs

$$= 74\,000 - 40\,000$$

$$= 34\,000$$

$$\text{£}34\,000$$

- b Branch 2 sales: 90 000

$$\text{Branch 2 profit: } 90\,000 - 52\,000 = 38\,000$$

$$\text{Branch 3 sales: } 44\,000$$

$$\text{Branch 3 profit: } 44\,000 - 20\,000 = 24\,000$$

$$\text{Double the profit of Branch 3} = 24\,000 \times 2 = 48\,000$$

$$38\,000 < 48\,000$$

The Branch 2 manager is incorrect.

- 12 a $\frac{x}{5} = 20$

$$x = 5 \times 20$$

$$x = 100$$

- b $3a^2 + 2ab$ or $a(3a + 2b)$

- c Divide the cost of the box (p) by the number of tins of beans (t).

$$c = \frac{p}{t}$$

- 13 Bill before VAT:

$$\text{£}70 + (3 \times \text{£}35) = \text{£}175$$

Using multiplier for 20% increase, bill after VAT:

$$\text{£}175 \times 1.2 = \text{£}210$$

- 14 Let Roman's age = r , Freddie's age = f and Claudia's age = c .

$$r + f + c = 33 \quad (1)$$

$$r + f = 27$$

$$\text{So } c + 27 = 33$$

$$c = 6 \quad (2)$$

Roman is 9 years older than Claudia, so:

$$r - 9 = 6$$

$$r = 15 \quad (3)$$

From (1), (2) and (3):

$$15 + f + 6 = 33$$

$$f = 33 - 15 - 6$$

$$f = 12$$

Freddie is 12.

- 15 a $x + y = 360 - 2 \times 129$ (angles in a quadrilateral = 360°)

$$= 360 - 258$$

$$= 102^\circ$$

$$x + y = 102 \quad (1)$$

$$2x = y \quad (2)$$

By substitution in (1): $x + 2x = 102$

$$3x = 102^\circ$$

$$x = 34^\circ$$

$$y = 2 \times 34 = 68^\circ$$

- b It has two equal opposite angles, and two different angles. It has two pairs of equal adjacent sides, and one line of symmetry. So it is a kite.

- 16 Let number of 10p coins be p , number of 20p coins be q , and number of 50p coins be r .

$$r = 4q \text{ (check it's the right way round: } r \text{ will be a bigger number than } q\text{)}$$

$$p = \frac{1}{2}q \text{ (check it's the right way round: } p \text{ will be a smaller number than } q\text{)}$$

$$2p = q$$

$$p = 8, \text{ so eliminate } p:$$

$$2 \times 8 = q$$

$$q = 16$$

From first equation:

$$r = 4 \times 16 = 64$$

$$\text{So } p = 8, q = 16 \text{ and } r = 64.$$

$$\text{Amount in meter, in pence} = (8 \times 10) + (16 \times 20) + (64 \times 50)$$


$$= 80 + 320 + 3200$$

$$= 3600 \text{ pence}$$

$$= \text{£}36$$


- 17 a There is a loose **positive correlation**: there tend to be more deckchairs hired on warmer days.

- b Draw a line of best fit on the graph and read off the value at 20° .



Jeff can expect to hire out approximately 42 deckchairs.
(Depending on how you draw your line of best fit, any answer up to 46 is acceptable.)


- 18 a Number taking just French and German:
 $33 - 16 - 9 - 2 = 6$
 Number taking just German and Spanish:
 $28 - 15 - 2 - 6 = 5$
 Number taking only Spanish:
 $40 - 9 - 2 - 5 = 24$



- b Total = $\xi = 7 + 16 + 9 + 2 + 6 + 15 + 5 + 24 = 84$
 c $2 + 5 = 7$ students take both German and Spanish.
 $P(\text{German} \cap \text{Spanish}) = \frac{7}{84} = \frac{1}{12}$
- 19 The price has been reduced by $\frac{2}{9}$, so the sale price is $\frac{7}{9}$.
 350 represents 7 parts of original price
 $350 \div 7 = 50$
 50 is 1 part = $\frac{1}{9}$ of original price
 Original price:
 $50 \times 9 = 450$
 £450
- 20 Volume for sculpture:
 $\frac{4}{3}\pi 3^3 + \frac{4}{3}\pi 2^3$
 $= \frac{4 \times 27}{3}\pi + \frac{4 \times 8}{3}\pi$
 $= 36\pi + \frac{32}{3}\pi$
 $= 46\frac{2}{3}\pi$
 $= 146.61 \text{ m}^3$ to 2 d.p.
- 21 $\begin{pmatrix} -3 \\ 2 \end{pmatrix} - \begin{pmatrix} -1 \\ -4 \end{pmatrix}$
 $= \begin{pmatrix} -3 + 1 \\ 2 + 4 \end{pmatrix}$
 $= \begin{pmatrix} -2 \\ 6 \end{pmatrix}^*$
- 22 a Write an equation showing the relationship between dollars (d) and pounds (p):
 $d = 1.272p$
 $p = \frac{d}{1.272}$
 Lucy buys \$500:
 $500 = 1.272p$
 $p = \frac{500}{1.272}$
 $= 393.08$ (2 d.p.)
 It costs Lucy £393.08.

- b Lucy brings home $500 - 427 = \$73$
 $p = 0.789d$
 $= 0.789 \times 73$
 $= 57.60$ (2 d.p.)
 Lucy gets back £57.60.

- 23 $\cos = \frac{\text{adjacent}}{\text{hypotenuse}}$
 $\cos 42^\circ = \frac{7}{XZ}$
 $XZ = \frac{7}{\cos 42^\circ}$
 $= 9.42$ cm (2 d.p.)
- 24 x is a negative number because negative \times negative \times negative = negative, or: when multiplying you need an even number of negative signs to make a positive, and with a cube number you are multiplying 3 numbers that are exactly the same. So if the original number is negative, the cube will be negative too.
- 25 a Construct the perpendicular bisector of the line CG.



- b Distance measured on page is 4.1 cm
 1 cm represents 200 m, so:
 4.1 cm represents $200 \times 4.1 = 820$ m
 $= 0.82$ km (0.8 km to 1 d.p.)
- 26 a $4.927^2 + \sqrt[3]{8.135} = 26.2865163054$ (10 d.p. from calculator display)
 $= 26.287$ to 3 d.p.
- b $4.927 \approx 5$
 $5^2 = 25$
 $8.135 \approx 8$
 $\sqrt[3]{8.135} \approx \sqrt[3]{8} = 2$
 $25 + 2 = 27$, so the answer to part a is sensible.
- 27 Depreciation of 18% means each year it has $100 - 18 = 82\%$ of its original value.
 Whole amount = $1 \times 17\,000$
 Multiplier for depreciation = 0.82
 $17\,000 \times 0.82^3 = 9373.256$
 Value after 3 years = £9373 to nearest pound.
- 28 a y -intercept (c) = 2
 gradient (m) = $\frac{-12}{6} = -2$
 Comparing with $y = mx + c$:
 $y = -2x + 2$
- b Parallel line must have the same gradient (m) = -2
 $y = -2x$

*This differs from the question in the Exam Practice book due to an error in our first edition. This question has now been re-checked and corrected.